

Supplementary Table 1: Conidia production (conidia per mL) of 18 isolates of *Mycogone pernicioso* growing on PDA medium at 25°C for 7 days.

No	Isolates	Concentration of conidia production Conidia/ mL (10 ⁶)
1	Hp1	0.003 ±0.001 ^a
2	Hp2	0.000
3	Hp3	0.002 ±0.001 ^a
4	Hp4	0.006 ±0.002 ^a
5	Hp5	0.067 ±0.029 ^a
6	Hp6	0.233 ±0.029 ^a
7	Hp7	0.467 ±0.029 ^a
8	Hp8	0.000
9	Hp9	0.037 ±0.046 ^{ab}
10	Hp10	0.517 ±0.076 ^a
11	Hp11	0.317 ±0.029 ^a
12	Hp12	0.283 ±0.029 ^a
13	Hp13	0.02 ±0.003 ^a
14	Hp14	0.03 ±0.004 ^a
15	Hp15	0.067 ±0.029 ^a
16	Hp16	0.067 ±0.058 ^a
17	Hp17	0.10 ±0.050 ^a
18	Hp18	0.60 ±0.180 ^a

Results followed by the same letter in the column do not differ significantly (Duncan's Multiple Range Test, P < 0.05). *P = 0.0001