

Supplementary Materials

Stabilization of Hypoxia-Inducible Factors and BNIP3 Promoter Methylation Contribute to Acquired Sorafenib Resistance in Human Hepatocarcinoma Cells

Carolina Méndez-Blanco, Flavia Fondevila, Paula Fernández-Palanca, Andrés García-Palomo, Jos van Pelt, Chris Verslype, Javier González-Gallego and José L. Mauriz




Figure S1. Full-length immunoblot images from Figures 2, 3 and 4: (a) Complete Western blot images of HIF-1 α , HIF-2 α , β -actin and PCNA from Figure 2 are represented. HIF-2 α and PCNA were reincubated after stripping in the blots of HIF-1 α and β -actin from Figure 2a, respectively; (b) Full Western blot images of Bax, cleaved caspase-3 and PCNA from Figure 3b and immunoblots of HIF-1 α , HIF-2 α and PCNA from Figure 3c; (c) Total BNIP3 and PCNA immunoblots from Figure 4. PCNA were reincubated after stripping of BNIP3 blots. PVDF membranes from (a) and (b) were cut to incubate with different antibodies simultaneously.



© 2019 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).