

Article title: QTL mapping and successful introgression of the spring wheat derived QTL *Fhb1* for Fusarium head blight resistance in three European triticale populations

Journal: Theoretical Applied Genetics

Authors: Ollier Marine^{1,2,3,6}, Talle Vincent¹, Brisset Anne-Laure¹, Le Bihan Zoé¹, Duerr Simon^{1,5}, Lemmens Marc¹, Goudemand Ellen³, Robert Olivier^{3,4}, Hilbert Jean-Louis², Buerstmayr Hermann¹

1. BOKU-University of Natural Resources and Life Sciences Vienna, Department of Agrobiotechnology, IFA-Tulln, Institute of Biotechnology in Plant Production, Konrad Lorenz Str. 20, 3430 Tulln, Austria

2. EA 7394, USC INRA 1411, Institut Charles Viollette (ICV), Agro-food and biotechnology research institute, Université de Lille, INRA, ISA, Univ. Artois, Univ. Littoral Côte d'Opale, Cité Scientifique, 59655 Villeneuve d'Ascq, France

3. Florimond-Desprez Veuve & Fils SAS, 3 rue Florimond-Desprez, BP 41, 59242 Cappelle-en-Pévèle, France

4. Deceased on February 21, 2017

5. Current address: Saatzucht Donau GmbH & Co KG, Breeding Station Reichersberg, Austria.

6. Current address: Bayer Crop Science, Le petit Boissay, Toury, France.

Corresponding author: Marine Ollier, marine.ollier@bayer.com

ESM_1 : Overview of the climatic conditions observed at IFA-Tulln (Austria) for the years 2014 to 2017, with the level of precipitation (mm) draw in solid line, and the average temperature (°C) draw in dashed line.

