

Table S2. Demographics and Clinical Characteristics of 141 cases of CRC

Characteristic	patients						P value
			Non-hypertension		Hypertension		
	n=141	%	n=114	%	n=27	%	
Sex							
male	85	60.3	68	59.6	17	63.0	0.752
female	56	39.7	46	40.4	10	37.0	
Age (years)							
≤ 60	93	34.0	80	70.2	13	51.9	0.030
>60	48	66.0	34	29.8	14	48.1	
BMI index							
<18.5	16	11.3	15	13.2	1	3.7	0.338
18.5-24.9	107	75.9	84	73.7	23	85.2	
≥25	18	12.8	15	13.2	3	11.1	
Tumour site							
left colon	47	33.3	37	32.5	10	37.0	0.528
right colon	39	27.7	30	26.3	9	33.3	
rectum	55	39.0	47	41.2	8	29.6	
Tumour differentiation							
Poorly	16	11.3	13	11.4	3	11.1	0.827
Moderately	88	62.4	70	61.4	18	66.7	
Well	19	13.5	15	13.2	4	14.8	
-	18	12.8	16	14.0	2	7.4	
TNM stage							
I	1	0.7	1	0.9	0	0.0	0.926
II	17	12.1	13	11.4	4	14.8	
III	22	15.6	18	15.8	4	14.8	
IV	101	71.6	82	71.9	19	70.4	
Lymph node metastasis							
positive	72	51.1	58	50.9	14	51.9	0.927
negative	69	48.9	56	49.1	13	48.1	
Liver metastasis							
positive	68	48.2	56	49.1	12	44.4	0.662
negative	73	51.8	58	50.9	15	55.6	
Lung metastasis							
positive	39	27.7	30	26.3	9	33.3	0.464

negative	102	72.3	84	73.7	18	66.7	
Organ metastasis(Others)							
positive	23	16.3	19	16.7	4	14.8	0.815
negative	118	83.7	95	83.3	23	85.2	
Gene mutation							
KRAS	22	15.6	17	14.9	5	18.5	0.100
BRAF	3	2.1	1	0.9	2	7.4	
other	2	1.4	1	0.9	1	3.7	
-	114	80.9	95	83.3	19	70.4	

Note. The correlation was examined using Chi-square test.