

## Supplementary Data

SUPPLEMENTARY TABLE S1.  
DEMOGRAPHIC CHARACTERISTICS

Patients ( <i>n</i> )	261
Age at diagnosis (mean ± SD), years	9.6 ± 4.0
Age group at diagnosis, <i>n</i> (%), years	
≤6	59 (22.6)
6 to <13	154 (59.0)
13 to <18	48 (18.4)
Sex, <i>n</i> (%)	
Female	131 (50.2)
Male	130 (49.8)
Race, <i>n</i> (%)	
NHW	114 (43.7)
NHB	4 (1.5)
Hispanic	64 (24.5)
Asian and Pacific Islander	29 (11.1)
AIAN	1 (0.4)
Unknown	49 (18.8)
Insurance type, <i>n</i> (%)	
Private	194 (74.3)
Public	65 (24.9)
Unknown	2 (0.8)
First A1c (mean ± SD), %	10.8 ± 2.5
A1c at 0 months (mean ± SD)	11.4 ± 2.2
A1c at 18 months (mean ± SD)	8.2 ± 1.9
Max A1c post nadir (mean ± SD)	8.4 ± 1.8
Time to Max A1c post nadir in months (mean ± SD)	13.3 ± 3.6
Pump use within 18 month post diagnosis, <i>n</i> (%)	
Yes	105 (40.2)
No	156 (59.8)
CGM use within 18 months post diagnosis, <i>n</i> (%)	
Yes	108 (41.4)
No	153 (58.6)

AIAN, American Indian or Alaska Native; CGM, continuous glucose monitor; NHB, non-Hispanic black; NHW, non-Hispanic white.