

Table S2. Effects of ticagrelor combined with aspirin and ticagrelor combined with placebo on coagulation activation evaluated with f1.2 and d-Dimer in venous blood from baseline to 24 h and from 2 h to 24 h in 44 healthy male volunteers

	f1.2 (nmol L⁻¹)		d-Dimer (mcg mL⁻¹)	
	Ticagrelor + aspirin	Ticagrelor + placebo	Ticagrelor + aspirin	Ticagrelor + placebo
Baseline	0.094 (0.084; 0.126)	0.095 (0.079; 0.159)	0.104 (0.085; 0.166)	0.142 (0.089; 0.230)
2 h	0.080 (0.076; 0.116)	0.091 (0.070; 0.141)	0.098 (0.080; 0.142)	0.138 (0.089; 0.213)
24 h	0.091 (0.078; 0.114)	0.091 (0.071; 0.149)	0.109 (0.082; 0.150)	0.130 (0.085; 0.190)
p-Value ^a	0.30	0.36	0.13	<0.0001
p-Value ^b	0.17		0.046	
GMR ^a	1.11 (0.91; 1.36)	0.91 (0.74; 1.12)	0.95 (0.90; 1.01)	0.88 (0.82; 0.93)
GMR ^b	1.22		1.09	

	(0.92; 1.63)		(1.00; 1.19)	
p-Value ^c	0.07	0.70	0.04	0.27
p-Value ^d	0.31		0.02	
GMR ^c	1.21 (0.98; 1.48)	1.04 (0.85; 1.28)	1.08 (1.01; 1.15)	0.96 (0.90; 1.03)
GMR ^d	1.16 (0.87; 1.55)		1.12 (1.02; 1.23)	

^aFrom baseline to 24 h. ^bDifferences between treatment groups in the interval from baseline to 24 h. ^cFrom 2 h to 24 h. ^dDifferences between treatment groups in the interval from 2 h to 24 h. Values are given as median (quartiles) and geometric mean ratio (GMR) (95% confidence interval) for prothrombin fragment 1.2 (f1.2) and d-Dimer, respectively.