

Supplementary Table 1: Results for trials where only end of treatment depression scores were available

	n			Variance Ratio Meta Analysis			r (M, SD)		Coefficient of Variance Ratio Meta-Analysis		
	Trials	AD	Placebo	VR (95% CI)	p	Q(df)	AD	Placebo	CVR (95% CI)	P	Q (df)
AD all	84	10879	7346	0.98 (0.96-1.00)	.07	52.53 (83)	.36**	.54**	1.15 (1.11-1.18)	.00	92.67 (83)
SSRI	40	5424	3763	0.99(0.96-1.03)	.71	28.60 (39)	.63**	.37*	1.13 (1.08-1.17)	.00	35.23 (39)**
SNRI	13	1440	1288	0.99 (0.94-1.05)	.73	4.50 (12)	-.49	.62*	1.15 (1.07-1.24)	.00	9.82 (12)**
Atypical	31	4245	3092	0.97 (0.94-1.00)	.07	12.63 (30)	.43*	.14	1.12 (1.07-1.17)	.00	34.37 (30)
Tricyclics	13	670	643	0.93 (0.86-1.01)	.07	9.18(10)	.09	.72**	1.37 (1.20-1.56)	.00	15.64 (12)

Notes:

p (M, SD) is the pearson correlation of mean pre-post differences in depression (M) and the respective standard deviation (SD). Q(df) ist the heterogeneity index for the meta-analysis. * p < .05, ** p < .01.