

Appendix 2 – Drop-out analysis	No questioner data (n=432)		Questioner data (n=1288)		Received questioner (n=1720)		Fisher's exact test
<b>Age, n (%)</b>							
≤50	17	(3.9)	39	(3.0)	56	(3.3)	0.003
51-60	170	(39.4)	401	(31.1)	571	(33.2)	
61-65	128	(29.6)	488	(37.9)	616	(35.8)	
66-70	117	(27.1)	360	(28.0)	477	(27.7)	
<b>T-stage, n (%)</b>							
T1a	25	(5.8)	45	(3.5)	70	(4.1)	0.04
T1b	6	(1.4)	8	(0.6)	14	(0.8)	
T1c	320	(74.1)	953	(74.0)	1273	(74.0)	
T2	81	(18.8)	282	(21.9)	363	(21.1)	
<b>PSA, n (%)</b>							
≤3	48	(11.1)	72	(5.6)	120	(7.0)	<0.001
3.1-7	292	(67.6)	922	(71.6)	1214	(70.6)	
7.1-10	92	(21.3)	294	(22.8)	386	(22.4)	
<b>Proportion positive cores, n (%)</b>							
≤12.5%	137	(31.7)	378	(29.3)	515	(29.9)	0.15
12.5-25%	107	(24.8)	380	(29.5)	487	(28.3)	
25.1-50%	126	(29.2)	359	(27.9)	485	(28.2)	
>50%	27	(6.2)	111	(8.6)	138	(8.0)	
Missing data	35	(8.1)	60	(4.7)	95	(5.5)	
<b>Mode of detection, n (%)</b>							
Screening	203	(47.0)	709	(55.0)	912	(53.0)	0.024
LUTS	145	(33.6)	377	(29.3)	522	(30.3)	
Other symptoms	64	(14.8)	160	(12.4)	224	(13.0)	
Missing data	20	(4.6)	42	(3.3)	62	(3.6)	
<b>Treatment according to NPCR, n (%)</b>							
AS	202	(46.8)	476	(37.0)	678	(39.4)	0.002
RP	187	(43.3)	660	(51.2)	847	(49.2)	
RT	43	(10.0)	152	(11.8)	195	(11.3)	