

Microaerobic conditions caused the overwhelming dominance of *Acinetobacter* spp. and the marginalization of *Rhodococcus* spp. in diesel fuel/crude oil mixture amended enrichment cultures

Fruzsina Révész^{1,2}, Perla Abigail Figueroa-Gonzalez³, Alexander J. Probst³, Balázs Kriszt^{1,2}, Sinchan Banerjee¹, Sándor Szoboszlay², Gergely Maróti⁴, András Táncsics^{1,2,*}

¹Regional University Center of Excellence in Environmental Industry, Szent István University, Gödöllő, Hungary

²Department of Environmental Safety and Ecotoxicology, Szent István University, Gödöllő, Hungary


³Biofilm Centre, University of Duisburg-Essen, 45141, Essen, Germany

⁴Institute of Plant Biology, Biological, Research Centre of the Hungarian Academy of Sciences, Szeged, Hungary


Journal: Archives of Microbiology

*Corresponding author: András Táncsics, Szent István University, Páter K. u. 1., 2100 Gödöllő, Hungary, tancsics.andras@fh.szie.hu, tel.: 06 28 522 000 #1611

(a)



(b)



Supplementary Figure 1: Cluster analysis of the 16S rDNA-based T-RFLP electropherograms by (a) Jaccard and, (b) Bray-Curtis algorithm.