

Supplementary

# Clinical and Radiographic Evaluation of Nanohydroxyapatite Powder in Combination with Polylactic Acid/Polyglycolic Acid Copolymer as Bone Replacement Graft in the Surgical Treatment of Intra-bony Periodontal Defects: A Retrospective Case Series Study

Simone Verardi <sup>1</sup>, Teresa Lombardi <sup>2</sup> and Claudio Stacchi <sup>3,\*</sup>

**Table S1.** Distribution of treated sites and clinical outcomes.

Patient No.	Tooth	PD			CAL		
		Baseline (mm)	12-month (mm)	PD reduction (mm)	Baseline (mm)	12-month (mm)	CAL gain (mm)
1	43	11	4	7	12	7	5
2	32	9	5	4	10	7	3
3	44	7	4	3	9	7	2
4	31	9	4	5	12	10	2
5	11	12	4	8	14	11	3
6	12	8	2	6	10	5	5
7	11	8	4	4	10	7	3
8	21	11	4	7	13	8	5
9	46	8	4	4	11	7	4
10	44	8	3	5	9	4	5
11	36	9	4	5	11	5	6
12	46	8	5	3	10	6	4
13	35	7	3	4	8	4	4
14	14	7	3	4	9	6	3
15	15	7	4	3	8	5	3
16	35	7	4	3	11	6	5
17	15	8	4	4	8	6	2
18	26	10	6	4	12	7	5
19	46	8	5	3	9	6	3
20	16	9	4	5	10	5	5
21	36	8	4	4	9	6	3
22	43	8	5	3	10	7	3
23	15	7	5	2	8	6	2
24	43	7	4	3	8	4	4
25	16	7	3	4	8	4	4
<b>Mean</b>		8.32	4.04	4.28	9.96	6.24	3.72
<b>SD (±)</b>		1.41	0.84	1.46	1.69	1.71	1.17

PD: probing depth; CAL: clinical attachment level; No.: number; mm: millimeters; SD: standard deviation.



© 2020 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).