

Supplementary Material

Exploring deep learning for complex trait genomic prediction in polyploid outcrossing species

L.M. Zingaretti^{1*}, S.A. Gezan², L.F. Ferrão³, L.F. Osorio⁴, A. Monfort^{1,5}, P.R. Muñoz³, V.M. Whitaker⁴, M. Pérez-Enciso^{1,6*}

¹ Centre for Research in Agricultural Genomics (CRAG), CSIC-IRTA-UAB-UB Consortium, 08193 Bellaterra, Barcelona, Spain

² School of Forest Resources and Conservation, University of Florida, 363 Newins-Ziegler Hall, PO Box 110410, Gainesville, FL 32611-0410, USA

³ Blueberry Breeding and Genomics Lab, Horticultural Sciences Department, University of Florida, Gainesville, FL 32611, USA

⁴ IFAS Gulf Coast Research and Education Center, University of Florida, 14625 CR 672, Wimauma, FL 33598, USA

⁵ Institut de Recerca i Tecnologia Agroalimentàries (IRTA), 08193 Barcelona, Spain

⁶ ICREA, Passeig de Lluís Companys 23, 08010 Barcelona, Spain

*** Correspondence:**

Corresponding authors:

laura.zingaretti@cragenomica.es

miguel.perez@uab.es

Figure S1: a) Principal component (PC) representation of the strawberry genotypes, each season (T2, T4, T6, T8 and T10) is in a different color; b) Principal component representation of the blueberry genotypes, training and validation individuals are colored differently.

