

Supplementary table 2: Overall sociodemographic and clinical characteristics of 7,989 CaPSURE participants with PCa and univariate analysis comparing characteristics between CAM users versus non-users.

<b>Characteristics</b>	<b>All Participants:</b>	<b>CAM non-user:</b>	<b>CAM user:</b>	<b>P-value</b>
<b>All participants n (%)</b>	7989 (100)	3484 (44)	4505 (56)	
<b>Age at diagnosis (mean ± SD)</b>	66±9	66±9	65±8	<0.01
<b>Race/ethnicity n (%)</b>				
White	7226 (90)	3022 (87)	4204 (93)	<0.01
Black	551 (7)	355 (10)	196 (4)	
Latino	90 (1)	50 (1)	40 (1)	
Other	122 (2)	57 (2)	65 (1)	
<b>Education level n (%)</b>				
High school	2605 (33)	1421 (41)	1184 (26)	<0.01
College	4373 (55)	1505 (43)	2868 (64)	
Unknown	1011 (13)	558 (16)	453 (10)	
<b>Income category in \$/year n (%)</b>				
<30,000	1676 (21)	909 (26)	767 (17)	<0.01
30,000 to <50,000	1451 (18)	575 (17)	876 (19)	
50,000 to <75,000	1266 (16)	480 (14)	786 (17)	
≥75,000	1930 (24)	613 (18)	1317 (29)	
Unknown	1666 (21)	907 (26)	759 (17)	
<b>Insurance type n (%)</b>				
Medicare	7351 (92)	3184 (91)	4167 (93)	0.18
Private	261 (3)	125 (4)	136 (3)	
Other/unknown	377 (5)	175 (5)	202 (5)	
<b>Study site n (%)</b>				
Academic	628 (8)	189 (5)	439 (10)	<0.01
Community	7112 (89)	3175 (91)	3937 (87)	
Veterans	249 (3)	120 (3)	129 (3)	
<b>Study region n (%)</b>				
West	1277 (16)	456 (13)	821 (18)	<0.01
East	3298 (41)	1516 (44)	1782 (40)	
Midwest	2247 (28)	915 (26)	1332 (30)	
South	1167 (15)	597 (17)	570 (13)	
<b>Gleason grade at diagnostic biopsy n (%)</b>				
2 to 6	4792 (60)	2083 (60)	2709 (60)	0.12
7 (3+4)	1430 (18)	594 (17)	836 (19)	
7 (4+3) or 8 to 10	1565 (20)	718 (21)	847 (19)	
Missing	202 (3)	89 (3)	113 (3)	
<b>Clinical T-stage</b>				
T1	4196 (53)	1752 (50)	2444 (54)	<0.01
T2	3096 (39)	1418 (41)	1678 (37)	
T3	194 (2)	105 (3)	89 (2)	
T4	13 (0.2)	8 (0.2)	5 (0.1)	
Missing	490 (6)	201 (6)	289 (6)	
<b>PSA at diagnosis in ng/ml</b>	6.0 (4.5 - 9.0)	6.2 (4.7 – 10.1)	5.8 (4.4 – 8.4)	<0.01

<b>Characteristics</b>	<b>All Participants:</b>	<b>CAM non-user:</b>	<b>CAM user:</b>	<b>P-value</b>
<b>median (SD)</b>				
<b>BMI at diagnosis in kg/m<sup>2</sup> median (IQR)</b>	26.9 (24.7 - 29.7)	27.0 (24.8 - 29.7)	26.7 (24.7 - 29.6)	0.21
<b>Comorbidities at diagnosis n ± SD</b>	1.9 ± 1.4	1.8 ± 1.4	2.0 ± 1.5	<0.01
<b>Primary treatment n (%)</b>				
Radical prostatectomy	4036 (51)	1665 (48)	2371 (53)	<0.01
Cryotherapy	275 (3)	111 (3)	164 (4)	
Brachytherapy	1090 (14)	486 (14)	604 (13)	
Radiation therapy	650 (8)	309 (9)	341 (8)	
Androgen deprivation therapy	935 (12)	488 (14)	447 (10)	
Chemotherapy	8 (0.1)	3 (0.1)	5 (0.1)	
Non-invasive management	995 (13)	422 (12)	573 (13)	
<b>Study participation (i.e., time from diagnosis to final questionnaire) in years median (IQR)</b>	4.3 (1.8 – 9.8)	2.2 (1.1 – 4.5)	7.0 (3.5 – 12.0)	<0.01