

Supplemental Figure

Deficiency of IgE protects mice from experimental abdominal aortic aneurysms

Jie Li,^{*†} Zhiyong Deng,^{†‡} Xian Zhang,[†] Feng Liu,^{*} Chongzhe Yang,^{*†‡} Guo-Ping Shi[†]

^{*}Department of Geriatrics, National Key Clinic Specialty, Guangzhou First People's Hospital, School of Medicine, South China University of Technology, Guangzhou, Guangzhou, China

[‡]Department of Geriatrics, Guangzhou First People's Hospital, Guangzhou Medical University, Guangzhou, China

[†]Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA 02115, USA




Figure S1. FACS analyses of CD11b⁺Ly6G⁺ neutrophil (A), CD11b⁺F4/80⁺ macrophage (B), CD11b⁺Ly6C^{hi}, CD11b⁺Ly6C^{int}, and CD11b⁺Ly6C^{lo} monocyte (C), CD4⁺ and CD8⁺ T cells (D), and CD117⁺FcεR1⁺ mast cells (E) in spleens from 8 weeks old *Apoe*^{-/-} and *Apoe*^{-/-}*Ige*^{-/-} mice. Data are mean±SEM from three independent experiments.