

Calcium-Induced Activity and Folding of a Repeat in Toxin Lipase from Antarctic *Pseudomonas fluorescens* Strain AMS8

Nur Shidaa Mohd Ali ^{1,2}, Abu Bakar Salleh ¹, Raja Noor Zaliha Raja Abd Rahman ^{1,3}, Thean Chor Leow ^{1,4} and Mohd Shukuri Mohamad Ali ^{1,2,*}

¹ Enzyme and Microbial Technology Research Center, Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia; nur_shidaa@yahoo.com (N.S.M.A.); abubakar@upm.edu.my (A.B.S.); rnzaliha@upm.edu.my (R.N.Z.R.A.R.); adamleow@upm.edu.my (T.C.L.)

² Department of Biochemistry, Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia.

³ Department of Microbiology, Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia.

⁴ Department of Cell and Molecular Biology, Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia.

* Correspondence: mshukuri@upm.edu.my; Tel.: +603-9769 6721




Figure S1: The predicted structure of AMS8 lipase obtained from Ali et al. (2013).