

Bayesian Spatiotemporal Modeling of Routinely Collected Data to Assess the Effect of Health Programs in Malaria Incidence During Pregnancy in Burkina Faso

Toussaint Rouamba^{1,2}, Sekou Samadoulougou^{3,4}, Halidou Tinto¹, Victor A. Alegana^{5,6}, and Fati Kirakoya-Samadoulougou²

1 Clinical Research Unit of Nanoro, Institute for Research in Health Sciences, National Center for Scientific and Technological Research, 528, Avenue Kumda-Yoore, BP 218 Ouagadougou CMS 11, Ouagadougou, Burkina Faso

2 Center for research in epidemiology, Biostatistics and Clinical Research, School of Public Health, University libre de Bruxelles (ULB), Route de Lennik, 808 B-1070 Bruxelles. Brussels, Belgium

3 Centre for Research on Planning and Development (CRAD), Laval University, Quebec, G1V 0A6, Canada

4 Evaluation Platform on Obesity Prevention, Quebec Heart and Lung Institute, Quebec, G1V 4G5, Canada.

5. Kenya Medical Research Institute - Wellcome Trust Research Programme, P.O. Box 43640-00100, Nairobi, Kenya

6. Geography and Environmental Science, University of Southampton, SO17 1BJ Southampton, UK

*Correspondence to rouambatoussaint@gmail.com

Supplementary Information 6

Figure S6. Posterior distributions of the regression parameters on antilog scale. The left panel contains trace-plots while the right panel are density estimates.

