

Supplementary Materials

Table S1. Quantification of salicylic acid production by endophytic bacteria isolated from various plant species based on HPLC method.

Plant species	Isolates	Isolates Identified as	Tissues	SA production (µg/mL)
<i>Taxus brevifolia</i>	EB-2	<i>Micrococcus luteus</i>	Fruit	NM*
	EB-7	<i>Enhydrobacter aerosaccus</i>	Leaves	NM
	EB-17	<i>Brachybacterium rhamnosum</i>	Leaves	NM
<i>Salix babylonica</i>	EB-42	<i>Pseudomonas syringae</i> pv. <i>phaseolicola</i>	Leaves	NM
	EB-43	<i>Xanthomonas campestris</i> pv. <i>campestris</i>	Leaves	NM
	EB-44	<i>Pseudomonas tremiae</i>	Leaves	57.05
	EB-45	<i>Xanthomonas campestris</i> pv. <i>campestris</i>	Leaves	NM
	EB-46	<i>Pantoea vagans</i>	Leaves	NM
	EB-47	<i>Curtobacterium herbarum</i>	Leaves	46.22
	Eb-48	<i>Curtobacterium plantarum</i>	Leaves	NM
	EB-49	<i>Agrobacterium tumefaciens</i>	Leaves	NM
	EB-50	<i>Xanthomonas gardneri</i>	Leaves	44.24
	EB-51	<i>Bacillus altitudinis</i>	Leaves	NM
	Eb-53	<i>Pseudomonas rhizosphaerae</i>	Leaves	NM
	EB-55	<i>Curtobacterium citreum</i>	Leaves	NM
	EB-56	<i>Agrobacterium tumefaciens</i>	Leaves	NM
	EB-57	<i>Curtobacterium flaccumfaciens</i> pv. <i>flaccumfaciens</i>	Leaves	NM
	EB-58	<i>Curtobacterium plantarum</i>	Leaves	NM
	EB-59	<i>Pseudomonas psychrotolerans</i>	Leaves	NM
	EB-60	<i>Curtobacterium flaccumfaciens</i> pv. <i>flaccumfaciens</i>	Leaves	NM
<i>Metasequoia glyptostroboides</i>	EB-62	<i>Agrobacterium larrymoorei</i>	Leaves	NM
	EB-63	<i>Enhydrobacter aerosaccus</i>	Leaves	NM
	EB-65	<i>Brevundimonas vesicularis</i>	Leaves	NM
<i>Ginkgo biloba</i>	EB-83	<i>Pseudomonas syringae</i> pv. <i>phaseolicola</i>	Cones	NM
	EB-84	<i>Micrococcus luteus</i>	Cones	NM
	EB-87	<i>Staphylococcus hominis</i> subsp. <i>novobiophageicus</i>	Leaves	NM
	EB-124	<i>Bacillus subtilis</i> subsp. <i>subtilis</i>	Fruits	NM

*NM-Not measurable amount of salicylic acid produced by endophytic bacteria, because minimum peak height was less than 0.1mV.