

New Eco-Friendly Synthesized Thermosets from Isoeugenol-Based Epoxy Resins

Quentin Ruiz ^{1,2}, Sylvie Pourchet ^{1,*}, Vincent Placet ², Laurent Plasseraud ¹ and Gilles Boni ^{1,*}

¹ ICMUB Institute, Université de Bourgogne Franche-Comté, UMR 6302 CNRS-UB, F-21000 Dijon, France; quentin_ruiz@etu.u-bourgogne.fr (Q.R.); laurent.plasseraud@u-bourgogne.fr (L.P.)

² FEMTO-ST Institute, Université de Bourgogne Franche-Comté, UMR 6174 CNRS-UFC-ENSMM-UTBM, Department of Applied Mechanics, F-25000 Besancon, France; vincent.placet@univ-fcomte.fr

* Correspondence: sylvie.pourchet@u-bourgogne.fr (S.P.); gilles.boni@u-bourgogne.fr (G.B.)

Figure S1. ¹H NMR spectrum of BioIgenol (CDCl₃)

Figure S2. ¹³C{¹H} NMR spectrum of BioIgenol (CDCl₃)

Figure S3. ¹H NMR Spectrum of DiEP1 and DiEP2 (CDCl₃)

Figure S4. ¹³C{¹H} NMR Spectrum of DiEP1 and DiEP2 (CDCl₃)

Figure S5. ¹H NMR spectrum of BioIgenox (CDCl₃)

Figure S6. ¹³C{¹H} NMR spectrum of BioIgenox (CDCl₃)




Figure S1. ^1H NMR spectrum of BioIgenol (CDCl_3)




Figure S2. $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum of BioIgenol (CDCl_3)




Figure S3. ^1H NMR Spectrum of DiEP1 and DiEP2 (CDCl_3)




Figure S4. $^{13}\text{C}\{^1\text{H}\}$ NMR Spectrum of DiEP1 and DiEP2 (CDCl_3)




Figure S5. ¹H Spectrum of BioIgenox (CDCl₃)




Figure S6. ¹³C{¹H} spectrum of BioIgenox (CDCl₃)