

Figure S1. Summary of cocultures with permissible Raji and Sup T cell lines and SRV antibody reactive, PCR negative donor PBMCs.

Day	Manipulations/Observations	PCR
-2	Stimulate PBMC with Staph Entertoxin A	Donors 1-6 negative Donors 1-6 Raji negative Donors 1-6 SupT
0	Coculture PBMC with Raji and Sup T	negative
4	No CPE. Add one half volume fresh media. Clumping but no CPE.	Not tested Donors 1-6 Raji negative Donors 1-6 SupT
7	Add fresh cells to double volume. Clumping but no CPE	negative Donors 1-6 Raji negative Donors 1-6 SupT
11	Replace one half volume with fresh media.	negative Donors 1-6 Raji negative Donors 1-6 SupT
14	No CPE but enlarged cells only in donor #4/ Sup T. Add one fourth volume fresh media.	negative Donors 1-6 Raji negative Donors 1-6 SupT
18	No CPE. Clumps and debris in Raji cultures Sup T. Add one fourth volume fresh cells.	negative Donors 1-6 Raji negative Donors 1-6 SupT
21	No CPE. Add one fourth volume fresh cells.	negative Donors 1-6 Raji negative Donors 1-6 SupT
25	No CPE. Terminate culture.	negative Donors 1-6 Raji negative Donors 1-6 SupT