Supporting information, S1

The health status of transgender and gender nonbinary adults in the United States

Ethan C. Cicero^{1*}, Sari L. Reisner²⁻⁵, Elizabeth I. Merwin^{6, 7, #a}, Janice C. Humphreys⁶, Susan G. Silva^{6,8}

S1 Table. Significant Bivariate Model Results: *A Posteriori* Contrasts for Study Group, Individual Factors, Seasonality, and Health Status Outcomes.

	Pairwise contrasts		
Factor	TW/TM	TW/GNB	TM/GNB
Personal characteristic			
Sexual minority			
P value	0.0444	0.0046	<0.001
OR	1.49	0.57	0.38
95% CI	1.01, 2.21	0.38, 0.84	0.24, 0.59
Socioeconomic position			
Uninsured			
P value	0.0106	0.0733	0.7214
OR	0.54	0.59	1.11
95% CI	0.33, 0.86	0.34, 1.05	0.63, 1.96
Health status outcome	·	•	
Fair/poor health			
P value	0.4891	0.0245	0.0081
OR	1.14	0.64	0.56
95% CI	0.78, 1.67	0.44, 0.94	0.37, 0.86

Bold indicates statistical significance at the 0.05 level. CI: confidence interval;

GNB: gender nonbinary adults; OR: odds ratio; TM: transgender men; TW: transgender women.

^{*}Corresponding author. E-mail: ethan.cicero@ucsf.edu; ¹ Department of Community Health Systems, University of California San Francisco School of Nursing, San Francisco, California; ² Department of Epidemiology, Harvard T.H. Chan School of Public Health, Boston, Massachusetts; ³ Department of Pediatrics, Harvard Medical School, Boston, Massachusetts; ⁴ Division of General Pediatrics, Boston Children's Hospital, Boston, Massachusetts; ⁵ The Fenway Institute, Fenway Health, Boston, Massachusetts; ⁶ School of Nursing, Duke University, Durham, North Carolina; ⁷ College of Nursing and Health Innovation, The University of Texas at Arlington, Arlington, Texas; ⁸ School of Medicine, Duke University, Durham, North Carolina