Supporting Information

A Specific Proteome of Hippocampal Regions CA2 and CA1 Reveals Proteins Linked to the Unique Physiology of Area CA2

Kyle J. Gerber^{1†}, Eric B. Dammer^{2,3†}, Duc M. Duong^{2,3}, Qiudong Deng^{2,3}, Serena M. Dudek⁴, Nicholas T. Seyfried^{2,3,5}, John R. Hepler^{1*}

Author Affiliations

- 1 Department of Pharmacology and Chemical Biology, Emory University School of Medicine, Rollins Research Center, Atlanta, Georgia, USA.
- 2 Department of Biochemistry, Emory University School of Medicine, Atlanta, Georgia, USA.
- 3 Center for Neurodegenerative Disease, Emory University.
- 4 Neurobiology Laboratory, National Institute of Environmental Health Sciences, National Institutes of Health, Research Triangle Park, North Carolina, USA.
- 5 Department of Neurology, Emory University School of Medicine, Atlanta, Georgia, USA.

Table of Contents

Supplemental Figure 1 – Box plots showing differential expression between CA1 and CA2 for proteins included in chord plots

Supplemental Figure 2 – Box plots showing differential expression between CA1 and CA2 across all mass spectrometry experiments for proteins included in chord plots

Available separately:

Supplemental Table 1 – Meta-analysis of quantitative proteomics

Supplemental Table 2 – Additional CA2 markers in TMT proteomics


Supplemental Table 3 – Information on dissected mouse samples


Supplemental Table 4 – Significance of differentially expressed proteins in modules

[†] Authors contributed equally.

^{*} Corresponding Author; Email address: jhepler@emory.edu.

Supplemental Figure 1 – Box plots showing differential expression between CA1 and CA2 for proteins included in chord plots





Supplemental Figure 2 – Box plots showing differential expression between CA1 and CA2 across all mass spectrometry experiments for proteins included in chord plots

