

Supplementary

Evaluation of the Use of an Inorganic Bone Matrix in the Repair of Bone Defects in Rats Submitted to Experimental Alcoholism

Iris Jasmin Santos German ^{1,2,3}, Karina Torres Pomini ¹, Ana Carolina Cestari Bighetti ¹, Jesus Carlos Andreo ¹, Carlos Henrique Bertoni Reis ⁴, André Luis Shinohara ¹, Geraldo Marco Rosa Júnior ^{5,6}, Daniel de Bortoli Teixeira ⁴, Marcelie Priscila de Oliveira Rosso ¹, Daniela Vieira Buchaim ^{4,7} and Rogério Leone Buchaim ^{1,4,*}

- ¹ Department of Biological Sciences (Anatomy), Bauru School of Dentistry, University of São Paulo (USP), Bauru, São Paulo, 17012-901, Brazil; irish_knaan@hotmail.com (I.J.S.G.); karinatorrespomini@gmail.com (K.T.P.); anacarolinacb25@gmail.com (A.C.C.B.); jcandreo@usp.br (J.C.A.); andreshinohara@yahoo.com.br (A.L.S.); marcelierosso@usp.br (M.P.O.R.)
 - ² Department of Dentistry, Faculty of Health Science, Universidad Iberoamericana (UNIBE), Santo Domingo 10203, Dominican Republic
 - ³ Mother and Teacher Pontifical Catholic University (PUCMM), Santo Domingo, Dominican Republic;
 - ⁴ Postgraduate Program in Structural and Functional Interactions in Rehabilitation, University of Marília (UNIMAR), Marília, São Paulo, 17525-902, Brazil; carlosbtreis@yahoo.com.br (C.H.B.R.); daniel.dbt@hotmail.com (D.B.T.); danibuchaim@usp.br (D.V.B.)
 - ⁵ University of the Ninth of July (UNINOVE), Bauru, São Paulo, 17011-102, Brazil; geraldomjr@yahoo.com.br
 - ⁶ University of the Sacred Heart (USC), Bauru, São Paulo, 17011-160, Brazil
 - ⁷ Medical School, University Center of Adamantina (UniFAI), Adamantina, São Paulo, 17800-000, Brazil;
- * Correspondence: rogerio@fob.usp.br

Received: 13 December 2019; Accepted: 27 January 2020; Published: 3 February 2020




Figure S1: Representative image of the methodology used to quantify the area of newly formed bone, graft particles, by Axio Vision software.