

BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

Global prevalence of surgical-site infection after appendectomy: a systematic review and meta-analysis

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2019-034266
Article Type:	Original research
Date Submitted by the Author:	12-Sep-2019
Complete List of Authors:	Danwang, Celestin; Universite Libre de Bruxelles, School Of Public Health; Bigna, Jean Joel; Centre Pasteur of Cameroon, Department of Epidemiology and Public Health Tochie, Joel Noutakdie; Universite de Yaounde I Faculte de Medecine et des Sciences Biomedicales, Mbonda, Aimé; Djohong District Hospital, Department of General Medicine Mbanga, Clarence; Mankon Sub-divisional Hospital Nzalie, Rolf ; Ngong District Hospital, North Region, Cameroon Guifo, Marc Leroy ; Universite de Yaounde I Faculte de Medecine et des Sciences Biomedicales Essomba, Arthur ; Universite de Yaounde I Faculte de Medecine et des Sciences Biomedicales
Keywords:	EPIDEMIOLOGY, Health policy < HEALTH SERVICES ADMINISTRATION & MANAGEMENT, Quality in health care < HEALTH SERVICES ADMINISTRATION & MANAGEMENT, Epidemiology < INFECTIOUS DISEASES, Public health < INFECTIOUS DISEASES

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Global prevalence of surgical-site infection after appendectomy: a 2 systematic review and meta-analysis

3
4 Celestin Danwang, MD^{1,2}, Jean Joel Bigna, MD^{3,4*}, Joel Noutakdie Tochie, MD¹, Aime
5 Mbonda, MD⁵, Clarence Mvalo Mbanga, MD⁶, Rolf Nyah Tuku Nzalie, MD⁷, Marc Leroy
6 Guifo, MD¹, Arthur Essomba, MD¹

- 7
8 1. Department of Surgery and Specialties, Faculty of Medicine and Biomedical Sciences,
9 University of Yaoundé I, Yaoundé, Cameroon.
10 2. School of Public Health, Université Libre de Bruxelles, Brussels, Belgium.
11 3. Department of Epidemiology and Public Health, Centre Pasteur of Cameroon, Yaoundé,
12 Cameroon.
13 4. School of Public Health, Faculty of Medicine, University of Paris Sud XI, Le Kremlin-
14 Bicêtre, Paris, France.
15 5. Department of General Medicine, Djohong District Hospital, Djohong, Cameroon.
16 6. Department of General Medicine, Mankon Sub-Divisional Hospital, Bamenda, Cameroon.
17 7. Department of General Medicine, Ngong District Hospital, Ngong, Cameroon.
18

19 *Corresponding author: Dr. Jean Joel Bigna, MD, MPH

20 Department of Epidemiology and Public Health, Centre Pasteur of Cameroon, Yaoundé,
21 Cameroon. E-Mail: bignarimjj@yahoo.fr

22
23 E-mail addresses:

24 CD: danram07@yahoo.fr; JJB: bignarimjj@yahoo.fr; JNT: joeltochie@gmail.com; AM:
25 aimembonda@yahoo.fr; CMM: mbangaclarence@gmail.com; RNN: nzaliierolf@yahoo.com;
26 GML: marcleroyguifo@gmail.com; AE: artgessomba@gmail.com

27
28 **Word count:** 2,618.

29 **Abstract**

30 **Objectives:** Although surgical-site infection (SSI) is one of the most studied healthcare-
31 associated infections, the global burden of SSI after appendectomy remains unknown. Hence,
32 we estimated the incidence of SSI after appendectomy at global and regional levels.

33 **Design:** Systematic review and meta-analysis.

34 **Participants:** Global population of patients with appendectomy.

35 **Data sources:** EMBASE, Medline, and Web of Science were searched to identify observational
36 studies, published between January 1, 2000 and December 30, 2018 and reporting on the
37 incidence of the SSI after appendectomy with no language restriction. A random-effect models
38 meta-analysis served to obtain the pooled incidence of SSI after 100 surgical procedures in
39 patients with appendicitis.

40 **Results:** In total, 226 studies (729,434 participants from 49 countries) were included in the
41 meta-analysis. Concerning the methodological quality, 59 (26.1%) studies had a low risk, 147
42 (65.0%) a moderate risk, and 20 (8.8%) a high risk of bias. We found an overall incidence of
43 SSIs of 7.0 per 100 surgical procedures (95% prediction interval: 1.0-17.6) for appendectomy
44 varying from 0 to 37.4 per 100 surgical procedures. Subgroup analysis for identifying sources
45 of heterogeneity showed that the incidence varied from 5.8 in Europe to 12.6 per 100 surgical
46 procedures in Africa, $p < 0.0001$. The incidence of SSI after appendectomy increased when the
47 level of income decreased; from 6.2 in high-income countries to 11.1 per 100 surgical
48 procedures in low-income countries ($p = 0.015$). Open appendectomy (11.0 per 100 surgical
49 procedures) was found to have a higher incidence of SSI compared to laparoscopy (4.6 per 100
50 surgical procedures), $p = 0.0002$.

51 **Conclusion:** This study suggests a high burden of SSIs after appendectomy in some regions
52 (especially Africa) and in low-income countries. Strategies are needed to implement and
53 vulgarize WHO guidelines to decrease the burden of SSI after appendectomy in these regions.

1
2
3 54 **Registration:** PROSPERO, CRD42017075257.
4
5
6 55
7
8 56
9

10 57 **Keywords:**
11

12 58 Surgical wound infection; Global Health; Hospital infections; Cross infection; Healthcare
13
14
15 59 associated infection
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

60 **Strengths and limitations of this study**

- 61 • This systematic review with the meta-analysis is the first figuring-out a comprehensive
62 global summary of the existing knowledge on the incidence of SSIs after appendectomy.
- 63 • It is also the first to summarize the burden of SSIs after appendectomy by all World Health
64 Organization (WHO) regions and by country level of income.
- 65 • Using strong and robust methodological and statistical procedures, we found an overall high
66 incidence of SSIs, seven per 100 surgical procedures, with substantial heterogeneity
67 according to WHO regions and country level of income.
- 68 • WHO Afro region had the highest incidence. The incidence also increased with decreasing
69 country level of income.
- 70 • All sources of heterogeneity for the variation in the incidence of SSI after appendectomy
71 were not identified.

72 **Introduction**

73 Defined as an acute inflammation of the vermiform appendix (1), evidence abounds that acute
74 appendicitis is the most common abdominal surgical emergency (2), with an incidence of
75 almost 100 per 100,000 person-years reported in Australia, Europe and North America (3,4).
76 Evidence suggests appendectomy, a surgical remove of the vermiform appendix as first-line
77 treatment for acute appendicitis, although antibiotic therapy may be efficacious for a selected
78 group of patients with uncomplicated acute appendicitis (5,6). Appendectomy is a relatively
79 safe surgical intervention with a case fatality rate of 2.1 - 2.4 per 1000 patients as reported in
80 studies conducted in Europe (7,8).

81 Innovations in appendectomy, especially with the advent of minimally invasive or laparoscopic
82 surgery in 1983 (9), which has replaced the traditional open appendectomy in most of high-
83 income countries, has led to a drastic reduction in the morbidity and mortality related to
84 appendectomy (10–12). Laparoscopic appendectomy is now recognized as the gold standard
85 surgical approach for uncomplicated acute appendicitis owing to its merits over open surgery;
86 due to less postoperative pain, reduced postoperative ileus, shorter hospital stay, rapid
87 postoperative recovery, and better aesthetic scars (13–17).

88 However, regardless of the surgical technique (laparoscopic or open surgery), appendectomy
89 remains a sceptical surgical intervention associated with a substantial risk of surgical-site
90 infections (SSIs). SSIs after appendectomy are postoperative nosocomial infections affecting
91 the incision site, deep tissues, organs at the operative site within 30 days after the surgical
92 procedure (18–21). SSI following appendectomy is a serious post-operative medical concern
93 that increases the financial burden for both healthcare systems and patient, and also have a
94 negative impact on the patients' health related quality of life (22–27).

95 SSI is both the most frequently studied and the leading healthcare-associated infections reported
96 hospital-wide in low- and middle-income countries (28). A recently published prospective

1
2
3 97 international multicentre cohort study suggested a high burden of SSIs after any gastrointestinal
4
5 98 surgery in low-income countries compared to high-income countries (29). Actually, there is no
6
7
8 99 global systematic review with meta-analysis reporting the burden of SSI after appendectomy or
9
10 100 comparing the burden between regions and between country level of income. It would be
11
12 101 interesting to have such accurately estimated data to construct efficient strategies to curb
13
14 102 globally the burden of SSIs after appendectomy. In an effort to fill this gap, the current
15
16
17 103 systematic review and meta-analysis aimed at summarizing contemporary data on the
18
19 104 occurrence of SSIs after appendectomy.
20

21 105

23 106 **Methods**

24 107 **Search strategy and selection criteria**

25
26
27
28
29 108 We searched EMBASE, Medline, and Web of Science (Web of Science Core Collection,
30
31 109 Current Contents Connect, KCI-Korean Journal Database, SciELO Citation Index, Russian
32
33 110 Science Citation Index) to identify observational studies, published between January 1, 2000
34
35 111 and December 30, 2018 and reporting data on the incidence of SSIs after appendectomy. No
36
37 112 language restriction was applied. The initial search strategy was designed for EMBASE and
38
39 113 was adapted for the use in others databases. The search strategy as illustrated in the study
40
41 114 protocol (30), was based on the combination of relevant text words and medical subject
42
43 115 headings related to SSIs. Moreover, the references of all relevant articles found were scrutinized
44
45 116 for potential additional data sources. When a full text was not available, it was requested via
46
47 117 the corresponding author by email. For duplicates or studies published in more than one report,
48
49 118 the one reporting the largest sample size was considered. We excluded letters, reviews,
50
51 119 commentaries and editorials, studies lacking key data and/or explicit method description as well
52
53 120 as studies in which relevant data on SSIs after appendectomy was impossible to extract even
54
55 121 after contacting the corresponding author.
56
57
58
59
60

1
2
3 122 Two reviewers (CD and AM) independently screened the titles and abstract of articles for
4
5 123 eligibility. Full texts of potentially eligible articles were retrieved and screened for final
6
7 124 inclusion. Disagreements between the two reviewers were solved by discussion and when a
8
9 125 consensus was not reached, a third reviewer (JNT) resolved discrepancies.

12 126 **Data analysis**

15 127 A standardized and pretested data extraction form was used by five reviewers (CD, JNT, AM,
16
17 128 RNZ, CMM) to independently extract data from individual studies. A sixth reviewer (JJB)
18
19 129 independently extracted data for accuracy. The last name of the first author, year of publication,
20
21 130 country, study design, age groups, sample size, mean or median age, proportion of males,
22
23 131 specific conditions of the study population, the surgical method (open surgery or laparoscopy),
24
25 132 and incidence of SSIs after appendectomy in the study population (or enough data to compute
26
27 133 this estimate). For multinational studies, data was disaggregated, with the results shown within
28
29 134 individual country.

33 135 A meta-analysis was used to summarize data concerning incidence of SSIs, by pooling together
34
35 136 data of studies reporting the incidence of SSIs. Study-specific estimates were then pooled
36
37 137 through a Dersimonian and Laird random-effects meta-analysis model to obtain an overall
38
39 138 summary estimate of the incidence across studies, after stabilizing the variance of individual
40
41 139 studies using the Freeman-Tukey double arc-sine transformation (31). Incidence was expressed
42
43 140 by 100 surgical procedures with their 95% confidence interval and 95% prediction interval.
44
45 141 Heterogeneity was evaluated by the χ^2 test on Cochrane's Q statistic (32) which is quantified
46
47 142 by I^2 values, assuming that I^2 values of 25%, 50% and 75% represent low, medium and high
48
49 143 heterogeneity respectively (33). Where substantial heterogeneity ($I^2 > 50\%$) was detected, a
50
51 144 subgroup analysis was performed to detect its possible sources using the following grouping
52
53 145 variables: type of surgery (laparoscopy or open), World Health Organization regions, and
54
55 146 country level of income. A p value <0.05 was indicative of significant difference. The meta-
56
57
58
59
60

1
2
3 147 regression analysis was performed to estimate the explained heterogeneity of each covariate
4
5 148 included in the subgroup analysis. Inter-rater agreement for study inclusion was assessed using
6
7 149 Cohen's κ coefficient (34). Funnel plots analysis and Egger's test ($p < 0.10$) were performed to
8
9 150 detect the presence of publication bias (35). Since we believe that the incidence estimates of interest
10
11 151 would likely be published even if substantially different from previously reported estimates, we have
12
13 152 not reported adjusted incidence estimate in the case of publication bias.

14
15
16 153 To assess the methodological quality of each study, two reviewers (CD and CMM) used an
17
18 154 adapted version of the tool of bias assessment for prevalence studies developed by Hoy and
19
20 155 colleagues (36).

21
22
23 156 This review was registered in the International Prospective Register of Systematic Reviews
24
25 157 (PROSPERO) under the registration number CRD42017075257. The protocol has been
26
27 158 published in a peer-review journal (30).

28 29 30 159 **Patient and public involvement**

31
32 160 Patients or the public were not involved in the design, or conduct, or reporting, or dissemination
33
34 161 of our research.

35
36
37 162

38 39 40 163 **Results**

41
42 164 Overall, 619 records were initially identified. After removal of duplicates, screening of study
43
44 165 titles, abstracts, and full texts; 226 studies including 729,434 patients were finally retained for
45
46 166 meta-analysis (Supplementary Figure 1). The full list of included studies is in the Appendix.
47
48 167 Concerning the methodological quality, 59 (26.1%) studies had a low risk, 147 (65.0%) a
49
50 168 moderate risk and 20 (8.8%) a high risk of bias. Supplementary Table 1 presents characteristics
51
52 169 of included studies. Among the included studies, 154 were done in high-income, 36 upper-
53
54 170 middle, 27 lower-middle, and nine in low-income countries. Overall, most of studies were from
55
56 171 Europe ($n = 68$) and Americas ($n = 67$). SSIs were defined according to Center of Disease
57
58
59
60

1
2
3 172 Control and Prevention criteria in 50 studies while 25 studies used other criteria. The definition
4
5 173 of SSIs was not clearly given in 151 studies.

7
8 174 The overall incidence of SSI after appendectomy was 7.0 per 100 surgical procedures (95%
9
10 175 prediction interval: 1.0-17.6) varying from 0% to 37.4% with substantial heterogeneity and
11
12 176 publication bias. The sensitive analysis including only studies with low risk of bias yielded a
13
14 177 very close incidence to crude analysis (Table 1).

16
17 178 According to country level of income (Figure 1), the incidence of SSI after appendectomy
18
19 179 increased when the level of income decreased; from 6.2 in high income countries to 11.1 per
20
21 180 100 surgical procedures in low income countries ($p = 0.015$) (Table 1).

23
24 181 The incidence varied widely across WHO regions (Figure 2). The incidence varied from 5.8 in
25
26 182 Europe to 12.6 per 100 surgical procedures in Africa, $p < 0.0001$ (Table 1). Two regions
27
28 183 (Europe and Americas) had an incidence < 6 per 100 surgical procedures, three an incidence
29
30 184 between 6-10 per 100 procedures (South-East Asia, Eastern Mediterranean, and Western
31
32 185 Pacific), and one an incidence > 10 per 100 procedures (Africa) (Table 1). The incidence also
33
34 186 varied widely in different regions. The incidence varied from 0.2 to 32.0 in Africa, from 1.9 to
35
36 187 37.4 in Western Pacific, from 1.3 to 33.8 in Eastern Mediterranean, from 1.2 to 25.8 in South-
37
38 188 East Asia, from 0.1 to 37.4 in Americas, and from 0 to 20.0 per 100 surgical procedures in
39
40 189 Europe (Figure 2).

42
43
44 190 Open appendectomy with an incidence of 11.0 (95% prediction interval: 0.0-39.3) per 100
45
46 191 surgical procedures was found to have a higher incidence of SSI compared to laparoscopic
47
48 192 appendectomy with an incidence of 4.6 (95% prediction interval: 0.0-14.3) per 100 surgical
49
50 193 procedures, $p = 0.0002$ (Figure 3).

51
52
53 194 Heterogeneity of the overall incidence of SSI after appendectomy was explained by WHO
54
55 195 regions (17.1%), country level of income (11.1%), and type of surgical procedure (0.1%).

56
57
58 196
59
60

197 Discussion

198 This first systematic review and meta-analysis of data of 729434 surgical procedures in 226
199 studies from 49 countries found an overall incidence of SSIs of 7.0 per 100 surgical procedures
200 for appendectomy varying from 0 to 37.4 per 100 surgical procedures with substantial
201 heterogeneity according to WHO regions, country level of income, and type of surgical
202 procedure. The incidence increased with decreasing country level of income and was higher
203 when using open surgery compared to laparoscopy. The incidence significantly varied by WHO
204 regions with Africa having the highest burden followed by Western Pacific, Eastern-
205 Mediterranean, and South-East Asia.

206 Health care-associated infections are acquired by patients when receiving care and are the most
207 frequent adverse event affecting patient safety worldwide. This includes SSIs after
208 appendectomy (37). As reported in a systematic review and meta-analysis, SSIs were the
209 leading infection in hospitals in developed countries (28). The high incidence we found in this
210 study suggests that SSIs after appendectomy remains a global public concern. WHO reported
211 that of every 100 hospitalized patients at any given time, seven in developed and 15 in
212 developing countries will acquire at least one health care-associated infection (37). SSIs are
213 mainly caused by micro-organisms resistant to commonly-used antimicrobials, which can be
214 multidrug-resistant. Indeed, more than 50% of SSIs can be antibiotic-resistant (38). The leading
215 micro-organisms identified in SSIs are *Staphylococcus aureus*, coagulase-negative
216 staphylococci, and *Escherichia coli* as reported by National Healthcare Safety Network (38). It
217 is important to worry since *Staphylococcus aureus* and *Escherichia coli* are the micro-
218 organisms with highest proportion of antibiotic resistance, respectively resistant to
219 oxacillin/methicillin in 43% of cases and to fluoroquinolones in 25% of cases (38). A recent
220 international prospective cohort study shown that 21.6% of patients with SSI after any
221 gastrointestinal surgery had an infection that was resistant to the prophylactic antibiotic used

1
2
3 222 (29). There are many factors that can favour SSI including patient-related and procedural-
4
5 223 related variable (39). These factors can be classified in two categories; non-modifiable like age
6
7 224 and sex and modifiable including nutritional status, tobacco use, correct use of antibiotics,
8
9
10 225 obesity, diabetes, prolonged surgery duration, pre-surgery hospital stay of at least two days,
11
12 226 lower volume of hospital and surgeons, and the intraoperative techniques (37). Strategies to
13
14 227 curb the burden of SSIs should therefore focus on addressing these identified factors.

16
17 228 In our present study looking at specifically SSI after appendectomy, we also found that SSI was
18
19 229 higher in low income countries. Interestingly, there was a trend with increasing incidence when
20
21 230 the country income decreased. The WHO Africa region essentially constituted with sub-
22
23 231 Saharan Africa was the region with highest incidence in this study. The WHO estimates that
24
25 232 the endemic burden of health care-associated infections is two to three time significantly higher
26
27 233 in low- and middle-income countries than in high-income nations (37). The highest burden
28
29 234 found in Africa may be associated with the fact most of countries in this continent are low
30
31 235 income countries compared to other regions. Indeed, factors associated with increased risk of
32
33 236 SSI after appendectomy may be higher in low-income settings. The burden of diabetes, obesity,
34
35 237 and undernutrition are increasing in low-income countries (40,41). There is also inadequate use
36
37 238 of antimicrobial in low- and middle-income countries (42,43) and micro-organisms are more
38
39 239 resistant to prophylactic antibiotics used to prevent SSI in low-income countries compared to
40
41 240 high-income countries (29). Lower level income is also associated with lower volume of
42
43 241 surgeon and hospital, factors recognised as associated increased risk of SSIs (37). The higher
44
45 242 incidence found in low income countries may also be explained by the fact open surgery is the
46
47 243 most used surgical procedure in this setting. Indeed, we found as in other studies that open
48
49 244 surgery is associated with higher incidence of SSIs compared to laparoscopy (44,45).
50
51 245 Laparoscopy is generally indicated for uncomplicated appendicitis where the dissemination of
52
53 246 micro-organism is lower compared open surgery indicated for perforated appendicitis with
54
55
56
57
58
59
60

1
2
3 247 peritonitis for example. Moreover, only few low-income countries have the necessary
4
5 248 infrastructure to carry out laparoscopy procedures compared to high-income countries (46–48).
6
7
8 249 Our findings have important implications for healthcare providers and health policy makers.
9
10 250 SSIs are among the most preventable healthcare-associated infections (49,50). They still
11
12 251 represent a significant burden in terms of patient morbidity and mortality and additional costs
13
14 252 for healthcare systems (37). The prevention of SSI has received considerable attention from
15
16
17 253 surgeons, infection control professionals, health policy makers, the media and the public since
18
19 254 there is a perception among the public that SSIs may reflect a poor quality of care (51).
20
21 255 However, special attention is needed for low-income countries and Africa. Strategy to curb the
22
23
24 256 burden of SSIs after appendectomy as for other surgery procedures should be focused on
25
26 257 strategies than can help to address factors associated with increased risk of SSIs. Therefore,
27
28 258 strategies should be a package including how to address the factors cited above. The 26 WHO
29
30 259 recommendations to avoid SSIs should be vulgarized and implemented (37), especially in low-
31
32
33 260 income countries. Strengthening the healthcare systems of low-income countries and of
34
35 261 countries in WHO Afro region is also a paramount by education of healthcare providers and
36
37 262 skilling them on the use of very less invasive surgical procedures.
38
39
40 263 This study should however be interpreted in the context of some drawbacks. Firstly, the same
41
42 264 definition of SSIs was not used by all the included studies. This may lead to an overestimation
43
44 265 or underestimation of the SSIs incidence by individual studies (depending on the definition
45
46 266 used). Secondly, few studies reported on the associated conditions of the study population since
47
48
49 267 this can modify the risk for developing SSIs. We were not therefore able to measure the impact
50
51 268 on our outcome of interest. Thirdly, only a quarter of studies had low risk of bias, however our
52
53 269 analysis including only studies with low risk of bias yielded an estimate close to the crude
54
55
56 270 incidence. Fourth, the various geographic regions and countries were variably represented, with
57
58
59
60

1
2
3 271 some countries with only one study or even no study, which could affect the generalizability of
4
5 272 our findings.

6
7 273 Despite these limitations, this is the first systematic review and meta-analysis providing a global
8
9 274 estimate of the burden of SSIs after appendectomy. A protocol had been published before, and
10
11 275 we used rigorous methodological and statistical procedures to obtain and pool data.
12
13 276 Furthermore, subgroup analyses were conducted to investigate the various factors likely
14
15 277 affecting our estimate.

16
17 278 This systematic review and meta-analysis compiled data from more than 700,000 people with
18
19 279 appendicitis in 49 countries and pointed a high incidence of SSIs after appendectomy, at 7 per
20
21 280 100 surgical procedures. This estimate seemed higher in some WHO regions (especially Africa)
22
23 281 and in low-income countries. These data suggest that less invasive procedure is associated with
24
25 282 low incidence of SSIs after appendectomy. Strategies are needed to implement already known
26
27 283 guidelines to decrease the burden of SSI after appendectomy. However, in low-income
28
29 284 countries which have weak health systems, cost-effectiveness studies are needed to inform
30
31 285 policies regarding the best strategies for decreasing the burden of SSI after appendectomy.
32
33 286

34
35
36
37
38 286

39 40 287 **Contributors**

41
42
43 288 CD and JJB conceived the idea of the study and developed the protocol. JJB, CD, and JNT did
44
45 289 the literature search. CD, AM, and JNT selected the studies, CD, JNT, RNZ, AM, CMM, JJB
46
47 290 extracted the relevant information. CD, JJB, and CMM synthesized the data. CD, JNT, CMM,
48
49 291 and JJB wrote the first draft of the paper. CD, JJB, JNT, AB, RNZ, CMM, GML, and AE
50
51 292 critically revised successive drafts of the paper and approved the final version. GML and AE
52
53 293 supervised the overall work, CD and JJB are the guarantors of the review.

54 55 294 **Acknowledgements**

56
57
58
59 295 None.
60

1
2
3 296 **Funding**
4

5 297 This research received no specific grant from any funding agency in the public, commercial or
6
7 298 not-for-profit sectors.
9

10 299 **Competing interests**
11

12 300 We declare no competing interests.
13

14 301 **Patient consent**
15

16 302 Not applicable.
17
18

19 303 **Data sharing statement**
20

21 304 All data generated for this study are in the manuscript and its supporting files.
22
23
24 305

25
26 306 **Figures Legend**
27

28 307 Figure 1. Global incidence of SSI (surgical site infection) after appendectomy by level of
29
30 308 country income
31
32

33 309 Figure 2. Global incidence of SSI (surgical site infection) after appendectomy by WHO regions
34

35 310 Figure 3. Global incidence of SSI (surgical site infection) after appendectomy by type of
36
37 311 surgical procedures
38
39

40 312

41
42 313
43

44 314 **References**
45

46
47 315 1. Giesen LJX, van den Boom AL, van Rossem CC, den Hoed PT, Wijnhoven BPL.
48
49 316 Retrospective Multicenter Study on Risk Factors for Surgical Site Infections after
50
51 317 Appendectomy for Acute Appendicitis. *Dig Surg*. 2016 Sep 16;34(2):103–7.
52
53

54 318 2. Navarro JF, Tárraga PL, Rodríguez JM, López MC. Validity of tests performed to
55
56 319 diagnose acute abdominal pain in patients admitted at an emergency department. *Rev Espanola*
57
58 320 *Enfermedades Dig Organo Of Soc Espanola Patol Dig*. 2009 Sep;101(9):610–8.
59
60

- 321 3. Ohmann C, Franke C, Kraemer M, Yang Q. [Status report on epidemiology of acute
322 appendicitis]. *Chir Z Alle Geb Oper Medizen*. 2002 Aug;73(8):769–76.
- 323 4. Körner H, Söreide JA, Pedersen EJ, Bru T, Söndena K, Vatten L. Stability in incidence
324 of acute appendicitis. A population-based longitudinal study. *Dig Surg*. 2001;18(1):61–6.
- 325 5. Varadhan KK, Neal KR, Lobo DN. Safety and efficacy of antibiotics compared with
326 appendectomy for treatment of uncomplicated acute appendicitis: meta-analysis of
327 randomised controlled trials. *BMJ*. 2012 Apr 5;344:e2156.
- 328 6. Masoomi H, Nguyen NT, Dolich MO, Mills S, Carmichael JC, Stamos MJ.
329 Laparoscopic appendectomy trends and outcomes in the United States: data from the
330 Nationwide Inpatient Sample (NIS), 2004-2011. *Am Surg*. 2014 Oct;80(10):1074–7.
- 331 7. Kotaluoto S, Ukkonen M, Pauniahho S-L, Helminen M, Sand J, Rantanen T. Mortality
332 Related to Appendectomy; a Population Based Analysis over Two Decades in Finland. *World
333 J Surg*. 2017;41(1):64–9.
- 334 8. Blomqvist PG, Andersson RE, Granath F, Lambe MP, Ekblom AR. Mortality after
335 appendectomy in Sweden, 1987-1996. *Ann Surg*. 2001 Apr;233(4):455–60.
- 336 9. Semm K. Endoscopic appendectomy. *Endoscopy*. 1983 Mar;15(2):59–64.
- 337 10. Xiao Y, Shi G, Zhang J, Cao J-G, Liu L-J, Chen T-H, et al. Surgical site infection after
338 laparoscopic and open appendectomy: a multicenter large consecutive cohort study. *Surg
339 Endosc*. 2015 Jun;29(6):1384–93.
- 340 11. Varela JE, Wilson SE, Nguyen NT. Laparoscopic surgery significantly reduces surgical-
341 site infections compared with open surgery. *Surg Endosc*. 2010 Feb;24(2):270–6.
- 342 12. Bregendahl S, Nørgaard M, Laurberg S, Jepsen P. Risk of complications and 30-day
343 mortality after laparoscopic and open appendectomy in a Danish region, 1998-2007;
344 a population-based study of 18,426 patients. *Pol Przegl Chir*. 2013 Jul;85(7):395–400.
- 345 13. Sauerland S, Jaschinski T, Neugebauer EA. Laparoscopic versus open surgery for

- 1
2
3 346 suspected appendicitis. *Cochrane Database Syst Rev.* 2010 Oct 6;(10):CD001546.
4
5 347 14. Aziz O, Athanasiou T, Tekkis PP, Purkayastha S, Haddow J, Malinovski V, et al.
6
7 348 Laparoscopic versus open appendectomy in children: a meta-analysis. *Ann Surg.* 2006
8
9 349 Jan;243(1):17–27.
10
11
12 350 15. Markides G, Subar D, Riyad K. Laparoscopic versus open appendectomy in adults with
13
14 351 complicated appendicitis: systematic review and meta-analysis. *World J Surg.* 2010
15
16 352 Sep;34(9):2026–40.
17
18
19 353 16. Wei B, Qi C-L, Chen T-F, Zheng Z-H, Huang J-L, Hu B-G, et al. Laparoscopic versus
20
21 354 open appendectomy for acute appendicitis: a metaanalysis. *Surg Endosc.* 2011 Apr;25(4):1199–
22
23 355 208.
24
25
26 356 17. Li X, Zhang J, Sang L, Zhang W, Chu Z, Li X, et al. Laparoscopic versus conventional
27
28 357 appendectomy--a meta-analysis of randomized controlled trials. *BMC Gastroenterol.* 2010 Nov
29
30 358 3;10:129.
31
32
33 359 18. Horan TC, Gaynes RP, Martone WJ, Jarvis WR, Emori TG. CDC definitions of
34
35 360 nosocomial surgical site infections, 1992: a modification of CDC definitions of surgical wound
36
37 361 infections. *Infect Control Hosp Epidemiol.* 1992 Oct;13(10):606–8.
38
39
40 362 19. European Centre for Disease Prevention and Control. Surveillance of surgical site
41
42 363 infections in Europe 2010-2011. Luxembourg: Publications Office; 2014.
43
44
45 364 20. Culver DH, Horan TC, Gaynes RP, Martone WJ, Jarvis WR, Emori TG, et al. Surgical
46
47 365 wound infection rates by wound class, operative procedure, and patient risk index. National
48
49 366 Nosocomial Infections Surveillance System. *Am J Med.* 1991 Sep 16;91(3B):152S-157S.
50
51
52 367 21. Horan TC, Andrus M, Dudeck MA. CDC/NHSN surveillance definition of health care-
53
54 368 associated infection and criteria for specific types of infections in the acute care setting. *Am J*
55
56 369 *Infect Control.* 2008 Jun;36(5):309–32.
57
58
59 370 22. Badia JM, Casey AL, Petrosillo N, Hudson PM, Mitchell SA, Crosby C. Impact of
60

- 1
2
3 371 surgical site infection on healthcare costs and patient outcomes: a systematic review in six
4
5 372 European countries. *J Hosp Infect.* 2017 May;96(1):1–15.
6
7
8 373 23. Thompson KM, Oldenburg WA, Deschamps C, Rupp WC, Smith CD. Chasing zero:
9
10 374 the drive to eliminate surgical site infections. *Ann Surg.* 2011 Sep;254(3):430–6; discussion
11
12 375 436-437.
13
14 376 24. Hawn MT, Vick CC, Richman J, Holman W, Deierhoi RJ, Graham LA, et al. Surgical
15
16 377 site infection prevention: time to move beyond the surgical care improvement program. *Ann*
17
18 378 *Surg.* 2011 Sep;254(3):494–9; discussion 499-501.
19
20
21 379 25. Alexander JW, Solomkin JS, Edwards MJ. Updated recommendations for control of
22
23 380 surgical site infections. *Ann Surg.* 2011 Jun;253(6):1082–93.
24
25
26 381 26. Andersson RE. Short-term complications and long-term morbidity of laparoscopic and
27
28 382 open appendectomy in a national cohort. *Br J Surg.* 2014 Aug;101(9):1135–42.
29
30
31 383 27. Impact of wound edge protection devices on surgical site infection after laparotomy:
32
33 384 multicentre randomised controlled trial (ROSSINI Trial) | *The BMJ* [Internet]. [cited 2018 Aug
34
35 385 22]. Available from: <https://www.bmj.com/content/347/bmj.f4305>
36
37
38 386 28. Allegranzi B, Nejad SB, Combescure C, Graafmans W, Attar H, Donaldson L, et al.
39
40 387 Burden of endemic health-care-associated infection in developing countries: systematic review
41
42 388 and meta-analysis. *The Lancet.* 2011 Jan 15;377(9761):228–41.
43
44
45 389 29. GlobalSurg Collaborative. Surgical site infection after gastrointestinal surgery in high-
46
47 390 income, middle-income, and low-income countries: a prospective, international, multicentre
48
49 391 cohort study. *Lancet Infect Dis.* 2018 May;18(5):516–25.
50
51
52 392 30. Danwang C, Mazou TN, Tochie JN, Nzalie RNT, Bigna JJ. Global prevalence and
53
54 393 incidence of surgical site infections after appendectomy: a systematic review and meta-analysis
55
56 394 protocol. *BMJ Open.* 2018 Aug 30;8(8):e020101.
57
58
59 395 31. Barendregt JJ, Doi SA, Lee YY, Norman RE, Vos T. Meta-analysis of prevalence. *J*
60

- 1
2
3 396 Epidemiol Community Health. 2013 Nov 1;67(11):974–8.
4
5 397 32. Cochran: The combination of estimates from different... - Google Scholar [Internet].
6
7 398 [cited 2018 Aug 23]. Available from:
9
10 399 [https://scholar.google.com/scholar_lookup?hl=en&volume=10&publication_year=1954&pag](https://scholar.google.com/scholar_lookup?hl=en&volume=10&publication_year=1954&pages=101-129&journal=Biometrics&author=GW+Cochran.&title=The+combination+of+estimates+from+different+experiments)
11
12 400 [es=101-](https://scholar.google.com/scholar_lookup?hl=en&volume=10&publication_year=1954&pages=101-129&journal=Biometrics&author=GW+Cochran.&title=The+combination+of+estimates+from+different+experiments)
13
14 401 [129&journal=Biometrics&author=GW+Cochran.&title=The+combination+of+estimates+fro](https://scholar.google.com/scholar_lookup?hl=en&volume=10&publication_year=1954&pages=101-129&journal=Biometrics&author=GW+Cochran.&title=The+combination+of+estimates+from+different+experiments)
15
16 402 [m+different+experiments](https://scholar.google.com/scholar_lookup?hl=en&volume=10&publication_year=1954&pages=101-129&journal=Biometrics&author=GW+Cochran.&title=The+combination+of+estimates+from+different+experiments)
17
18 403 33. Huedo-Medina TB, Sánchez-Meca J, Marín-Martínez F, Botella J. Assessing
19
20 404 heterogeneity in meta-analysis: Q statistic or I² index? Psychol Methods. 2006 Jun;11(2):193–
21
22 405 206.
23
24 406 34. Viera AJ, Garrett JM. Understanding interobserver agreement: the kappa statistic. Fam
25
26 407 Med. 2005 May;37(5):360–3.
27
28 408 35. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a
29
30 409 simple, graphical test. BMJ. 1997 Sep 13;315(7109):629–34.
31
32 410 36. Hoy D, Brooks P, Woolf A, Blyth F, March L, Bain C, et al. Assessing risk of bias in
33
34 411 prevalence studies: modification of an existing tool and evidence of interrater agreement. J Clin
35
36 412 Epidemiol. 2012 Sep;65(9):934–9.
37
38 413 37. World Health Organization. Global guidelines for the prevention of surgical site
39
40 414 infection. 2016.
41
42 415 38. Sievert DM, Ricks P, Edwards JR, Schneider A, Patel J, Srinivasan A, et al.
43
44 416 Antimicrobial-resistant pathogens associated with healthcare-associated infections: summary
45
46 417 of data reported to the National Healthcare Safety Network at the Centers for Disease Control
47
48 418 and Prevention, 2009-2010. Infect Control Hosp Epidemiol. 2013 Jan;34(1):1–14.
49
50 419 39. Buggy D. Can anaesthetic management influence surgical-wound healing? Lancet Lond
51
52 420 Engl. 2000 Jul 29;356(9227):355–7.
53
54
55
56
57
58
59
60

- 1
2
3 421 40. Seidell JC, Halberstadt J. The global burden of obesity and the challenges of prevention.
4
5 422 *Ann Nutr Metab.* 2015;66 Suppl 2:7–12.
6
7
8 423 41. Checkley W, Ghannem H, Irazola V, Kimaiyo S, Levitt NS, Miranda JJ, et al.
9
10 424 Management of NCD in low- and middle-income countries. *Glob Heart.* 2014 Dec;9(4):431–
11
12 425 43.
13
14 426 42. Versporten A, Zarb P, Caniaux I, Gros M-F, Drapier N, Miller M, et al. Antimicrobial
15
16 427 consumption and resistance in adult hospital inpatients in 53 countries: results of an internet-
17
18 428 based global point prevalence survey. *Lancet Glob Health.* 2018 Jun;6(6):e619–29.
19
20
21 429 43. Klein EY, Van Boeckel TP, Martinez EM, Pant S, Gandra S, Levin SA, et al. Global
22
23 430 increase and geographic convergence in antibiotic consumption between 2000 and 2015. *Proc*
24
25 431 *Natl Acad Sci U S A.* 2018 10;115(15):E3463–70.
26
27
28 432 44. Foster D, Kethman W, Cai LZ, Weiser TG, Forrester JD. Surgical Site Infections after
29
30 433 Appendectomy Performed in Low and Middle Human Development-Index Countries: A
31
32 434 Systematic Review. *Surg Infect.* 2018 Apr;19(3):237–44.
33
34
35 435 45. Marchi M, Pan A, Gagliotti C, Morsillo F, Parenti M, Resi D, et al. The Italian national
36
37 436 surgical site infection surveillance programme and its positive impact, 2009 to 2011.
38
39 437 *Eurosurveillance.* 2014 May 29;19(21):20815.
40
41
42 438 46. Udwardia TE. Diagnostic laparoscopy. *Surg Endosc.* 2004 Jan;18(1):6–10.
43
44
45 439 47. Adisa AO, Lawal OO, Arowolo OA, Alatise OI. Local adaptations aid establishment of
46
47 440 laparoscopic surgery in a semiurban Nigerian hospital. *Surg Endosc.* 2013 Feb;27(2):390–3.
48
49 441 48. Alfa-Wali M, Osaghae S. Practice, training and safety of laparoscopic surgery in low
50
51 442 and middle-income countries. *World J Gastrointest Surg.* 2017 Jan 27;9(1):13–8.
52
53
54 443 49. Haley RW, Culver DH, White JW, Morgan WM, Emori TG, Munn VP, et al. The
55
56 444 efficacy of infection surveillance and control programs in preventing nosocomial infections in
57
58 445 US hospitals. *Am J Epidemiol.* 1985 Feb;121(2):182–205.
59
60

- 1
2
3 446 50. Harbarth S, Sax H, Gastmeier P. The preventable proportion of nosocomial infections:
4
5 447 an overview of published reports. *J Hosp Infect.* 2003 Aug;54(4):258–66; quiz 321.
6
7 448 51. Birgand G, Lepelletier D, Baron G, Barrett S, Breier A-C, Buke C, et al. Agreement
8
9 449 among Healthcare Professionals in Ten European Countries in Diagnosing Case-Vignettes of
10
11 450 Surgical-Site Infections. *PLOS ONE.* 2013 Jul 9;8(7):e68618.
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

Table 1. Summary statistics of meta-analysis incidence of surgery site infections after appendectomy


	Incidence per 100 surgical procedures (95%CI)	95% Prediction interval	N studies	N participants	H (95%CI)	I ² (95%CI)	P heterogeneity	P Egger test	P difference
Global	7.0 (6.4-7.7)	1.0-17.7	226	729,434	8.9 (8.7-9.1)	98.7 (98.7-98.8)	< 0.0001	< 0.0001	-
- Low risk of bias	6.9 (6.0-7.9)	1.6-15.2	59	204,450	6.7 (6.3-7.1)	97.7 (97.4-98.0)	< 0.0001	< 0.0001	-
By Level of income									
- Low	11.1 (5.5-18.2)	0.0-42.2	9	1,496	3.8 (3.0-4.8)	93.1 (89.0-95.6)	< 0.0001	0.735	0.015
- Lower-middle	9.2 (6.3-12.6)	0.0-31.6	27	10,379	5.1 (4.6-5.7)	96.2 (95.3-96.9)	< 0.0001	0.960	
- Upper-middle	8.5 (6.5-10.8)	0.3-25.3	36	26,557	5.4 (2.9-5.9)	96.6 (95.9-97.1)	< 0.0001	0.392	
- High	6.2 (5.6-6.9)	0.9-15.3	154	691,002	9.5 (9.2-9.8)	98.9 (98.8-99.0)	< 0.0001	< 0.0001	
By WHO regions									
- Africa	12.6 (3.3-26.4)	0.0-72.5	8	3,001	9.1 (7.9-10.5)	98.8 (98.4-99.1)	< 0.0001	0.628	< 0.0001
- Western Pacific	9.6 (8.1-11.2)	2.3-20.8	43	30,822	3.8 (3.5-4.2)	93.2 (91.7-94.4)	< 0.0001	0.150	
- Eastern Mediterranean	8.2 (6.4-10.2)	1.7-18.6	23	7,779	2.6 (2.2-3.1)	85.3 (79.1-89.6)	< 0.0001	0.515	
- South-East Asia	7.6 (4.7-11.1)	0.0-24.6	16	5,782	3.8 (3.2-4.5)	93.0 (90.1-95.0)	< 0.0001	0.0001	
- Americas	5.9 (5.2-6.6)	1.9-11.7	67	401,931	7.5 (7.1-7.9)	98.2 (98.0-98.4)	< 0.0001	0.0004	
- Europe	5.8 (4.6-7.0)	0.0-19.1	68	276,793	10.4 (10.0-10.8)	99.1 (99.0-99.1)	< 0.0001	< 0.0001	

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

By type of surgical procedure										
- Laparoscopy with open surgery	4.6 (2.5-7.2)	0.0-15.6	10	4,892	3.2 (2.6-4.2)	90.7 (85.0-94.2)	< 0.0001	0.942	0.0002	
- Laparoscopy	4.6 (3.4-5.9)	0.0-14.3	40	33,873	4.4 (4.0-4.8)	94.7 (93.6-95.7)	< 0.0001	0.0002		
- Open surgery	11.0 (7.9-14.4)	0.0-39.3	44	13,120	5.7 (5.2-6.1)	96.9 (96.4-97.3)	< 0.0001	0.077		

WHO: World Health Organization

For peer review only




Global burden of surgical-site infection after appendectomy: a systematic review and meta-analysis

APPENDIX

Celestin **Danwang**, Jean Joel **Bigna**, Joel Noutakdie **Tochie**,
Aime **Mbonda**, Clarence Mvalo **Mbanga**, Rolf Nyah Tuku **Nzalie**,
Marc Leroy **Guifo**, Arthur **Essomba**


<i>Supplementary Figure 1. Study Flow</i>	2
<i>Supplementary Figure 2. Funnel plot for publication bias</i>	3
<i>Supplementary Table 1. Characteristics of included studies</i>	4
<i>Reference list of included studies. References</i>	6



Supplementary Figure 1. Study flow

review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Supplementary Figure 2. Funnel plot for publication bias

Supplementary Table 1 : Characteristics of included studies

Characteristics	N = 226
Year of publication, range	2000-2018
%Male, range	0-100 (n = 195)
Mean/median age, range	7-74 (n = 186)
%HIV	0-13.1 (n = 2)
%Diabetes	0-95.7 (n = 34)
%Obesity	0-7.4 (n = 18)
Design, n	
- Cross sectional	120
- Cohort study	99
- Case control	7
WHO regions, n	
- Africa	8
- Americas	67
- Eastern Mediterranean	23
- Europe	68
- Multiregional	1
- South-East Asia	16
- Western Pacific	43
Level of income, n	
- Low	9
- Lower-middle	27
- Upper-middle	36
- High	154
Timing of data collection	
- Retrospective	123
- Prospective	101
- Unclear	2
Sampling	
- Consecutive	131
- Systematic	37
- Random	32
- Exhaustive	11
- Unclear	15
Number of sites	
- Multisite	51
- One site	170
- Unclear	5
Pattern of appendicitis, range	
- %Catarrhal	0-100 (n = 84)
- %Perforated	0-100 (n = 110)
- %Suppurated	0-100 (n = 70)
- %Gangrenous	0-46.7 (n = 89)
%With administered antibiotics	24.1-100 (n = 109)
%With administered analgesics	64.5-100 (n = 20)
%With diet > 6 or 8 hours	50-100 (n = 3)
Type of surgery	
- %Open surgery	0-100 (n = 134)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- %Laparoscopy	0-100 (n = 187)
Mean/median time to complete the intervention (in hours), range	0.1-2.2 (n = 106)
Type of anesthesia, n	
- General	118
- Spinal and general	2
- Unclear	106
SSI definition, n	
- CDC-NNIS criteria	50
- Other criteria	25
- Not reported/Unclear	151

For peer review only

References

1. Adejumo A.A., N.M., Mshelia, Y.M., Saleh. Clinicopathological presentation and management outcome of appendicitis in gombe, north-east nigeria: a 7-year retrospective audit. *Niger J Med.* 2015;24(4):337-43.
2. Aguiló J., S., Peiró, C., Muñoz, J., García del Caño, M., Garay, V., Viciano, et al. Adverse outcomes in the surgical treatment of acute appendicitis. *Efectos adversos en la cirugía de la apendicitis aguda.* 2005;78(5):312-7.
3. Adhikary S., S., Tyagi, G., Sapkota, A., Afaq, B.K., Bhattarai, C.S., Agrawal. Port exteriorization appendectomy: is it the future? *Nepal Med Coll J.* 2008;10(1):30-4.
4. Ahmad M., K., Ali, H., Latif, S., Naz, K., Said. Comparison of primary wound closure with delayed primary closure in perforated appendicitis. *J Ayub Med Coll Abbottabad.* 2014;26(2):153-7.
5. Ahmed I., J., Burr, M., Castillo, D., Collins, J.A., Cook, M., Campbell, et al. Single port/incision laparoscopic surgery compared with standard three-port laparoscopic surgery for appendectomy: A randomized controlled trial. *Surg Endosc Interv Tech.* 2015;29(1):77-85.
6. Akkoyun I., A., Taş Tuna. Advantages of abandoning abdominal cavity irrigation and drainage in operations performed on children with perforated appendicitis. *J Pediatr Surg.* 2012;47(10):1886-90.
7. Al-Saadi A.S., A.H., Al-Wadan, S.A., Hamarnah, H., Amin. Is abandoning routine peritoneal cultures during appendectomy justified? *Saudi Med J.* 2007;28(12):1827-9.
8. Al-Temimi M.H., M.A., Berglin, E.G., Kim, D.J., Tessier, S.D., Johna. Endostapler versus Hem-O-Lok clip to secure the appendiceal stump and mesoappendix during laparoscopic appendectomy. *Am J Surg.* 2017;214(6):1143-8.

- 1
2
3 9. Ali N., S., Aliyu. Appendicitis and its surgical management experience at the University of
4 Maiduguri Teaching Hospital Nigeria. *Niger J Med.* 2012;21(2):223-6.
5
6
7
- 8
9 10. Ali K., H., Latif, S., Ahmad. Frequency of wound infection in non-perforated appendicitis
10 with use of single dose preoperative antibiotics. *J Ayub Med Coll Abbottabad.*
11 2015;27(2):378-80.
12
13
- 14
15
16 11. Almström M., J.F., Svensson, B., Patkova, A., Svenningsson, T., Wester. In-hospital
17 surgical delay does not increase the risk for perforated appendicitis in children. *Ann Surg.*
18 2017;265(3):616-21.
19
20
21
- 22
23 12. Álvarez-Moreno C., A.M., Pérez-Fernández, V.D., Rosenthal, J., Quintero, E., Chapeta-
24 Parada, C., Linares, et al. Surgical site infection rates in 4 cities in Colombia: Findings of
25 the International Nosocomial Infection Control Consortium (INICC). *Am J Infect Control.*
26 2014;42(10):1089-92.
27
28
29
- 30
31
32 13. Andert A., H.P., Alizai, C.D., Klink, N., Neitzke, C., Fitzner, C., Heidenhain, et al. Risk
33 factors for morbidity after appendectomy. *Langenbeck's Arch Surg.* 2017;402(6):987-93.
34
35
36
- 37
38 14. Andersson R.E. Short-term complications and long-term morbidity of laparoscopic and
39 open appendectomy in a national cohort. *Br J Surg.* 2014;101(9):1135-42.
40
41
42
- 43
44 15. Aranda-Narváez J.M., A.J., González-Sánchez, N., Marín-Camero, C., Montiel-Casado, P.,
45 López-Ruiz, B., Sánchez-Pérez, et al. Conservative approach versus urgent appendectomy
46 in surgical management of acute appendicitis with abscess or phlegmon. *Resultados del*
47 *tratamiento conservador inicial y de la cirugía urgente en la apendicitis aguda evolucionada.*
48 2010;102(11):648-52.
49
50
51
- 52
53
54 16. Aranda-Narváez J.M., T., Prieto-Puga Arjona, B., García-Albiach, M.C., Montiel-Casado,
55 A.J., González-Sánchez, B., Sánchez-Pérez, et al. Postappendectomy surgical site infection:
56 Overall rate and type according to open/laparoscopic approach. *Infección de sitio quirúrgico*
57
58
59
60

- 1
2
3 tras apendicectomía urgente: tasa global y tipo según la vía de abordaje
4 (abierta/laparoscópica). 2014;32(2):76-81.
5
6
7
8
- 9 17. Arthur T., R., Gartrell, B., Manoharan, D., Parker. Emergency appendectomy in Australia:
10 findings from a multicentre, prospective study. ANZ J Surg. 2017;87(9):656-60.
11
12
13
- 14 18. Asefa Z. Acute appendicitis in Yirgalem Hospital, southern Ethiopia. Ethiop Med J.
15 2002;40(2):155-62.
16
17
- 18 19. Assefa Z., A., G/yesuse. Acute appendicitis in children admitted to zewditu memorial
19 hospital. Ethiop Med J. 2014;52(4):189-95.
20
21
22
- 23 20. Atif M.L., F., Sadaoui, A., Bezzaoucha, C.A., BezzaouchaKaddache, R., Boukari, S.,
24 Djelato, et al. Intra-abdominal abscesses and laparoscopic versus open appendectomies.
25 Infect Control Hosp Epidemiol. 2009;30(7):713-5
26
27
28
29
- 30 21. Bae E., A., Dehal, V., Franz, M., Joannides, N., Sakis, J., Scurlock, et al. Postoperative
31 antibiotic use and the incidence of intra-abdominal abscess in the setting of suppurative
32 appendicitis: a retrospective analysis. Am J Surg.2016;212(6):1121-5.
33
34
35
36
37
38
39
- 40 22. Bae S.U., W.K., Jeong, S.K., Baek. Single-port laparoscopic interval appendectomy for
41 perforated appendicitis with a periappendiceal abscess. Ann Coloproctol. 2016;32(3):105-
42 10.
43
44
45
46
47
- 48 23. Baek H.N., Y.H., Jung, Y.H., Hwang. Laparoscopic versus open appendectomy for
49 appendicitis in elderly patients. J Korean Soc Coloproctology. 2011;27(5):241-5.
50
51
52
- 53 24. Bali İ., F., Karateke, S., Özyazıcı, A., Kuvvetli, C., Oruç, E., Menekşe, et al. Comparison
54 of intracorporeal knotting and endoloop for stump closure in laparoscopic appendectomy.
55 Laparoskopik appendektomide intrakorporal düğüm ve endoloop ile güdük kapama
56 yöntemlerinin karşılaştırılması. 2015;21(6):446-9.
57
58
59
60

- 1
2
3
4
5 25. Bhangu A., Richardson, C., Torrance, A., Pinkney, T., Collaborative, Natl Surg Res.
6 Multicentre observational study of performance variation in provision and outcome of
7 emergency appendicectomy. *British Journal Of Surgery*. 2013;100(9):1240-52.
8
9
10
11
12 26. Bansal V., S., Altermatt, D., Nadal, C., Berger. Lack of benefit of preoperative antimicrobial
13 prophylaxis in children with acute appendicitis: A prospective cohort study. *Infection*.
14 2012;40(6):635-41.
15
16
17
18
19
20 27. Batajoo H., N.K., Hazra. Laparoscopic versus open appendectomy in acute appendicitis. *J*
21 *Nepal Health Res Counc*. 2012;10(22):239-42.
22
23
24
25
26 28. Saranga Bharathi R., V., Sharma, A., Chakladar, P., Kumari. Port exteriorisation
27 appendectomy-our experience. *Med J Armed Forces India*. 2011;67(2):147-51.
28
29
30
31
32 29. Biçakci U., B., Tander, M., Günaydin, R., Rizalar, E., Aritürk, S.H., Ayyildiz, et al. The
33 comparison of open and laparoscopic appendectomy: Is there any outcome difference
34 between non-complicated and complicated appendicitis? *Balkan Med J*. 2011;28(3):304-6.
35
36
37
38
39 30. Bickel A., M., Gurevits, R., Vamos, S., Ivry, A., Eitan. Perioperative hyperoxygenation and
40 wound site infection following surgery for acute appendicitis : A randomized, prospective,
41 controlled trial. *Arch Surg*. 2011;146(4):464-70.
42
43
44
45
46 31. Blackwood B.P., C.D., Gause, J.C., Harris, C.M., Theodorou, I., Helenowski, T.B., Lautz,
47 et al. Overweight and obese pediatric patients have an increased risk of developing a
48 surgical site infection. *Surg Infect*. 2017;18(4):491-7.
49
50
51
52
53
54 32. Blakely M.L., R., Williams, M.S., Dassinger, J.W., Eubanks III, P., Fischer, E.Y., Huang,
55 et al. Early vs interval appendectomy for children with perforated appendicitis. *Arch Surg*.
56 2011;146(6):660-5.
57
58
59
60

- 1
2
3 33. Bonadio W., K., Rebillot, O., Ukwuoma, C., Saracino, A., Iskhakov. Management of
4 Pediatric Perforated Appendicitis: Comparing Outcomes Using Early Appendectomy
5 versus Solely Medical Management. *Pediatr Infect Dis J.* 2017;36(10):937-41.
6
7
8
9
10
11 34. Boomer L.A., J.N., Cooper, K.J., Deans, P.C., Minneci, K., Leonhart, K.A., Diefenbach, et
12 al. Does delay in appendectomy affect surgical site infection in children with appendicitis?
13 *J Pediatr Surg.* 2014;49(6):1026-9.
14
15
16
17
18 35. Boomer L.A., J.N., Cooper, S., Anandalwar, S.C., Fallon, D., Ostlie, C.M., Leys, et al.
19 Delaying appendectomy does not lead to higher rates of surgical site infections. *Ann Surg.*
20 2016;264(1):164-8.
21
22
23
24
25
26 36. Bozkurt M.A., M.G., Ünsal, S., Kapan, B., Kankaya, M.U., Kalaycii, H., Aliiş. Two
27 different methods for appendiceal stump closure: Metal clip and Hem-o-lok clip. *J*
28 *Laparoendosc Adv Surg Techn.* 2014;24(8):571-3.
29
30
31
32
33 37. Brandt C., U., Hott, D., Sohr, F., Daschner, P., Gastmeier, H., Rüden. Operating room
34 ventilation with laminar airflow shows no protective effect on the surgical site infection rate
35 in orthopedic and abdominal surgery. *Ann Surg.* 2008;248(5):695-700.
36
37
38
39
40
41 38. Cairo S.B., M.V., Raval, M., Browne, H., Meyers, D.H., Rothstein. Association of same-
42 day discharge with hospital readmission after appendectomy in pediatric patients. *JAMA*
43 *Surg.* 2017;152(12):1106-12.
44
45
46
47
48 39. Cameron D.B., P., Melvin, D.A., Graham, C.C., Glass, S.K., Serres, M.P., Kronman, et al.
49 Extended Versus Narrow-spectrum Antibiotics in the Management of Uncomplicated
50 Appendicitis in Children: A Propensity-matched Comparative Effectiveness Study. *Ann*
51 *Surg.* 2017.
52
53
54
55
56
57 40. Cao J.-G., F., Tao, X.-J., Zhou, X.-G., Wang, S.-S., Wang, H., Zhang, et al. Trends and
58 outcomes of laparoscopic appendectomy in China: A multicenter, retrospective cohort
59
60

- 1
2
3 study. *Surg Pract.* 2015;19(4):166-72.
4
5
6
7
8 41. Cervantes-Sánchez C.R., R., Gutiérrez-Vega, J.A., Vázquez-Carpizo, P., Clark, C., Athié-
9 Gutiérrez. Syringe pressure irrigation of subdermic tissue after appendectomy to decrease
10 the incidence of postoperative wound infection. *World J Surg.* 2000;24(1):38-42.
11
12
13
14
15 42. Chamisa I. A clinicopathological review of 324 appendices removed for acute appendicitis
16 in Durban, South Africa: a retrospective analysis. *Ann R Coll Surg Engl.* 2009;91(8):688-
17 92.
18
19
20
21
22 43. Chaudhary I.A., Samiullah, A.A., Mallhi, Z., Afridi, A., Bano. Is it necessary to invaginate
23 the stump after appendectomy? *Pak J Med Sci.* 2005;21(1):35-8.
24
25
26
27
28 44. Chen D., H., Shi, H., Dong, K., Liu, K., Ding. Gasless single-incision laparoscopic
29 appendectomy. *Surg Endosc Interv Tech.* 2011;25(5):1472-6.
30
31
32
33
34 45. Chen C.-Y., Y.-C., Chen, H.-N., Pu, C.-H., Tsai, W.-T., Chen, C.-H., Lin. Bacteriology of
35 acute appendicitis and its implication for the use of prophylactic antibiotics. *Surg Infect.*
36 2012;13(6):383-90.
37
38
39
40
41 46. Chen C.-C., C.-T., Ting, M.-J., Tsai, W.-C., Hsu, P.-C., Chen, M.-D., Lee, et al.
42 Appendectomy timing: Will delayed surgery increase the complications? *J Chin Med*
43 *Assoc.* 2015;78(7):395-9.
44
45
46
47
48 47. Chiang R.-A., S.-L., Chen, Y.-C., Tsai, M.-J., Bair. Comparison of primary wound closure
49 versus open wound management in perforated appendicitis. *J Formos Med Assoc.*
50 2006;105(10):791-5.
51
52
53
54
55
56 48. Chiang R.-A., S.-L., Chen, Y.-C., Tsai. Delayed primary closure versus primary closure for
57 wound management in perforated appendicitis: A prospective randomized controlled trial.
58 *J Chin Med Assoc.* 2012;75(4):156-9.
59
60

- 1
2
3
4
5
6 49. Cho M., J., Kang, I.-K., Kim, K.Y., Lee, S.-K., Sohn. Underweight body mass index as a
7 predictive factor for surgical site infections after laparoscopic appendectomy. *Yonsei Med*
8 *J.* 2014;55(6):1611-6.
9
10
11
12
13 50. Choudhary S.K., S.K., Dhakaita. Appendicular mass-early appendicectomy vs interval
14 appendicectomy. *Intl J Pharma Bio Sci.* 2014;5(1):B400-B4.
15
16
17
18
19 51. Clyde C., T., Bax, A., Merg, M., MacFarlane, P., Lin, S., Beyersdorf, et al. Timing of
20 intervention does not affect outcome in acute appendicitis in a large community practice.
21 *Am J Surg.* 2008;195(5):590-3.
22
23
24
25
26 52. Coakley B.A., E.S., Sussman, T.S., Wolfson, A.S., Bhagavath, J.J., Choi, N.E., Ranasinghe,
27 et al. Postoperative antibiotics correlate with worse outcomes after appendectomy for
28 nonperforated appendicitis. *J Am Coll Surg.* 2011;213(6):778-83.
29
30
31
32
33
34 53. Crandall M., M.B., Shapiro, M., Worley, M.A., West. Acute uncomplicated appendicitis:
35 case time of day influences hospital length of stay. *Surg Infect (Larchmt).* 2009;10(1):65-
36 9.
37
38
39
40
41
42 54. Dede K., T., Mersich, A., Zaránd, I., Besznyák, Z., Baranyai, B., Atkári, et al. Laparoscopic
43 or open appendectomy? *Laparoszkópos vagy nyílt appendectomia?* 2008;149(50):2357-61.
44
45
46
47
48 55. Dhiman N., A., Chi, T.M., Pawlik, D.T., Efron, E.R., Haut, E.B., Schneider, et al. Increased
49 complications after appendectomy in patients with cerebral palsy: Are special needs patients
50 at risk for disparities in outcomes? *Surgery.* 2013;154(3):479-85.
51
52
53
54
55 56. Dimitriou I., B., Reckmann, O., Nephuth, M., Betzler. Single institution's experience in
56 laparoscopic appendectomy as a suitable therapy for complicated appendicitis.
57 *Langenbeck's Arch Surg.* 2013;398(1):147-52.
58
59
60

- 1
2
3 57. Durkin M.J., K.V., Dicks, A.W., Baker, S.S., Lewis, R.W., Moehring, L.F., Chen, et al.
4 Seasonal variation of common surgical site infections: Does season matter? *Infect Control*
5 *Hosp Epidemiol.* 2015;36(9):1011-6.
6
7
8
9
10
11 58. Ein S.H., A., Nasr, A., Ein. Open appendectomy for pediatric ruptured appendicitis: a
12 historical clinical review of the prophylaxis of wound infection and postoperative intra-
13 abdominal abscess. *Can J Surg.* 2013;56(3):E7-E12.
14
15
16
17
18 59. Fukuda H. Patient-related risk factors for surgical site infection following eight types of
19 gastrointestinal surgery. *J Hosp Infect.* 2016;93(4):347-54.
20
21
22
23
24 60. Gandaglia G., K.R., Ghani, A., Sood, J.R., Meyers, J.D., Sammon, M., Schmid, et al. Effect
25 of minimally invasive surgery on the risk for surgical site infections results from the national
26 surgical quality improvement program (nsqip) database. *JAMA Surg.* 2014;149(10):1039-
27 44.
28
29
30
31
32
33 61. Garcell H.G., A.V., Arias, C.A., Pancorbo Sandoval, E.G., García, M.E., Valle Gamboa,
34 A.B., Sado, et al. Incidence and etiology of surgical site infections in appendectomies: A 3-
35 year prospective study. *Oman Med J.* 2017;32(1):31-5.
36
37
38
39
40
41 62. Ghnnam W.M. Elderly versus young patients with appendicitis 3 years experience. *Alex J*
42 *Med.* 2012;48(1):9-12.
43
44
45
46
47 63. Giesen L.J., A.L., van den Boom, C.C., van Rossem, P.T., den Hoed, B.P., Wijnhoven.
48 Retrospective Multicenter Study on Risk Factors for Surgical Site Infections after
49 Appendectomy for Acute Appendicitis. *Dig Surg.* 2017;34(2):103-7.
50
51
52
53
54
55 64. Giiti G.C., H.D., Mazigo, J., Heukelbach, W., Mahalu. HIV, appendectomy and
56 postoperative complications at a reference hospital in Northwest Tanzania: Cross-sectional
57 study. *AIDS Res Ther.* 2010;7.
58
59
60

- 1
2
3 65. Golub A.V., R.S., Kozlov, V.G., Pleshkov, A.P., Moskalev, R.A., Alibegov, M.A.,
4 Chelombitko. Surgical Site Infections after Open Appendectomy and Effectiveness of
5 Complex Approach to Their Prevention. *Khirurgiia (Mosk)*. 2016(6):68-76.
6
7
8
9
10
11 66. Gross T.S., C., McCracken, K.F., Heiss, M.L., Wulkan, M.V., Raval. The contribution of
12 practice variation to length of stay for children with perforated appendicitis. *J Pediatr Surg*.
13 2016;51(8):1292-7.
14
15
16
17
18 67. Gurien L.A., D.L., Wyrick, S.D., Smith, M.S., Dassinger. Optimal timing of appendectomy
19 in the pediatric population. *J Surg Res*. 2016;202(1):126-31.
20
21
22
23
24 68. Hamzaoglu I, B., Baca, D.E., Böler, E., Polat, Y., Özer. Is umbilical flora responsible for
25 wound infection after laparoscopic surgery? *Surg Laparoscopy Endosc Percutaneous Tech*.
26 2004;14(5):263-7.
27
28
29
30
31
32 69. Harmon L.A., M.L., Davis, D.C., Jupiter, R.C., Frazee, J.L., Regner. Computed tomography
33 to operating room in less than 3 hours minimizes complications from appendicitis. *Am J*
34 *Surg*. 2016;212(2):246-50.
35
36
37
38
39 70. Helling T.S., D.F., Soltys, S., Seals. Operative versus non-operative management in the care
40 of patients with complicated appendicitis. *Am J Surg*. 2017;214(6):1195-200.
41
42
43
44
45 71. Helmer K.S., E.K., Robinson, K.P., Lally, J.C., Vasquez, K.L., Kwong, T.H., Liu, et al.
46 Standardized patient care guidelines reduce infectious morbidity in appendectomy patients.
47 *Am J Surg*. 2002;183(6):608-13.
48
49
50
51
52 72. Hesami M.A., H., Alipour, H., Nikoupour Daylami, B., Alipour, S., Bazargan-Hejazi, A.,
53 Ahmadi. Irrigation of abdomen with imipenem solution decreases surgical site infections in
54 patients with perforated appendicitis: A randomized clinical trial. *Iran Red Crescent MedJ*.
55 2014;16(4).
56
57
58
59
60

- 1
2
3 73. Horvath P., J., Lange, R., Bachmann, F., Struller, A., Königsrainer, M., Zdichavsky.
4 Comparison of clinical outcome of laparoscopic versus open appendectomy for complicated
5 appendicitis. *Surg Endosc Interv Tech.* 2017;31(1):199-205.
6
7
8
9
10
11 74. Hughes M.J., E., Harrison, S., Paterson-Brown. Post-operative antibiotics after
12 appendectomy and post-operative abscess development: A retrospective analysis. *Surg*
13 *Infect.* 2013;14(1):56-61.
14
15
16
17 75. Hussain M.I., M.K., Alam, H.H., Al-Qahatani, M.H., Al-Akeely. Role of postoperative
18 antibiotics after appendectomy in non-perforated appendicitis. *J Coll Phys Surg Pak.*
19 2012;22(12):756-9.
20
21
22
23
24
25
26 76. Iqbal M., M., Jawaid, A., Qureshi, S., Iqbal. Effect of povidone-iodine irrigation on post
27 appendectomy wound infection: Randomized control trial. *J Postgrad Med Inst.*
28 2015;29(3):160-4.
29
30
31
32
33 77. Iñigo J.J., B., Bermejo, B., Oronoz, J., Herrera, A., Tarifa, F., Pérez, et al. Surgical site
34 infection in general surgery: 5-year analysis and assessment of the National Nosocomial
35 Infection Surveillance (NNIS) index. *Infección de sitio quirúrgico en un servicio de cirugía*
36 *general Análisis de cinco años y valoración del índice National Nosocomial Infection*
37 *Surveillance (NNIS).* 2006;79(4):224-30.
38
39
40
41
42
43
44 78. Javadi S.M.R., S.Y., Zarghami, P., Ghaderzadeh, M., Ghorbanpoor, H.R., Makarchian, A.,
45 Derakhshanfar, et al. Comparison of small access and classic McBurney's incisions for open
46 appendectomy: A randomized controlled trial. *Shiraz E Med J.* 2017;18(10).
47
48
49
50
51 79. Jenkins P.C., M.K., Oerline, A.J., Mullard, M.J., Englesbe, D.A., Campbell, M.R.,
52 Hemmila. Hospital variation in outcomes following appendectomy in a regional quality
53 improvement program. *Am J Surg.* 2016;212(5):857-62.
54
55
56
57
58
59 80. Kang J., B.N., Bae, G., Gwak, I., Park, H., Cho, K., Yang, et al. Comparative study of a
60

1
2
3 single-incision laparoscopic and a conventional laparoscopic appendectomy for the
4 treatment of acute appendicitis. *J Korean Soc Coloproctology*. 2012;28(6):304-8.
5
6
7
8

- 9 81. Kapischke M., A., Pries, A., Caliebe. Short term and long term results after open vs.
10 Laparoscopic appendectomy in childhood and adolescence: A subgroup analysis. *BMC*
11 *Pediatr*. 2013;13(1).
12
13
14
15
16 82. Karam P.A., A., Hiuser, D., Magnuson, F.G.F., Seifarth. Intracorporeal hybrid single port
17 vs conventional laparoscopic appendectomy in children. *Pediatr Med Chir*. 2016;38(3):89-
18 92.
19
20
21
22
23
24 83. Karam P.A., A., Mohan, M.R., Buta, F.G., Seifarth. Comparison of Transumbilical
25 Laparoscopically Assisted Appendectomy to Conventional Laparoscopic Appendectomy in
26 Children. *Surg Laparoscopy Endosc Percutaneous Tech*. 2016;26(6):508-12.
27
28
29
30
31 84. Kasatpibal N., S., Jamulitrat, V., Chongsuvivatwong. Standardized incidence rates of
32 surgical site infection: A multicenter study in Thailand. *Am J Infect Control*.
33 2005;33(10):587-94.
34
35
36
37
38
39 85. Kasatpibal N., M., Nørgaard, H.T., Sørensen, H.C., Schönheyder, S., Jamulitrat, V.,
40 Chongsuvivatwong. Risk of surgical site infection and efficacy of antibiotic prophylaxis: A
41 cohort study of appendectomy patients in Thailand. *BMC Infect Dis*. 2006;6.
42
43
44
45
46 86. Kato Y., T., Marusasa, S., Ichikawa, G.J., Lane, T., Okazaki, A., Yamataka. Lapprotector
47 use decreases incisional wound infections in cases of perforated appendicitis: a prospective
48 study. *Asian J Surg*. 2008;31(3):101-3.
49
50
51
52
53
54 87. Kell M.R., K., Power, D.C., Winter, C., Power, C., Shields, W.O., Kirwan, et al. Predicting
55 outcome after appendectomy. *Ir J Med Sci*. 2003;172(2):63-5.
56
57
58
59 88. Khan M.N., T., Fayyad, T.D., Cecil, B.J., Moran. Laparoscopic versus open appendectomy:
60

- 1
2
3 the risk of postoperative infectious complications. *JLS*. 2007;11(3):363-7.
4
5
6
7
8 89. Khan K.I., S., Mahmood, M., Akmal, A., Waqas. Comparison of rate of surgical wound
9 infection, length of hospital stay and patient convenience in complicated appendicitis
10 between primary closure and delayed primary closure. *J Pak Med Assoc*. 2012;62(8):596-
11 8.
12
13
14
15
16 90. Khan I., M.I., Khan, M., Jawed, U., Shaikh, S., Ahmed, A., Arif. To compare the frequency
17 of superficial surgical site infection after laparoscopic versus open appendectomy. *Med*
18 *Forum Monthly*. 2014;25(11):52-5.
19
20
21
22
23
24 91. Khiria L.S., R., Ardhnari, N., Mohan, P., Kumar, R., Nambiar. Laparoscopic
25 appendectomy for complicated appendicitis: Is it safe and justified? A retrospective
26 analysis. *Surg Laparoscopy Endosc Percutaneous Tech*. 2011;21(3):142-5.
27
28
29
30
31 92. Kılıç Ş.S., S., Ekinçi, İ., Karnak, A.Ö., Çiftçi, F.C., Tanyel, M.E., Şenocak. Drainage
32 systems' effect on surgical site infection in children with perforated appendicitis. *Drenaj*
33 *Sistemlerinin perforé apandisitli çocuklarda cerrahi alan enfeksiyonuna etkisi*.
34 2016;7(5):591-4.
35
36
37
38
39
40 93. Kim M.J., F.J., Fleming, D.D., Gunzler, S., Messing, R.M., Salloum, J.R.T., Monson.
41 Laparoscopic appendectomy is safe and efficacious for the elderly: An analysis using the
42 National Surgical Quality Improvement Project database. *Surg Endosc Interv Tech*.
43 2011;25(6):1802-7.
44
45
46
47
48
49 94. Kim J.H., H.Y., Kim, S.K., Park, J.S., Lee, D.S., Heo, S.W., Park, et al. Single-incision
50 laparoscopic appendectomy versus conventional laparoscopic appendectomy: Experiences
51 from 1208 cases of single-incision laparoscopic appendectomy. *Ann Surg*.
52 2015;262(6):1054-8.
53
54
55
56
57
58 95. Kim J.K., J., Kang, W.R., Kim, E.J., Park, S.H., Baik, K.Y., Lee. Does Conversion
59
60

- 1
2
3 Adversely Impact the Clinical Outcomes for Patients with Complicated Appendicitis? J
4 Laparoendosc Adv Surg Techn. 2016;26(8):635-40.
5
6
7
8
9 96. Kiriakopoulos A., D., Tsakayannis, D., Linos. Laparoscopic management of complicated
10 appendicitis. JSLS. 2006;10(4):453-6.
11
12
13
14
15 97. Kirshtein B., Z.H., Perry, S., Mizrahi, L., Lantsberg. Value of laparoscopic appendectomy
16 in the elderly patient. World J Surg. 2009;33(5):918-22.
17
18
19
20
21 98. Kiudelis M., P., Ignatavicius, K., Zviniene, S., Grizas. Analysis of intracorporeal knotting
22 with invaginating suture versus endoloops in appendiceal stump closure. Wideochir Inne
23 Tech Ma?oinwazyjne. 2013;8(1):69-73.
24
25
26
27
28 99. Kleif J., L., Rasmussen, S., Fonnes, P., Tibæk, A., Daoud, H., Lund, et al. Enteral
29 Antibiotics are Non-inferior to Intravenous Antibiotics After Complicated Appendicitis in
30 Adults: A Retrospective Multicentre Non-inferiority Study. World J Surg.
31 2017;41(11):2706-14.
32
33
34
35
36
37
38 100. Koizumi N., H., Kobayashi, Y., Nakase, T., Takagi, K., Fukumoto. Efficacy of
39 transumbilical laparoscopic-assisted appendectomy for appendicitis: a four-year experience
40 at a single center. Surg Today. 2015;45(10):1245-9.
41
42
43
44
45 101. Shimizu T., M., Ishizuka, K., Kubota. The preoperative serum C-reactive protein level
46 is a useful predictor of surgical site infections in patients undergoing appendectomy. Surg
47 Today. 2015;45(11):1404-10.
48
49
50
51
52
53 102. Kumamoto K., H., Imaizumi, N., Hokama, T., Ishiguro, K., Ishibashi, K., Baba, et al.
54 Recent trend of acute appendicitis during pregnancy. Surg Today. 2015;45(12):1521-6.
55
56
57
58 103. Kumar B., A., Samad, T.W., Khanzada, M.H., Laghari, A.R., Shaikh. Superiority of
59 laparoscopic appendectomy over open appendectomy: The Hyderabad experience. Rawal
60

- 1
2
3 Med J. 2008;33(2):165-8.
4
5
6
7
8 104. Kumar S., A., Jalan, B.N., Patowary, S., Shrestha. Laparoscopic appendectomy versus
9 open appendectomy for acute appendicitis: A prospective comparative study. Kathmandu
10 Univ Med J. 2016;14(55):244-8.
11
12
13
14
15 105. Lacher M., O.J., Muensterer, G.R., Yannam, C.J., Aprahamian, L., Perger, M., Megison,
16 et al. Feasibility of single-incision pediatric endosurgery for treatment of appendicitis in
17 415 children. J Laparoendosc Adv Surg Techn. 2012;22(6):604-8.
18
19
20
21
22 106. Lasses-Martínez B., E., Ortiz-Oshiro, J.L., Cabañas-Ojeda, P., Benito-Expósito, C.,
23 Fernández-Pérez, J., Alvarez Fernández-Represa. Cost is not a drawback to perform
24 laparoscopic appendectomy in an academic hospital. Surg Laparoscopy Endosc
25 Percutaneous Tech. 2014;24(4):e123-e7.
26
27
28
29
30
31
32 107. Le D., W., Rusin, B., Hill, J., Langell. Post-operative antibiotic use in nonperforated
33 appendicitis. Am J Surg.2009;198(6):748-52.
34
35
36
37
38
39 108. Lee P., K., Waxman, B., Taylor, S., Yim. Use of wound-protection system and
40 postoperative wound-infection rates in open appendectomy: A randomized prospective
41 trial. Arch Surg. 2009;144(9):872-5.
42
43
44
45
46 109. Lee S.L., S., Shekherdimian, V.Y., Chiu. Comparison of pediatric appendicitis
47 outcomes between teaching and nonteaching hospitals. J Pediatr Surg. 2010;45(5):894-7.
48
49
50
51 110. Lee J.A., K.Y., Sung, J.H., Lee, D.S., Lee. Laparoscopic appendectomy with a single
52 incision in a single institute. J Korean Soc Coloproctology. 2010;26(4):260-4.
53
54
55
56
57 111. Lee S.-Y., H.-M., Lee, C.-S., Hsieh, J.-H., Chuang. Transumbilical laparoscopic
58 appendectomy for acute appendicitis: A reliable one-port procedure. Surg Endosc Interv
59
60

- 1
2
3 Tech. 2011;25(4):1115-20.
4
5
6
7 112. Lee S.L., A., Yaghoubian, A., Kaji. Laparoscopic vs open appendectomy in children:
8 Outcomes comparison based on age, sex, and perforation status. Arch Surg.
9 2011;146(10):1118-21.
10
11
12
13
14
15 113. Lee S.M., G.S., Hwang, D.S., Lee. Single-incision laparoscopic appendectomy using
16 homemade glove port at low cost. J Minimal Access Surg. 2016;12(2):124-8.
17
18
19
20
21 114. Levy S.M., G., Holzmann-Pazgal, K.P., Lally, K., Davis, L.S., Kao, K., Tsao. Quality
22 check of a quality measure: Surgical wound classification discrepancies impact risk-
23 stratified surgical site infection rates in pediatric appendicitis. J Am Coll Surg.
24 2013;217(6):969-73.
25
26
27
28
29
30 115. Li P., Q., Xu, Z., Ji, Y., Gao, X., Zhang, Y., Duan, et al. Comparison of surgical stress
31 between laparoscopic and open appendectomy in children. J Pediatr Surg. 2005;40(8):1279-
32 83.
33
34
35
36
37
38
39 116. Lim S.G., E.J., Ahn, S.Y., Kim, I.Y., Chung, J.-M., Park, S.H., Park, et al. A clinical
40 comparison of laparoscopic versus open appendectomy for complicated appendicitis. J
41 Korean Soc Coloproctology. 2011;27(6):293-7.
42
43
44
45
46 117. Geraldo José de Souza Lima, Silva, Alcino Lázaro da, Castro, Eduardo Godoy, Abras,
47 Gustavo Munayer, Pires, Lívio José Suretti, Leite, Rodrigo Fabiano Guedes. Efetividade e
48 segurança da apendicectomia videoassistida em porta única transumbilical em adolescentes
49 e adultos X1 Effectiveness and safeness of single-port trans-umbilical laparoscopic
50 appendectomy done in adolescents and adults. Revista do Colégio Brasileiro de Cirurgiões.
51 2008;35(4):244-51.
52
53
54
55
56
57
58
59 118. Lin H.-F., J.-M., Wu, L.-M., Tseng, K.-H., Chen, S.-H., Huang, I.-R., Lai. Laparoscopic
60

- 1
2
3 Versus Open Appendectomy for Perforated Appendicitis. *J Gastrointest Surg.*
4 2006;10(6):906-10.
5
6
7
8
9 119. Litz C.N., S.M., Farach, P.D., Danielson, N.M., Chandler. Obesity and single-incision
10 laparoscopic appendectomy in children. *J Surg Res.* 2016;203(2):283-6.
11
12
13
14 120. Liu C., W., Wang, Y., Sun, M., Xu, H., Zhuang, H., Chen, et al. Efficacy and
15 complications of laparoscopic appendectomy for pediatric appendicitis. *Int J Clin Exp Med.*
16 2017;10(9):13784-9.
17
18
19
20
21
22 121. Mahmood M.M., A., Shahab, M.A., Razzaq. Surgical site infection in open versus
23 laparoscopic appendectomy. *Pak J Med Health Sci.* 2016;10(3):1076-8.
24
25
26
27
28 122. Mbah N., W.Ek., Opara, N.P., Agwu. Waiting time among acute abdominal
29 emergencies in a Nigerian teaching hospital: Causes of delay and consequences. *Niger J*
30 *Surg Res.* 2006;8(1):69-73.
31
32
33
34
35
36 123. Memon G.A., A.I., Memon, S.K.A., Shah, R.A., Sahito, Habib-Ur-Rehman, S., Leghari,
37 et al. An experience of treatment outcome in acute appendicitis with antibiotics and
38 appendectomy at a tertiary care hospital. *Med Forum Monthly.* 2017;28(3):136-40.
39
40
41
42
43 124. Menezes M., L., Das, M., Alagtal, J., Haroun, P., Puri. Laparoscopic appendectomy is
44 recommended for the treatment of complicated appendicitis in children. *Pediatr Surg Int.*
45 2008;24(3):303-5.
46
47
48
49
50 125. Merenda M., A., Litarski, P., Kabziński, D., Janczak. Laparoscopic appendectomy as an
51 alternative to conventional procedure - results in our own material. *Pol Przegl Chir.*
52 2013;85(6):323-8.
53
54
55
56
57
58 126. Michailidou M., M.G., Sacco Casamassima, S.D., Goldstein, C., Gause, O., Karim, J.H.,
59 Salazar, et al. The impact of obesity on laparoscopic appendectomy: Results from the ACS
60

- 1
2
3 National Surgical Quality Improvement Program pediatric database. *J Pediatr Surg.*
4 2015;50(11):1880-4.
5
6
7
8
9 127. Michailidou M., S.D., Goldstein, M.G., Sacco Casamassima, J.H., Salazar, R., Elliott,
10 J., Hundt, et al. Laparoscopic versus open appendectomy in children: The effect of surgical
11 technique on healthcare costs. *Am J Surg.* 2015;210(2):270-5.
12
13
14
15
16 128. Mickovic I.N., Z., Golubovic, S., Mickovic, D., Vukovic, S., Trajkovic, S.S.,
17 Antunovic, et al. A comparative analysis of laparoscopic appendectomy in relation to the
18 open appendectomy in children. *Uporedna analiza laparoscopske apendektomije u odnosu*
19 *na otvorenu apendektomiju kod dece.* 2016;17(1):49-53.
20
21
22
23
24
25
26 129. Ming P.C., T.Y., Yee Yan, L.H., Tat. Risk factors of postoperative infections in adults
27 with complicated appendicitis. *Surg Laparoscopy Endosc Percutaneous Tech.*
28 2009;19(3):244-8.
29
30
31
32
33 130. Mingmalairak C., P., Ungbhakorn, V., Paocharoen. Efficacy of antimicrobial coating
34 suture coated polyglactin 910 with tricosan (Vicryl Plus) compared with polyglactin 910
35 (Vicryl) in reduced surgical site infection of appendicitis, double blind randomized control
36 trial, preliminary safety report. *J Med Assoc Thailand.* 2009;92(6):770-5.
37
38
39
40
41
42
43 131. Miyano G., T., Okazaki, Y., Kato, T., Marusasa, T., Takahashi, G.J., Lane, et al. Open
44 versus laparoscopic treatment for pan-peritonitis secondary to perforated appendicitis in
45 children: A prospective analysis. *J Laparoendosc Adv Surg Techn.* 2010;20(7):655-7.
46
47
48
49
50 132. Moazzez A., R.J., Mason, N., Katkhouda. Thirty-day outcomes of laparoscopic versus
51 open appendectomy in elderly using ACS/NSQIP database. *Surg Endosc Interv Tech.*
52 2013;27(4):1061-71.
53
54
55
56
57 133. Mohammad Taghi Rajabi-Mashhadi, Mousavi, Seyed Hadi, Khosravi-Mashizi, M. H.,
58 Ghayour-Mobarhan, Majid, Sahebkar, Amirhossein. Optimum duration of perioperative
59
60

- 1
2
3 antibiotic therapy in patients with acute non-perforated appendicitis: a prospective
4 randomized trial. *Asian Biomedicine*. 2012;6(6):891-4.
5
6
7
- 8
9 134. Monge Jodra V., A., Robustillo Rodela, F., Martin Martinez, N., López Fresneña, S.,
10 Oña Compán, F., Calbo Torrecillas, et al. Standardized infection ratios for three general
11 surgery procedures: A comparison between Spanish hospitals and U.S. centers participating
12 in the national nosocomial infections surveillance system. *Infect Control Hosp Epidemiol*.
13 2003;24(10):744-8.
14
15
16
17
18
- 19
20 135. Mueck K.M., L.R., Putnam, K.T., Anderson, K.P., Lally, K., Tsao, L.S., Kao. Does
21 compliance with antibiotic prophylaxis in pediatric simple appendicitis matter? *J Surg Res*.
22 2017;216:1-8.
23
24
25
26
27
- 28
29 136. Muensterer O.J., C., Puga Nougues, O.O., Adibe, S.R., Amin, K.E., Georgeson, C.M.,
30 Harmon. Appendectomy using single-incision pediatric endosurgery for acute and
31 perforated appendicitis. *Surg Endosc Interv Tech*. 2010;24(12):3201-4.
32
33
34
35
- 36
37 137. Muensterer O.J., R., Keijzer. A simple vacuum dressing reduces the wound infection
38 rate of single-incision pediatric endosurgical appendectomy. *J Soc Laparoendoscopic Surg*.
39 2011;15(2):147-50.
40
41
42
- 43
44 138. Mustafa M.I.T., S.M., Chaudhry, R.I.T., Mustafa. Comparison of early outcome
45 between patients of open appendectomy with and without drain for perforated appendicitis.
46 *Pak J Med Health Sci*. 2016;10(3):890-3.
47
48
49
- 50
51 139. Nadler E.P., K.K., Reblock, H.R., Ford, B.A., Gaines. Monotherapy versus multi-drug
52 therapy for the treatment of perforated appendicitis in children. *Surg Infect*. 2003;4(4):327-
53 33.
54
55
56
57
- 58
59 140. Nataraja R.M., A., Bandi, S.A., Clarke, M.J., Haddad. Comparison of intra-abdominal
60 abscess formation following laparoscopic and open appendectomy in children. *J*

- Laparoendosc Adv Surg Techn. 2010;20(4):391-4.
141. Nataraja R.M., W.J., Teague, J., Galea, L., Moore, M.J., Haddad, T., Tsang, et al. Comparison of intraabdominal abscess formation after laparoscopic and open appendectomies in children. *J Pediatr Surg.* 2012;47(2):317-21.
142. Norton Pérez, Romero, Marcela, Castelblanco, María Isabel, Rodríguez, Emma Isabel. Infección del sitio operatorio de apendicectomías en un hospital de la orinoquia colombiana X1 Surgical site infection following appendectomy at a hospital in the Colombian Orinoco river basin (Colombian Orinoquia). *Revista Colombiana de Cirugía.* 2009;24(1):23-30.
143. Obayashi J., K., Ohyama, S., Manabe, K., Tanaka, H., Nagae, H., Shima, et al. Are there reliable indicators predicting post-operative complications in acute appendicitis? *Pediatr Surg Int.* 2015;31(12):1189-93.
144. Obinwa O., C., Peirce, M., Cassidy, T., Fahey, J., Flynn. A model predicting perforation and complications in paediatric appendectomy. *Int J Colorectal Dis.* 2015;30(4):559-65.
145. Ohene-Yeboah M., B., Togbe. An audit of appendicitis and appendectomy in Kumasi, Ghana. *West Afr J Med.* 2006;25(2):138-43.
146. Okkyung Suh, 신완균, 강성희, 양대현. Appropriate Duration of Prophylactic Antibiotics in Acute Nonperforated Appendicitis Z1 급성 비천공성 충수염 수술시 예방적 항균제의 사용기간. *Korean Journal of Clinical Pharmacy S1 한국임상약학회지.* 2002;12(2):65-70.
147. Francisco Gabriel Onieva, Roldán, Sara, Domínguez, José Ramón, Montero, Juan Pedro, Galnares, Alfonso, Peralta, Jordi. Abordaje laparoscópico frente a enfoque clásico

- 1
2
3 en el tratamiento de la apendicitis aguda X1 Laparoscopic approach versus classic open
4 procedure in the treatment of acute appendicitis. *Revista Colombiana de Cirugía*.
5 2017;32(1):26-31.
6
7
8
9
- 10 148. Page A.J., J.D., Pollock, S., Perez, S.S., Davis, E., Lin, J.F., Sweeney. Laparoscopic
11 Versus Open Appendectomy: An Analysis of Outcomes in 17,199 Patients Using
12 ACS/NSQIP. *J Gastrointest Surg*. 2010;14(12):1955-62.
13
14
15
16
- 17 149. Palesty J.A., X.J., Wang, R.C., Rutland, J., Leighton, S.J., Dudrick, A., Benbrahim.
18 Fifty-five consecutive laparoscopic appendectomy procedures without conversion. *JSL*.
19 2004;8(2):141-5.
20
21
22
23
24
- 25 150. Pandit R.K. Safe and feasible time limit for early appendectomy in appendiceal mass.
26 *Kathmandu Univ Med J*. 2016;14(55):210-4.
27
28
29
30
- 31 151. Parcells J.P., J.P., Mileski, F.T., Gnagy, A.F., Haragan, W.J., Mileski. Using
32 antimicrobial solution for irrigation in appendicitis to lower surgical site infection rates. *Am*
33 *J Surg*. 2009;198(6):875-80.
34
35
36
37
38
39
40
- 41 152. Park H.-C., M.J., Kim, B.H., Lee. Effect of a Standardized Protocol of Antibiotic
42 Therapy on Surgical Site Infection after Laparoscopic Surgery for Complicated
43 Appendicitis. *Surg Infect*. 2017;18(6):684-8.
44
45
46
47
48
- 49 153. Seongmun Park, Park, Min-Su, Lee, Kil-Yeon. Relationship between the Hospital Visit-
50 to-Operation Time Interval and the Risk of Appendiceal Perforation and Clinical Outcomes.
51 *Journal of Minimally Invasive Surgery*. 2018;21(1):31-7.
52
53
54
- 55 154. Reoyo Pascual J.F., R., León Miranda, C., Cartón Hernández, E., Alonso Alonso, R.M.,
56 Martínez Castro, J., Sánchez Manuel. Laparoscopic appendectomy by 'glove port'
57 system: Our first 100 cases. *Apendicectomía laparoscópica por sistema «glove port»:*
58
59
60

- 1
2
3 nuestros primeros 100 casos. 2017;69(6):467-71.
4
5
6
7 155. Patrice Lemieux., Pascal Rheaume., Isabelle Levesque., Emmanuel Bujold., Gaetan
8 Brochu. Laparoscopic appendectomy in pregnant patients: a review of 45 cases. Surg
9 Endosc.2009; 23:1701.
10
11
12
13
14 156. Patel S.C., G.F., Jumba, S., Akmal. Laparoscopic appendicectomy at the Aga Khan
15 Hospital, Nairobi. East Afr Med J. 2003;80(9):447-51.
16
17
18
19
20 157. Percy C., K., Almahmoud, T., Jackson, C., Hartline, A., Cahill, L., Spence, et al. Risky
21 business? Investigating outcomes of patients undergoing urgent laparoscopic
22 appendectomy on antithrombotic therapy. Am J Surg. 2017;214(6):1012-5.
23
24
25
26
27 158. Pishori T., A.R., Siddiqui, M., Ahmed. Surgical wound infection surveillance in general
28 surgery procedures at a teaching hospital in Pakistan. Am J Infect Control. 2003;31(5):296-
29 301.
30
31
32
33
34
35 159. Putnam L.R., T.G., Ostovar-Kermani, A., Le Blanc, K.T., Anderson, G., Holzmann-
36 Pazgal, K.P., Lally, et al. Surgical site infection reporting: more than meets the agar. J
37 Pediatr Surg. 2017;52(1):156-60.
38
39
40
41
42
43 160. Al-Qahtani S.M., H.M., Al-Amoudi, S., Al-Jehani, A.S., Ashour, M.R., Abd-Hammad,
44 O.R., Tawfik, et al. Post-appendectomy surgical site infection rate after using an
45 antimicrobial film incise drape: A prospective study. Surg Infect. 2015;16(2):155-8.
46
47
48
49
50 161. Quezada F., N., Quezada, R., Mejia, A., Brañes, O., Padilla, N., Jarufe, et al.
51 Laparoscopic versus open approach in the management of appendicitis complicated
52 exclusively with peritonitis: A single center experience. Int J Surg. 2015;13:80-3.
53
54
55
56
57
58 162. Raakow J., H.-G., Liesaus, P., Neuhaus, R., Raakow. Single-incision versus multiport
59 laparoscopic appendectomy: a case-matched comparative analysis. Surg Endosc Interv
60

- 1
2
3 Tech. 2015;29(6):1530-6.
4
5
6
7 163. Ríos J., C., Murillo, G., Carrasco, C., Humet. Increase in costs attributable to surgical
8 infection after appendicectomy and colectomy. Incremento de costes atribuible a la
9 infección quirúrgica de la apendicectomía y colectomía. 2003;17(3):218-25.
10
11
12
13
14 164. Rafiq M.S., M.M., Khan, A., Khan, H., Jan. Evaluation of postoperative antibiotics after
15 non-perforated appendectomy. J Pak Med Assoc. 2015;65(8):815-7.
16
17
18
19
20
21
22 165. Reinisch A., J., Heil, G., Woeste, W., Bechstein, J., Liese. The meteorological influence
23 on seasonal alterations in the course of acute appendicitis. J Surg Res. 2017;217:137-43.
24
25
26
27
28 166. Romano A., P., Parikh, P., Byers, N., Namias. Simple acute appendicitis versus non-
29 perforated gangrenous appendicitis: Is there a difference in the rate of post-operative
30 infectious complications? Surg Infect. 2014;15(5):517-20.
31
32
33
34
35
36 167. Romel Hilaire, Fernández, Zenén Rodríguez, García, Lázaro Ibrahim Romero, Sánchez,
37 Luis Pablo Rodríguez. Apendicectomía videolaparoscópica frente a apendicectomía
38 convencional X1 Laparoscopic versus conventional appendectomy. Revista Cubana de
39 Cirugía. 2014;53(1):30-40.
40
41
42
43
44
45 168. Romy S., M.-C., Eisenring, V., Bettschart, C., Petignat, P., Francioli, N., Troillet.
46 Laparoscope use and surgical site infections in digestive surgery. Ann Surg.
47 2008;247(4):627-32.
48
49
50
51
52
53 169. Rooh-ul-Muqim, M., Khan, M., Zarin. Experience of laparoscopic appendectomies
54 versus open appendectomies. Pak J Med Sci. 2010;26(2):324-8.
55
56
57
58
59 170. Van Rossem C.C., M.H.F., Schreinemacher, K., Treskes, R.M., Van Hogeand,
60 A.A.W., Van Geloven. Duration of antibiotic treatment after appendicectomy for acute

- 1
2
3 complicated appendicitis. *Br J Surg.* 2014;101(6):715-9.
4
5
6
7
8 171. Van Rossem C.C., M.D., Bolmers, M.H., Schreinemacher, A.A., van Geloven, W.A.,
9 Bemelman. Prospective nationwide outcome audit of surgery for suspected acute
10 appendicitis. *Br J Surg.* 2016;103(1):144-51.
11
12
13
14 172. Viet Hung N., T., Anh Thu, V.D., Rosenthal, D., Tat Thanh, N., Quoc Anh, N., Le Bao
15 Tien, et al. Surgical site infection rates in seven cities in Vietnam: Findings of the
16 international nosocomial infection control consortium. *Surg Infect.* 2016;17(2):243-9.
17
18
19
20
21 173. Rotermann M. Infection after cholecystectomy, hysterectomy or appendectomy. *Health*
22 *Rep.* 2004;15(4):11-23.
23
24
25
26 174. Saar S., P., Talving, J., Laos, T., Põdrämägi, M., Sokirjanski, T., Lustenberger, et al.
27 Delay Between Onset of Symptoms and Surgery in Acute Appendicitis Increases
28 Perioperative Morbidity: A Prospective Study. *World J Surg.* 2016;40(6):1308-14.
29
30
31
32
33 175. Saber A.A., M.H., Elgamal, T.H., El-Ghazaly, A.V., Dewoolkar, A., Akl. Simple
34 technique for single incision transumbilical laparoscopic appendectomy. *Int J Surg.*
35 2010;8(2):128-30.
36
37
38
39
40 176. Sadraei-Moosavi S.-M., N., Nikhbakhsh, A.-A., Darzi. Postoperative antibiotic therapy
41 after appendectomy in patients with non-perforated appendicitis. *Caspian J Int Med.*
42 2017;8(2):104-7.
43
44
45
46
47 177. Saha N., D.K., Saha, M.A., Rahman, M.K., Islam, M.A., Aziz. Comparison of post
48 operative morbidity between laparoscopic and open appendectomy in children.
49 *Mymensingh Med J.* 2010;19(3):348-52.
50
51
52
53
54 178. Sahn M., R., Kube, S., Schmidt, C., Ritter, M., Pross, H., Lippert. Current analysis of
55 endoloops in appendiceal stump closure. *Surg Endosc Interv Tech.* 2011;25(1):124-9.
56
57
58
59 179. Sahn M., M., Pross, R., Otto, A., Koch, I., Gastinger, H., Lippert. Clinical health service
60

- 1
2
3 research on the surgical therapy of acute appendicitis: Comparison of outcomes based on 3
4 German multicenter quality assurance studies over 21 years. *Ann Surg.* 2015;262(2):338-
5 46.
6
7
8
9
- 10 180. Salö M., E., Järbur, M., Hambræus, B., Ohlsson, P., Stenström, E., Arnbjörnsson. Two-
11 trocar appendectomy in children - description of technique and comparison with
12 conventional laparoscopic appendectomy. *BMC Surg.* 2016;16(1):52.
13
14
15
16
- 17 181. Sánchez-Santana T., J.A., del-Moral-Luque, P., Gil-Yonte, L., Bañuelos-Andrío, M.,
18 Durán-Poveda, G., Rodríguez-Caravaca. Effect of compliance with an antibiotic
19 prophylaxis protocol in surgical site infections in appendectomies. Prospective cohort
20 study. Efecto de la adecuación a protocolo de la profilaxis antibiótica en la incidencia de
21 infección quirúrgica en apendicectomías Estudio de cohortes prospectivo. 2017;85(3):208-
22 13.
23
24
25
26
27
28
- 29 182. Sauvain M.-O., K., Slankamenac, M.K., Muller, S., Wildi, U., Metzger, W., Schmid, et
30 al. Delaying surgery to perform CT scans for suspected appendicitis decreases the rate of
31 negative appendectomies without increasing the rate of perforation nor postoperative
32 complications. *Langenbeck's Arch Surg.* 2016;401(5):643-9.
33
34
35
36
37
38
- 39 183. Scarborough J.E., K.M., Bennett, T.N., Pappas. Racial disparities in outcomes after
40 appendectomy for acute appendicitis. *Am J Surg.* 2012;204(1):11-7.
41
42
43
44
- 45 184. Seifarth F.G., N., Kundu, A.D., Guerron, M.M., Garland, M.W., Gaffley, S., Worley, et
46 al. Umbilical Negative Pressure Dressing for Transumbilical Appendectomy in Children.
47 *JLS.* 2016;20(4).
48
49
50
- 51 185. Federico G. Seifarth, Kundu, Neilendu, Guerron, Alfredo D., Garland, Mary M.,
52 Gaffley, Michaela W. G., Worley, Sarah, et al. Umbilical Negative Pressure Dressing for
53 Transumbilical Appendectomy in Children. *JLS-JOURNAL OF THE SOCIETY OF*
54 *LAPAROENDOSCOPIC SURGEONS.* 2016;20(4).
55
56
57
58
59
- 60 186. Senekjian L., R., Nirula. Tailoring the operative approach for appendicitis to the patient:

- 1
2
3 A prediction model from national surgical quality improvement program data. *J Am Coll*
4 *Surg.* 2013;216(1):34-40.
5
6
7
8
9 187. Sesia S.B., M., Frech, F.-M., Häcker, J., Mayr. Laparoscopic "single-port"
10 appendectomy in children. *Laparoskopische "single port"-appendektomie im Kindesalter.*
11 2011;136(1):50-5.
12
13
14
15 188. Shaikh A.R., S., Khatoon, M., Arif. Evaluation of re-admission after open
16 appendectomy. *Rawal Med J.* 2011;36(2):100-3.
17
18
19
20
21
22 189. Shang Q., Q., Geng, X., Zhang, C., Guo. The efficacy of combined therapy with
23 metronidazole and broad-spectrum antibiotics on postoperative outcomes for pediatric
24 patients with perforated appendicitis. *Medicine.* 2017;96(47).
25
26
27
28
29 190. Shindholimath V., K., Thinakaran, T., Rao, Y., Veerappa. Laparoscopic management
30 of appendicular mass. *J Minimal Access Surg.* 2011;7(2):136-40.
31
32
33
34 191. Shimizu T., M., Ishizuka, K., Kubota. The preoperative serum C-reactive protein level
35 is a useful predictor of surgical site infections in patients undergoing appendectomy. *Surg*
36 *Today.* 2015;45(11):1404-10.
37
38
39
40
41 192. Siam B., A., Al-Kurd, N., Simanovsky, H., Awesat, Y., Cohn, B., Helou, et al.
42 Comparison of appendectomy outcomes between senior general surgeons and general
43 surgery residents. *JAMA Surg.* 2017;152(7):679-85.
44
45
46
47
48 193. 서승원, 김신곤. Acute Appendicitis in Pregnant Patients and Non-Pregnant Patients:
49
50
51 Recent Clinical Experience of the Tertiary Hospital Z1 임신부와 가임기 여성의 급성
52
53
54
55
56
57
58
59
60
충수염의 비교 고찰: 최근 3차 병원의 임상적 경험. *Annals of Surgical Treatment and*
Research S1 대한외과학회지. 2002;62(6):486-90.

- 1
2
3
4
5
6
7 194. Alfredo Silva, M, Guido Vargas, A, Amparo Moreno, H, Pablo Becerra. Utilidad del
8 retractor elástico abdominal para disminuir el riesgo de infección de herida operatoria en
9 apendicitis aguda X1 Use of an elasticwall retractor during appendectomy to reduce wound
10 infection. *Revista chilena de cirugía*. 2008;60(6):527-33.
11
12
13
14
15 195. Singh V.K., K., Nishant, B., Kharga, A.K., Kalita, P., Bhutia, J., Jain. Randomized
16 controlled trial comparing open, conventional, and single port laparoscopic appendectomy.
17 *J Clin Diagn Res*. 2017;11(10):PC05-PC10.
18
19
20
21
22
23
24 196. Siribumrungwong B., K., Srikuea, A., Thakkinstian. Comparison of superficial surgical
25 site infection between delayed primary and primary wound closures in ruptured
26 appendicitis. *Asian J Surg*. 2014;37(3):120-4.
27
28
29
30
31 197. Sivrikoz E., E., Karamanos, E., Beale, P., Teixeira, K., Inaba, D., Demetriades. The
32 effect of diabetes on outcomes following emergency appendectomy in patients without
33 comorbidities: A propensity score-matched analysis of National Surgical Quality
34 Improvement Program database. *Am J Surg*. 2015;209(1):206-11.
35
36
37
38
39 198. Soll C., P., Wyss, H., Gelpke, D.A., Raptis, S., Breitenstein. Appendiceal stump closure
40 using polymeric clips reduces intra-abdominal abscesses. *Langenbeck's Arch Surg*.
41 2016;401(5):661-6.
42
43
44
45
46 199. Sozutek A., T., Colak, M., Dirlik, K., Ocal, O., Turkmenoglu, A., Dag. A prospective
47 randomized comparison of single-port laparoscopic procedure with open and standard 3-
48 port laparoscopic procedures in the treatment of acute appendicitis. *Surg Laparoscopy*
49 *Endosc Percutaneous Tech*. 2013;23(1):74-8.
50
51
52
53
54 200. Srishewachart P., S., Narksut. Incidence of abnormal preoperative blood testing and
55 postoperative complication in appendectomy patients in Siriraj Hospital. *J Med Assoc*
56 *Thailand*. 2016;99(5):517-24.
57
58
59
60

- 1
2
3 201. Staszewicz W., M.-C., Eisenring, V., Bettschart, S., Harbarth, N., Troillet. Thirteen
4 years of surgical site infection surveillance in Swiss hospitals. *J Hosp Infect.* 2014;88(1):40-
5 7.
6
7
8
9
10 202. Suttie S.A., S., Seth, C.P., Driver, A.A., Mahomed. Outcome after intra- and extra-
11 corporeal laparoscopic appendectomy techniques. *Surg Endosc.* 2004;18(7):1123-5.
12
13
14
15 203. Svensson J.F., B., Patkova, M., Almström, S., Eaton, T., Wester. Outcome after
16 introduction of laparoscopic appendectomy in children: A cohort study. *J Pediatr Surg.*
17 2016;51(3):449-53.
18
19
20
21
22 204. Taguchi Y., S., Komatsu, E., Sakamoto, S., Norimizu, Y., Shingu, H., Hasegawa.
23 Laparoscopic versus open surgery for complicated appendicitis in adults: a randomized
24 controlled trial. *Surg Endosc Interv Tech.* 2016;30(5):1705-12.
25
26
27
28
29 205. Tanaka S., D., Kubota, S.H., Lee, K., Oba, M., Matsuyama. Effectiveness of
30 laparoscopic approach for acute appendicitis. *Osaka City Med J.* 2007;53(1):1-8.
31
32
33
34 206. Tijerina J., R., Velasco-Rodríguez, C., Vásquez, V., Melnikov, S., Rodriguez.
35 Effectiveness of a systemic antibiotic followed by topical ionized solution as surgical site
36 infection prophylaxis. *J Int Med Res.* 2010;38(4):1287-93.
37
38
39
40
41 207. The SCARLESS Study Group. Single port/incision laparoscopic surgery compared with
42 standard three-port laparoscopic surgery for appendectomy: A randomized controlled
43 trial. *Surg Endosc Interv Tech.* 2015;29(1):77-85.
44
45
46
47
48 208. Juan Pablo Toro, Barrera, Óscar Javier, Morales, Carlos Hernando. Superioridad clínica
49 de la apendicectomía laparoscópica sobre la técnica abierta: ¿adopción lenta de un nuevo
50 estándar de tratamiento? X1 Clinical superiority of laparoscopic appendectomy over the
51 open technique: sluggish adoption of a new standard of treatment? *Revista Colombiana de*
52 *Cirugía.* 2017;32(1):32-9.
53
54
55
56
57
58 209. Towfigh S., T., Clarke, W., Yacoub, A.H., Pooli, R.J., Mason, N., Katkhouda, et al.
59 Significant reduction of wound infections with daily probing of contaminated wounds : A
60

- 1
2
3 prospective randomized clinical trial. *Arch Surg.* 2011;146(4):448-52.
4
5
6
7 210. Troillet N., E., Aghayev, M.-C., Eisenring, A.F., Widmer. First Results of the Swiss
8 National Surgical Site Infection Surveillance Program: Who Seeks Shall Find. *Infect*
9 *Control Hosp Epidemiol.* 2017;38(6):697-704.
10
11
12
13 211. Tsioplis C., C., Brockschmidt, S., Sander, D., Henne-Bruns, M., Kornmann. Factors
14 influencing the course of acute appendicitis in adults and children. *Langenbeck's Arch Surg.*
15 2013;398(6):857-67.
16
17
18
19
20 212. Vahdad M.R., M., Nissen, A., Semaan, T., Klein, E., Palade, T., Boemers, et al.
21 Experiences with LESS-appendectomy in Children. *Arch Iran Med.* 2016;19(1):57-63.
22
23
24
25 213. Van Rossem C.C., M.H.F., Schreinemacher, A.A.W., Van Geloven, W.A., Bemelman,
26 G.J.D., Van Acker, B., Akkermans, et al. Antibiotic duration after laparoscopic
27 appendectomy for acute complicated appendicitis. *JAMA Surg.* 2016;151(4):323-9.
28
29
30
31
32 214. Van Rossem C.C., M.H.F., Schreinemacher, K., Treskes, R.M., Van Hogeand,
33 A.A.W., Van Geloven. Duration of antibiotic treatment after appendectomy for acute
34 complicated appendicitis. *Br J Surg.* 2014;101(6):715-9.
35
36
37
38
39 215. Wang-Chan A., F.H., Hetzer, C., Gingert, C., Gingert, E., Angst, E., Angst, et al.
40 Clinical relevance and effect of surgical wound classification in appendicitis: Retrospective
41 evaluation of wound classification discrepancies between surgeons, Swissnoso-trained
42 infection control nurse, and histology as well as surgical site infection rates by wound class.
43 *J Surg Res.* 2017;215:132-9.
44
45
46
47
48
49 216. Watanabe A., S., Kohnoe, H., Sonoda, K., Shirabe, K., Fukuzawa, S., Maekawa, et al.
50 Effect of intra-abdominal absorbable sutures on surgical site infection. *Surg Today.*
51 2012;42(1):52-9.
52
53
54
55
56 217. Willis Z.I., E.M., Duggan, B.T., Bucher, J.B., Pietsch, M., Milovancev, W., Wharton,
57 et al. Effect of a clinical practice guideline for pediatric complicated appendicitis. *JAMA*
58 *Surg.* 2016;151(5).
59
60

- 1
2
3
4
5 218. Ramírez-Wong F.M., T., Atencio-Espinoza, V.D., Rosenthal, E., Ramirez, S.L., Torres-
6 Zegarra, Z.R., Díaz Tavera, et al. Surgical Site Infections Rates in More Than 13,000
7 Surgical Procedures in Three Cities in Peru: Findings of the International Nosocomial
8 Infection Control Consortium. *Surg Infect*. 2015;16(5):572-6.
9
10
11
12
13 219. Wu J.-M., K.-H., Chen, H.-F., Lin, L.-M., Tseng, S.-H., Tseng, S.-H., Huang.
14 Laparoscopic appendectomy in pregnancy. *J Laparoendosc Adv Surg Techn Part A*.
15 2005;15(5):447-50.
16
17
18
19
20 220. Wu H.-S., H.-W., Lai, S.-J., Kuo, Y.-T., Lee, D.-R., Chen, C.-W., Chi, et al. Competitive
21 edge of laparoscopic appendectomy versus open appendectomy: A subgroup comparison
22 analysis. *J Laparoendosc Adv Surg Techn*. 2011;21(3):197-202.
23
24
25
26
27 221. Wu K., L., Yang, A., Wu, J., Wang, S., Xu, H., Zhao, et al. Single-site laparoscopic
28 appendectomy in children using conventional instruments: a prospective, randomized,
29 control trial. *Pediatr Surg Int*. 2014;31(2):167-71.
30
31
32
33
34 222. Wu T.-C., Q., Lu, Z.-Y., Huang, X.-H., Liang. Efficacy of emergency laparoscopic
35 appendectomy in treating complicated appendicitis for elderly patients. *Saudi Med J*.
36 2017;38(11):1108-12.
37
38
39
40
41 223. Yaghoubian A., C., de Virgilio, V., Chiu, S.L., Lee. "July effect" and appendicitis. *J*
42 *Surg Educ*. 2010;67(3):157-60.
43
44
45
46 224. Yagnik V., J., Rathod, A., Phatak. A retrospective study of two-port appendectomy and
47 its comparison with open appendectomy and three-port appendectomy. *Saudi J*
48 *Gastroenterol*. 2010;16(4):268-71.
49
50
51
52
53 225. Yousef Y., F., Yousef, M., Homsy, T., Dinh, K., Pandya, H., Stagg, et al.
54 Standardization of care for pediatric perforated appendicitis improves outcomes. *J Pediatr*
55 *Surg*. 2017;52(12):1916-20.
56
57
58
59
60 226. Zhang Z., Y., Wang, R., Liu, L., Zhao, H., Liu, J., Zhang, et al. Suprapubic single-

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

incision versus conventional laparoscopic appendectomy. J Surg Res. 2016;200(1):131-8.

For peer review only



PRISMA 2009 Checklist

Section/topic	#	Checklist item	Reported on page #
TITLE			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	1
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	2-3
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known.	5-6
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	6
METHODS			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	8
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	6
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	6
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	6
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	7
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	7
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	7
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	8
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).	7-8
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., I^2) for each meta-analysis.	7-8



PRISMA 2009 Checklist

Section/topic	#	Checklist item	Reported on page #
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	7-8
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	8
RESULTS			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	8
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	8
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	8
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	8-9
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	8-9
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).	8-9
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).	9
DISCUSSION			
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).	10
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	12
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	12
FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	14

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit: www.prisma-statement.org.

BMJ Open

Global incidence of surgical-site infection after appendectomy: a systematic review and meta-analysis

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2019-034266.R1
Article Type:	Original research
Date Submitted by the Author:	23-Nov-2019
Complete List of Authors:	Danwang, Celestin; Universite Libre de Bruxelles, School of Public Health; University of Yaoundé I, Department of Surgery and Specialties, Faculty of Medicine and Biomedical Sciences Bigna, Jean Joel; Centre Pasteur of Cameroon, Department of Epidemiology and Public Health Tochie, Joel Noutakdie; University of Yaoundé I, Department of Anesthesiology and Critical Care Medicine, Faculty of Medicine and Biomedical Sciences Mbonda, Aimé; Djohong District Hospital, Department of General Medicine Mbanga, Clarence; Mankon Sub-divisional Hospital Nzalie, Rolf ; Ngong District Hospital, North Region, Cameroon Guifo, Marc Leroy ; University of Yaoundé I, Department of Surgery and Specialties, Faculty of Medicine and Biomedical Sciences Essomba, Arthur ; University of Yaoundé I, Department of Surgery and Specialties, Faculty of Medicine and Biomedical Sciences
Primary Subject Heading:	Surgery
Secondary Subject Heading:	Global health, Infectious diseases, Public health
Keywords:	EPIDEMIOLOGY, Health policy < HEALTH SERVICES ADMINISTRATION & MANAGEMENT, Quality in health care < HEALTH SERVICES ADMINISTRATION & MANAGEMENT, Epidemiology < INFECTIOUS DISEASES, Public health < INFECTIOUS DISEASES

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

1 **Global incidence of surgical-site infection after appendectomy: a systematic** 2 **review and meta-analysis**

3 Celestin Danwang^{1,2}, Jean Joel Bigna^{3*}, Joel Noutakdie Tochie⁴, Aime Mbonda⁵, Clarence
4 Mvalo Mbanga⁶, Rolf Nyah Tuku Nzalie⁷, Marc Leroy Guifo¹, Arthur Essomba¹

- 5
- 6 1. Department of Surgery and Specialties, Faculty of Medicine and Biomedical Sciences,
7 University of Yaoundé I, Yaoundé, Cameroon.
- 8 2. School of Public Health, Université Libre de Bruxelles, Brussels, Belgium.
- 9 3. Department of Epidemiology and Public Health, Centre Pasteur of Cameroon, Yaoundé,
10 Cameroon.
- 11 4. Department of Anesthesiology and Critical Care Medicine, Faculty of Medicine and Biomedical
12 Sciences, University of Yaoundé I, Yaoundé, Cameroon.
- 13 5. Department of General Medicine, Djohong District Hospital, Djohong, Cameroon.
- 14 6. Department of General Medicine, Mankon Sub-Divisional Hospital, Bamenda,
15 Cameroon.
- 16 7. Department of General Medicine, Ngong District Hospital, Ngong, Cameroon.

17
18 *Corresponding author: Dr. Jean Joel Bigna, MD, MPH

19 Department of Epidemiology and Public Health, Centre Pasteur of Cameroon, Yaoundé,
20 Cameroon. E-Mail: bignarimjj@yahoo.fr

21
22 E-mail addresses:

23 CD: danram07@yahoo.fr; JJB: bignarimjj@yahoo.fr; JNT: joeltochie@gmail.com; AM:
24 aimembonda@yahoo.fr; CMM: mbangaclarence@gmail.com; RNN: nzalieolf@yahoo.com;
25 GML: marcleroyguifo@gmail.com; AE: artgessomba@gmail.com

26
27 **Word count:** 2,774.

28 **Abstract**

29 **Objectives:** Although surgical-site infection (SSI) is one of the most studied healthcare-
30 associated infections, the global burden of SSI after appendectomy remains unknown. Hence,
31 we estimated the incidence of SSI after appendectomy at global and regional levels.

32 **Design:** Systematic review and meta-analysis.

33 **Participants:** Patients with appendectomy.

34 **Data sources:** EMBASE, PubMed, and Web of Science were searched to identify observational
35 studies and clinical trials, published between January 1, 2000 and December 30, 2018 and
36 reporting on the incidence of the SSI after appendectomy; with no language restriction. A
37 random-effect models meta-analysis served to obtain the pooled incidence of SSI after 100
38 surgical procedures in patients with appendectomy.

39 **Results:** In total, 226 studies (729,434 participants from 49 countries) were included in the
40 meta-analysis. Concerning the methodological quality, 59 (26.1%) studies had a low risk, 147
41 (65.0%) a moderate risk, and 20 (8.8%) a high risk of bias. We found an overall incidence of
42 SSIs of 7.0 per 100 surgical procedures (95% prediction interval: 1.0-17.6) for appendectomy
43 varying from 0 to 37.4 per 100 surgical procedures. Subgroup analysis for identifying sources
44 of heterogeneity showed that the incidence varied from 5.8 in Europe to 12.6 per 100 surgical
45 procedures in Africa, $p < 0.0001$. The incidence of SSI after appendectomy increased when the
46 level of income decreased; from 6.2 in high-income countries to 11.1 per 100 surgical
47 procedures in low-income countries ($p = 0.015$). Open appendectomy (11.0 per 100 surgical
48 procedures) was found to have a higher incidence of SSI compared to laparoscopy (4.6 per 100
49 surgical procedures), $p = 0.0002$.

50 **Conclusion:** This study suggests a high burden of SSIs after appendectomy in some regions
51 (especially Africa) and in low-income countries. Strategies are needed to implement and
52 vulgarize WHO guidelines to decrease the burden of SSI after appendectomy in these regions.

1
2
3 53 **Registration:** PROSPERO, CRD42017075257.
4
5
6 54
7
8 55
9

10 56 **Keywords:**

11
12 57 Surgical wound infection; Global Health; Hospital infections; Cross infection; Healthcare
13
14
15 58 associated infection
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

59 **Strengths and limitations of this study**

- 60 • This meta-analysis is the first to summarize the global incidence of SSIs after
61 appendectomy.
- 62 • We investigated WHO regions, level of income, and surgical procedure as sources of
63 heterogeneity.
- 64 • We were not able to investigate all sources of heterogeneity because of missing information
65 in the original studies.
- 66 • There were few studies from low income countries and from Africa.

Introduction

Defined as an acute inflammation of the vermiform appendix,¹ evidence abounds that acute appendicitis is the most common abdominal surgical emergency,² with an incidence of almost 100 per 100,000 person-years reported in Australia, Europe and North America.^{3 4} Evidence suggests appendectomy, a surgical remove of the vermiform appendix as first-line treatment for acute appendicitis, although antibiotic therapy may be efficacious for a selected group of patients with uncomplicated acute appendicitis.⁵⁻⁷ Appendectomy is a relatively safe surgical intervention with a case fatality rate of 2.1 - 2.4 per 1000 patients as reported in studies conducted in Europe.^{8 9}

Innovations in appendectomy, especially with the advent of minimally invasive or laparoscopic surgery in 1983,¹⁰ which has replaced the traditional open appendectomy in most of high-income countries, has led to a drastic reduction in the morbidity and mortality related to appendectomy.¹¹⁻¹³ Laparoscopic appendectomy is now recognized as the gold standard surgical approach for uncomplicated acute appendicitis owing to its merits over open surgery; due to less postoperative pain, reduced postoperative ileus, shorter hospital stay, rapid postoperative recovery, and better aesthetic scars.¹⁴⁻¹⁹

However, regardless of the surgical technique (laparoscopic or open surgery), appendectomy remains a sceptical surgical intervention associated with a substantial risk of surgical-site infections (SSIs). SSIs after appendectomy are postoperative nosocomial infections affecting the incision site, deep tissues, organs at the operative site within 30 days after the surgical procedure.²⁰⁻²² SSI following appendectomy is a serious post-operative medical concern that increases the financial burden for both healthcare systems and patient, and also have a negative impact on the patients' health related quality of life.²³⁻²⁸

SSI is both the most frequently studied and the leading healthcare-associated infections reported hospital-wide in low- and middle-income countries.²⁹ A recently published prospective

1
2
3 international multicentre cohort study suggested a high burden of SSIs after any gastrointestinal
4 surgery in low-income countries compared to high-income countries.³⁰ Actually, there is no
5 global systematic review with meta-analysis reporting the burden of SSI after appendectomy or
6
7 comparing the burden between regions and between country level of income. It would be
8 interesting to have such accurately estimated data to construct efficient strategies to curb
9
10 globally the burden of SSIs after appendectomy. In an effort to fill this gap, the current
11
12 systematic review and meta-analysis aimed at summarizing contemporary data on the
13
14 occurrence of SSIs after appendectomy.
15
16
17
18
19
20
21
22
23

24 **Methods**

25 **Design**

26
27 This systematic review and meta-analysis was registered in the International Prospective
28
29 Register of Systematic Reviews (PROSPERO) under the registration number
30
31 CRD42017075257. The protocol has been published in a peer-review journal.³¹
32
33
34
35

36 **Eligibility criteria**

37
38 We considered observational studies (cross-sectional, case-control, and cohort) and clinical
39
40 trials of patients with appendectomy. Outcome of interest was incidence of SSI of enough data
41
42 (number of cases of SSI and sample size) to compute this estimate. We excluded letters,
43
44 reviews, commentaries and editorials, and studies lacking key data and/or explicit method
45
46 description as well as studies in which relevant data on SSIs after appendectomy was impossible
47
48 to extract even after contacting the corresponding author.
49
50
51
52

53 **Search strategy**

54
55 We searched EMBASE, PubMed, and Web of Science (Web of Science Core Collection,
56
57 Current Contents Connect, KCI-Korean Journal Database, SciELO Citation Index, Russian
58
59
60

1
2
3 Science Citation Index) to identify observational studies, published between January 1, 2000
4 and December 30, 2018. No language restriction was applied. The initial search strategy was
5 designed for EMBASE and was adapted for the use in others databases. The search strategy as
6 illustrated in the Supplementary Table 1 and in the study protocol,³¹ was based on the
7 combination of relevant text words and medical subject headings related to SSIs. Moreover, the
8 references of all relevant articles found were scrutinized for potential additional data sources.
9
10 When a full text was not available, it was requested via the corresponding author by email. For
11 duplicates or studies published in more than one report, the one reporting the largest sample
12 size was considered.
13
14
15
16
17
18
19
20
21
22

23 **Study selection**

24
25
26
27 Two reviewers (CD and AM) independently screened the titles and abstract of articles for
28 eligibility. Full texts of potentially eligible articles were retrieved and screened for final
29 inclusion. Disagreements between the two reviewers were solved by discussion and when a
30 consensus was not reached, a third reviewer (JNT) resolved discrepancies. Studies in other
31 languages than French, English, and Spanish were translated using Google Translate.
32
33
34
35
36
37

38 **Data extraction and management**

39
40
41 A standardized and pretested data extraction form was used by five reviewers (CD, JNT, AM,
42 RNZ, CMM) to independently extract data from individual studies. A sixth reviewer (JJB)
43 independently extracted data for accuracy. The last name of the first author, year of publication,
44 country, study design, age groups, sample size, mean or median age, proportion of males,
45 specific conditions of the study population, the surgical method (open surgery or laparoscopy),
46 and incidence of SSIs after appendectomy in the study population (or enough data to compute
47 this estimate) were extracted.
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 To assess the methodological quality of each study, two reviewers (CD and CMM) used an
4 adapted version of the tool of bias assessment for prevalence studies developed by Hoy and
5 colleagues.³²
6
7
8
9

10 **Data synthesis and analysis**

11
12
13 A meta-analysis was used to summarize data concerning incidence of SSIs, by pooling together
14 data of studies reporting the incidence of SSIs. Study-specific estimates were then pooled
15 through a Dersimonian and Laird random-effects meta-analysis model to obtain an overall
16 summary estimate of the incidence across studies, after stabilizing the variance of individual
17 studies using the Freeman-Tukey double arc-sine transformation.³³ Incidence was expressed by
18 100 surgical procedures with their 95% confidence interval and 95% prediction interval.
19 Heterogeneity was evaluated by the χ^2 test on Q statistic which is quantified by I^2 values,³⁴
20 assuming that I^2 values of 25%, 50% and 75% represent low, medium and high heterogeneity
21 respectively.³⁵ Where substantial heterogeneity ($I^2 > 50\%$) was detected, a subgroup analysis
22 was performed to detect its possible sources using the following grouping variables: type of
23 surgery (laparoscopy or open), World Health Organization regions, and country level of
24 income. A p value < 0.05 was indicative of significant difference. The meta-regression analysis
25 was performed to estimate the explained heterogeneity of each covariate included in the
26 subgroup analysis. Inter-rater agreement for study inclusion was assessed using Cohen's κ
27 coefficient.³⁶ Funnel plots analysis and Egger's test ($p < 0.10$) were performed to detect the
28 presence of publication bias.³⁷ Since we believe that the incidence estimates of interest would likely
29 be published even if substantially different from previously reported estimates, we have not reported
30 adjusted incidence estimate in the case of publication bias. Data were analysed using the '*meta*' package
31 in R, version 3.6.1.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55

56 **Patient and public involvement**

1
2
3 Patients or the public were not involved in the design, or conduct, or reporting, or dissemination
4
5 of our research.
6
7
8
9

10 **Results**

11 **Study selection and characteristics**

12
13 Overall, 619 records were initially identified. After removal of duplicates, screening of study
14
15 titles, abstracts, and full texts; 226 studies including 729,434 patients were finally retained for
16
17 meta-analysis (Supplementary Figure 1). The full list of included studies is in the Appendix.
18
19 Concerning the methodological quality, 59 (26.1%) studies had a low risk, 147 (65.0%) a
20
21 moderate risk and 20 (8.8%) a high risk of bias. Supplementary Table 2 presents characteristics
22
23 of included studies. Among the included studies, 154 were done in high-income, 36 upper-
24
25 middle, 27 lower-middle, and nine in low-income countries. Overall, most of studies were from
26
27 Europe (n = 68) and Americas (n = 67). SSIs were defined according to Center of Disease
28
29 Control and Prevention criteria in 50 studies while 25 studies used other criteria. The definition
30
31 of SSIs was not clearly given in 151 studies.
32
33
34
35
36
37
38

39 **Overall prevalence**

40
41 The overall incidence of SSI after appendectomy was 7.0 per 100 surgical procedures (95%
42
43 prediction interval: 1.0-17.6) varying from 0% to 37.4% with substantial heterogeneity and
44
45 publication bias (Supplementary Figure 2). The sensitive analysis including only studies with
46
47 low risk of bias yielded a very close incidence to crude analysis (Table 1).
48
49
50

51 **Sources of heterogeneity**

52
53 According to country level of income (Figure 1), the incidence of SSI after appendectomy
54
55 increased when the level of income decreased; from 6.2 in high income countries to 11.1 per
56
57 100 surgical procedures in low income countries ($p = 0.015$) (Table 1).
58
59
60

1
2
3 The incidence varied widely across WHO regions (Figure 2). The incidence varied from 5.8 in
4 Europe to 12.6 per 100 surgical procedures in Africa, $p < 0.0001$ (Table 1). Two regions
5 (Europe and Americas) had an incidence < 6 per 100 surgical procedures, three an incidence
6 between 6-10 per 100 procedures (South-East Asia, Eastern Mediterranean, and Western
7 Pacific), and one an incidence > 10 per 100 procedures (Africa) (Table 1). The incidence also
8 varied widely in different regions. The incidence varied from 0.2 to 32.0 in Africa, from 1.9 to
9 37.4 in Western Pacific, from 1.3 to 33.8 in Eastern Mediterranean, from 1.2 to 25.8 in South-
10 East Asia, from 0.1 to 37.4 in Americas, and from 0 to 20.0 per 100 surgical procedures in
11 Europe (Figure 2).

12
13
14
15
16
17
18
19
20
21
22
23
24 Open appendectomy with an incidence of 11.0 (95% prediction interval: 0.0-39.3) per 100
25 surgical procedures was found to have a higher incidence of SSI compared to laparoscopic
26 appendectomy with an incidence of 4.6 (95% prediction interval: 0.0-14.3) per 100 surgical
27 procedures, $p = 0.0002$ (Figure 3).

28
29
30
31
32
33 Heterogeneity of the overall incidence of SSI after appendectomy was explained by WHO
34 regions (17.1%), country level of income (11.1%), and type of surgical procedure (4.9%). In
35 the meta-regression analysis of 119 studies reporting the information of the use of antibiotics,
36 there was no association between the variation of SSI incidence and proportion of patients with
37 the use of antibiotics (coefficient: 0.0010 [95%CI: -0.0004; 0.0023]; $p = 0.170$). however, most
38 (79.5%) of these studies reported using antibiotics for all patients.

49 Discussion

50
51
52 This first systematic review and meta-analysis of data of 729,434 surgical procedures in 226
53 studies from 49 countries found an overall incidence of SSIs of 7.0 per 100 surgical procedures
54 for appendectomy varying from 0 to 37.4 per 100 surgical procedures with substantial
55 heterogeneity according to WHO regions, country level of income, and type of surgical
56
57
58
59
60

1
2
3 procedure. The incidence increased with decreasing country level of income and was higher
4
5 when using open surgery compared to laparoscopy. The incidence significantly varied by WHO
6
7 regions with Africa having the highest burden followed by Western Pacific, Eastern-
8
9 Mediterranean, and South-East Asia. We found no association between SSI incidence and
10
11 proportion of using antibiotics.
12
13

14 Health care-associated infections are acquired by patients when receiving care and are the most
15
16 frequent adverse event affecting patient safety worldwide. This includes SSIs after
17
18 appendectomy.³⁸ As reported in a previous systematic review and meta-analysis, SSIs were the
19
20 leading infection in hospitals in developed countries.²⁹ The high incidence we found in this
21
22 study suggests that SSIs after appendectomy remains a global public concern. WHO reported
23
24 that of every 100 hospitalized patients at any given time, seven in developed and 15 in
25
26 developing countries will acquire at least one health care-associated infection.³⁸ SSIs are mainly
27
28 caused by micro-organisms resistant to commonly-used antimicrobials, which can be
29
30 multidrug-resistant. Indeed, more than 50% of SSIs can be antibiotic-resistant.³⁹ The leading
31
32 micro-organisms identified in SSIs are *Staphylococcus aureus*, coagulase-negative
33
34 staphylococci, and *Escherichia coli* as reported by National Healthcare Safety Network.³⁹ It is
35
36 important to worry since *Staphylococcus aureus* and *Escherichia coli* are the micro-organisms
37
38 with highest proportion of antibiotic resistance, respectively resistant to oxacillin/methicillin in
39
40 43% of cases and to fluoroquinolones in 25% of cases.³⁹ A recent international prospective
41
42 cohort study shown that 21.6% of patients with SSI after any gastrointestinal surgery had an
43
44 infection that was resistant to the prophylactic antibiotic used.³⁰ There are many factors that can
45
46 favour SSI including patient-related and procedural-related variable.⁴⁰ These factors can be
47
48 classified in two categories; non-modifiable like age and sex and modifiable including
49
50 nutritional status, tobacco use, correct use of antibiotics, obesity, diabetes, prolonged surgery
51
52 duration, pre-surgery hospital stay of at least two days, lower volume of hospital and surgeons,
53
54
55
56
57
58
59
60

1
2
3 and the intraoperative techniques.³⁸ Strategies to curb the burden of SSIs should therefore focus
4
5 on addressing these identified factors. However, we were not able to find an association
6
7 between SSI with the use antibiotics, may be due to the low variability in the proportion of
8
9 antibiotics in the original studies.

10
11
12 In our present study looking at specifically SSI after appendectomy, we also found that SSI was
13
14 higher in low income countries. Interestingly, there was a trend with increasing incidence when
15
16 the country income decreased. The WHO Africa region essentially constituted with sub-
17
18 Saharan Africa was the region with highest incidence in this study. The WHO estimates that
19
20 the endemic burden of health care-associated infections is two to three time significantly higher
21
22 in low- and middle-income countries than in high-income nations.³⁸ The highest burden found
23
24 in Africa may be associated with the fact most of countries in this continent are low income
25
26 countries compared to other regions. Indeed, factors associated with increased risk of SSI after
27
28 appendectomy may be higher in low-income settings. The burden of diabetes, obesity, and
29
30 undernutrition are increasing in low-income countries.^{41 42} There is also inadequate use of
31
32 antimicrobial in low- and middle-income countries and micro-organisms are more resistant to
33
34 prophylactic antibiotics used to prevent SSI in low-income countries compared to high-income
35
36 countries.^{30 43 44} Lower level income is also associated with lower volume of surgeon and
37
38 hospital, factors recognised as associated increased risk of SSIs.³⁸ The higher incidence found
39
40 in low income countries may also be explained by the fact open surgery is the most used surgical
41
42 procedure in this setting. Indeed, we found as in other studies that open surgery is associated
43
44 with higher incidence of SSIs compared to laparoscopy.^{45 46} Laparoscopy is generally indicated
45
46 for uncomplicated appendicitis where the dissemination of micro-organism is lower compared
47
48 open surgery indicated for perforated appendicitis with peritonitis for example. Moreover, only
49
50 few low-income countries have the necessary infrastructure to carry out laparoscopy procedures
51
52 compared to high-income countries.⁴⁷⁻⁴⁹
53
54
55
56
57
58
59
60

1
2
3 Our findings have important implications for healthcare providers and health policy makers.
4
5 SSIs are among the most preventable healthcare-associated infections.^{50 51} They still represent
6
7 a significant burden in terms of patient morbidity and mortality and additional costs for
8
9 healthcare systems.³⁸ The prevention of SSI has received considerable attention from surgeons,
10
11 infection control professionals, health policy makers, the media and the public since there is a
12
13 perception among the public that SSIs may reflect a poor quality of care.⁵² However, special
14
15 attention is needed for low-income countries and Africa. Strategy to curb the burden of SSIs
16
17 after appendectomy as for other surgery procedures should be focused on strategies that can
18
19 help to address factors associated with increased risk of SSIs. Therefore, strategies should be a
20
21 package including how to address the factors cited above. The 26 WHO recommendations to
22
23 avoid SSIs should be vulgarized and implemented,³⁸ especially in low-income countries.
24
25 Strengthening the healthcare systems of low-income countries and of countries in WHO Afro
26
27 region is also a paramount by education of healthcare providers and skilling them on the use of
28
29 very less invasive surgical procedures.
30
31
32
33

34
35 This study should however be interpreted in the context of some drawbacks. Firstly, the same
36
37 definition of SSIs was not used by all the included studies. This may lead to an overestimation
38
39 or underestimation of the SSIs incidence by individual studies (depending on the definition
40
41 used). Secondly, few studies reported on the participants' characteristics and details on the
42
43 surgical procedure since this can modify the risk for developing SSIs. We were not therefore
44
45 able to measure the impact on our outcome of interest. Thirdly, only a quarter of studies had
46
47 low risk of bias, however our analysis including only studies with low risk of bias yielded an
48
49 estimate close to the crude incidence. Fourth, the various geographic regions and countries were
50
51 variably represented, with some countries with only one study or even no study, which could
52
53 affect the generalizability of our findings.
54
55
56
57
58
59
60

1
2
3 Despite these limitations, this is the first systematic review and meta-analysis providing a global
4 estimate of the burden of SSIs after appendectomy. A protocol had been published before, and
5
6 estimate of the burden of SSIs after appendectomy. A protocol had been published before, and
7
8 we used rigorous methodological and statistical procedures to obtain and pool data.
9
10 Furthermore, subgroup analyses were conducted to investigate the various factors likely
11
12 affecting our estimate.
13
14

17 **Conclusion**

19 This systematic review and meta-analysis compiled data from more than 700,000 people with
20
21 appendicitis in 49 countries and pointed a high incidence of SSIs after appendectomy, at 7 per
22
23 100 surgical procedures. This estimate seemed higher in some WHO regions (especially Africa)
24
25 and in low-income countries. These data suggest that less invasive procedure is associated with
26
27 low incidence of SSIs after appendectomy. Strategies are needed to implement already known
28
29 guidelines to decrease the burden of SSI after appendectomy. However, in low-income
30
31 countries which have weak health systems, cost-effectiveness studies are needed to inform
32
33 policies regarding the best strategies for decreasing the burden of SSI after appendectomy.
34
35
36
37
38
39

40 **Contributors**

41
42
43 CD and JJB conceived the idea of the study and developed the protocol. JJB, CD, and JNT did
44
45 the literature search. CD, AM, and JNT selected the studies, CD, JNT, RNZ, AM, CMM, JJB
46
47 extracted the relevant information. CD, JJB, and CMM synthesized the data. CD, JNT, CMM,
48
49 and JJB wrote the first draft of the paper. CD, JJB, JNT, AB, RNZ, CMM, GML, and AE
50
51 critically revised successive drafts of the paper and approved the final version. GML and AE
52
53 supervised the overall work, CD and JJB are the guarantors of the review.
54
55
56

57 **Acknowledgements**

58
59 None.
60

Funding

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests

We declare no competing interests.

Patient consent

Not applicable.

Data sharing statement

All data generated for this study are in the manuscript and its supporting files.

Figures Legend

Figure 1. Global incidence of SSI (surgical site infection) after appendectomy by level of country income

Figure 2. Global incidence of SSI (surgical site infection) after appendectomy by WHO regions

Figure 3. Global incidence of SSI (surgical site infection) after appendectomy by type of surgical procedures

References

1. Giesen LJX, van den Boom AL, van Rossem CC, et al. Retrospective Multicenter Study on Risk Factors for Surgical Site Infections after Appendectomy for Acute Appendicitis. *Dig Surg* 2017;34(2):103-07. doi: 10.1159/000447647 [published Online First: 09/16]
2. Navarro Fernández JA, Tárraga López PJ, Rodríguez Montes JA, et al. Validity of tests performed to diagnose acute abdominal pain in patients admitted at an emergency department. *Rev Esp Enferm Dig* 2009;101(9):610-18. doi: 10.4321/s1130-01082009000900003
3. Ohmann C, Franke C, Kraemer M, et al. Status report on epidemiology of acute appendicitis. *Chirurg* 2002;73(8):769-76. doi: 10.1007/s00104-002-0512-7

- 1
2
3 4. Körner H, Söreide JA, Pedersen EJ, et al. Stability in incidence of acute appendicitis. A
4 population-based longitudinal study. *Dig Surg* 2001;18(1):61-66. doi:
5 10.1159/000050099
6
- 7
8 5. Martínez Carrilero J. Safety and efficacy of antibiotics compared with appendicectomy for
9 treatment of uncomplicated acute appendicitis: meta-analysis of randomised controlled
10 trials. *Rev Clin Esp* 2012;212(9):460-60. doi: 10.1016/j.rce.2012.05.005
11
- 12
13 6. Varadhan KK, Neal KR, Lobo DN. Safety and efficacy of antibiotics compared with
14 appendicectomy for treatment of uncomplicated acute appendicitis: meta-analysis of
15 randomised controlled trials. *BMJ (Clinical research ed)* 2012;344:e2156-e56. doi:
16 10.1136/bmj.e2156
17
- 18
19 7. Masoomi H, Nguyen NT, Dolich MO, et al. Laparoscopic appendectomy trends and
20 outcomes in the United States: data from the Nationwide Inpatient Sample (NIS), 2004-
21 2011. *Am Surg* 2014;80(10):1074-77.
22
- 23
24 8. Kotaluoto S, Ukkonen M, Pauniahio S-L, et al. Mortality Related to Appendectomy; a
25 Population Based Analysis over Two Decades in Finland. *World J Surg* 2017;41(1):64-
26 69. doi: 10.1007/s00268-016-3688-6
27
- 28
29 9. Blomqvist PG, Andersson RE, Granath F, et al. Mortality after appendectomy in Sweden,
30 1987-1996. *Ann Surg* 2001;233(4):455-60. doi: 10.1097/00000658-200104000-00001
31
- 32
33 10. Semm K. Endoscopic appendectomy. *Endoscopy* 1983;15(2):59-64. doi: 10.1055/s-2007-
34 1021466
35
- 36
37 11. Xiao Y, Shi G, Zhang J, et al. Surgical site infection after laparoscopic and open
38 appendectomy: a multicenter large consecutive cohort study. *Surg Endosc*
39 2015;29(6):1384-93. doi: 10.1007/s00464-014-3809-y [published Online First: 10/11]
40
- 41
42 12. Varela JE, Wilson SE, Nguyen NT. Laparoscopic surgery significantly reduces surgical-
43 site infections compared with open surgery. *Surg Endosc* 2010;24(2):270-76. doi:
44 10.1007/s00464-009-0569-1 [published Online First: 06/17]
45
- 46
47 13. Bregendahl S, Nørgaard M, Laurberg S, et al. Risk of complications and 30-day mortality
48 after laparoscopic and open appendectomy in a Danish region, 1998-2007;
49 a population-based study of 18,426 patients. *Pol Przegl Chir* 2013;85(7):395-400. doi:
50 10.2478/pjs-2013-0060
51
- 52
53 14. Sauerland S, Jaschinski T, Neugebauer EA. Laparoscopic versus open surgery for suspected
54 appendicitis. *The Cochrane database of systematic reviews* 2010(10):CD001546-CD46.
55 doi: 10.1002/14651858.CD001546.pub3
56
- 57
58 15. Aziz O, Athanasiou T, Tekkis PP, et al. Laparoscopic versus open appendectomy in
59 children: a meta-analysis. *Ann Surg* 2006;243(1):17-27. doi:
60 10.1097/01.sla.0000193602.74417.14
16. Dai L, Shuai J. Laparoscopic versus open appendectomy in adults and children: A meta-
analysis of randomized controlled trials. *United European Gastroenterol J*
2017;5(4):542-53. doi: 10.1177/2050640616661931 [published Online First: 08/16]

17. Markides G, Subar D, Riyad K. Laparoscopic versus open appendectomy in adults with complicated appendicitis: systematic review and meta-analysis. *World J Surg* 2010;34(9):2026-40. doi: 10.1007/s00268-010-0669-z
18. Wei B, Qi C-L, Chen T-F, et al. Laparoscopic versus open appendectomy for acute appendicitis: a metaanalysis. *Surg Endosc* 2011;25(4):1199-208. doi: 10.1007/s00464-010-1344-z [published Online First: 09/17]
19. Li X, Zhang J, Sang L, et al. Laparoscopic versus conventional appendectomy--a meta-analysis of randomized controlled trials. *BMC Gastroenterol* 2010;10:129-29. doi: 10.1186/1471-230X-10-129
20. Horan TC, Gaynes RP, Martone WJ, et al. CDC definitions of nosocomial surgical site infections, 1992: a modification of CDC definitions of surgical wound infections. *Am J Infect Control* 1992;20(5):271-74. doi: 10.1016/s0196-6553(05)80201-9
21. Culver DH, Horan TC, Gaynes RP, et al. Surgical wound infection rates by wound class, operative procedure, and patient risk index. National Nosocomial Infections Surveillance System. *Am J Med* 1991;91(3B):152S-57S. doi: 10.1016/0002-9343(91)90361-z
22. Horan TC, Andrus M, Dudeck MA. CDC/NHSN surveillance definition of health care-associated infection and criteria for specific types of infections in the acute care setting. *Am J Infect Control* 2008;36(5):309-32. doi: 10.1016/j.ajic.2008.03.002
23. Badia JM, Casey AL, Petrosillo N, et al. Impact of surgical site infection on healthcare costs and patient outcomes: a systematic review in six European countries. *J Hosp Infect* 2017;96(1):1-15. doi: 10.1016/j.jhin.2017.03.004 [published Online First: 03/08]
24. Thompson KM, Oldenburg WA, Deschamps C, et al. Chasing zero: the drive to eliminate surgical site infections. *Ann Surg* 2011;254(3):430-37. doi: 10.1097/SLA.0b013e31822cc0ad
25. Hawn MT, Vick CC, Richman J, et al. Surgical site infection prevention: time to move beyond the surgical care improvement program. *Ann Surg* 2011;254(3):494-501. doi: 10.1097/SLA.0b013e31822c6929
26. Mehta JA, Sable SA, Nagral S. Updated recommendations for control of surgical site infections. *Ann Surg* 2015;261(3):e65-e65. doi: 10.1097/SLA.0b013e318289c5fd
27. Andersson RE. Short-term complications and long-term morbidity of laparoscopic and open appendectomy in a national cohort. *Br J Surg* 2014;101(9):1135-42. doi: 10.1002/bjs.9552 [published Online First: 06/30]
28. Pinkney TD, Calvert M, Bartlett DC, et al. Impact of wound edge protection devices on surgical site infection after laparotomy: multicentre randomised controlled trial (ROSSINI Trial). *BMJ (Clinical research ed)* 2013;347:f4305-f05. doi: 10.1136/bmj.f4305
29. Allegranzi B, Bagheri Nejad S, Combescure C, et al. Burden of endemic health-care-associated infection in developing countries: systematic review and meta-analysis.

- 1
2
3 *Lancet (London, England)* 2011;377(9761):228-41. doi: 10.1016/S0140-
4 6736(10)61458-4 [published Online First: 12/09]
5
6
7 30. GlobalSurg Collaborative. Surgical site infection after gastrointestinal surgery in high-
8 income, middle-income, and low-income countries: a prospective, international,
9 multicentre cohort study. *Lancet Infect Dis* 2018;18(5):516-25. doi: 10.1016/S1473-
10 3099(18)30101-4 [published Online First: 02/13]
11
12 31. Danwang C, Mazou TN, Tochie JN, et al. Global prevalence and incidence of surgical site
13 infections after appendectomy: a systematic review and meta-analysis protocol. *BMJ*
14 *open* 2018;8(8):e020101-e01. doi: 10.1136/bmjopen-2017-020101
15
16 32. Hoy D, Brooks P, Woolf A, et al. Assessing risk of bias in prevalence studies: modification
17 of an existing tool and evidence of interrater agreement. *J Clin Epidemiol*
18 2012;65(9):934-9. doi: 10.1016/j.jclinepi.2011.11.014 [published Online First:
19 2012/06/30]
20
21 33. Barendregt JJ, Doi SA, Lee YY, et al. Meta-analysis of prevalence. *Journal of epidemiology*
22 *and community health* 2013;67(11):974-78. doi: 10.1136/jech-2013-203104 [published
23 Online First: 08/20]
24
25 34. Cochran WG. The Combination of Estimates from Different Experiments. *Biometrics*
26 1954;10(1):101-29.
27
28 35. Higgins JPT, Thompson SG, Deeks JJ, et al. Measuring inconsistency in meta-analyses.
29 *BMJ (Clinical research ed)* 2003;327(7414):557-60. doi: 10.1136/bmj.327.7414.557
30
31 36. Viera AJ, Garrett JM. Understanding interobserver agreement: the kappa statistic. *Fam Med*
32 2005;37(5):360-63.
33
34 37. Egger M, Davey Smith G, Schneider M, et al. Bias in meta-analysis detected by a simple,
35 graphical test. *BMJ (Clinical research ed)* 1997;315(7109):629-34. doi:
36 10.1136/bmj.315.7109.629 [published Online First: 1997/10/06]
37
38 38. WHO. Global guidelines on the prevention of surgical site infection: WHO; 2016 [Available
39 from: [https://apps.who.int/iris/bitstream/handle/10665/250680/9789241549882-
40 eng.pdf?sequence=8](https://apps.who.int/iris/bitstream/handle/10665/250680/9789241549882-eng.pdf?sequence=8) accessed Nov 23 2019].
41
42 39. Sievert DM, Ricks P, Edwards JR, et al. Antimicrobial-resistant pathogens associated with
43 healthcare-associated infections: summary of data reported to the National Healthcare
44 Safety Network at the Centers for Disease Control and Prevention, 2009-2010. *Infect*
45 *Control Hosp Epidemiol* 2013;34(1):1-14. doi: 10.1086/668770 [published Online First:
46 11/27]
47
48 40. Buggy D. Can anaesthetic management influence surgical-wound healing? *Lancet (London,*
49 *England)* 2000;356(9227):355-57. doi: 10.1016/S0140-6736(00)02523-X
50
51 41. Seidell JC, Halberstadt J. The global burden of obesity and the challenges of prevention.
52 *Ann Nutr Metab* 2015;66 Suppl 2:7-12. doi: 10.1159/000375143 [published Online
53 First: 06/02]
54
55
56
57
58
59
60


- 1
- 2
- 3 42. Checkley W, Ghannem H, Irazola V, et al. Management of NCD in low- and middle-income
- 4 countries. *Glob Heart* 2014;9(4):431-43. doi: 10.1016/j.ghheart.2014.11.003
- 5
- 6 43. Versporten A, Zarb P, Caniaux I, et al. Antimicrobial consumption and resistance in adult
- 7 hospital inpatients in 53 countries: results of an internet-based global point prevalence
- 8 survey. *Lancet Glob Health* 2018;6(6):e619-e29. doi: 10.1016/S2214-109X(18)30186-
- 9 4 [published Online First: 04/23]
- 10
- 11 44. Klein EY, Van Boeckel TP, Martinez EM, et al. Global increase and geographic
- 12 convergence in antibiotic consumption between 2000 and 2015. *Proceedings of the*
- 13 *National Academy of Sciences of the United States of America* 2018;115(15):E3463-
- 14 E70. doi: 10.1073/pnas.1717295115 [published Online First: 03/26]
- 15
- 16 45. Foster D, Kethman W, Cai LZ, et al. Surgical Site Infections after Appendectomy
- 17 Performed in Low and Middle Human Development-Index Countries: A Systematic
- 18 Review. *Surg Infect (Larchmt)* 2018;19(3):237-44. doi: 10.1089/sur.2017.188
- 19 [published Online First: 10/23]
- 20
- 21 46. Marchi M, Pan A, Gagliotti C, et al. The Italian national surgical site infection surveillance
- 22 programme and its positive impact, 2009 to 2011. *Euro Surveill* 2014;19(21):20815.
- 23 doi: 10.2807/1560-7917.es2014.19.21.20815
- 24
- 25 47. Udwardia TE. Diagnostic laparoscopy. *Surg Endosc* 2004;18(1):6-10. doi: 10.1007/s00464-
- 26 002-8872-0 [published Online First: 09/10]
- 27
- 28 48. Adisa AO, Lawal OO, Arowolo OA, et al. Local adaptations aid establishment of
- 29 laparoscopic surgery in a semiurban Nigerian hospital. *Surg Endosc* 2013;27(2):390-
- 30 93. doi: 10.1007/s00464-012-2463-5 [published Online First: 07/18]
- 31
- 32 49. Alfa-Wali M, Osaghae S. Practice, training and safety of laparoscopic surgery in low and
- 33 middle-income countries. *World J Gastrointest Surg* 2017;9(1):13-18. doi:
- 34 10.4240/wjgs.v9.i1.13
- 35
- 36 50. Haley RW, Culver DH, White JW, et al. The efficacy of infection surveillance and control
- 37 programs in preventing nosocomial infections in US hospitals. *American journal of*
- 38 *epidemiology* 1985;121(2):182-205. doi: 10.1093/oxfordjournals.aje.a113990
- 39
- 40 51. Harbarth S, Sax H, Gastmeier P. The preventable proportion of nosocomial infections: an
- 41 overview of published reports. *J Hosp Infect* 2003;54(4):258-321. doi: 10.1016/s0195-
- 42 6701(03)00150-6
- 43
- 44 52. Birgand G, Lepelletier D, Baron G, et al. Agreement among healthcare professionals in ten
- 45 European countries in diagnosing case-vignettes of surgical-site infections. *PloS one*
- 46 2013;8(7):e68618-e18. doi: 10.1371/journal.pone.0068618
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

Table 1. Summary statistics of meta-analysis incidence of surgery site infections after appendectomy

	Incidence per 100 surgical procedures (95%CI)	95% Prediction interval	N studies	N participants	H (95%CI)	I ² (95%CI)	P heterogeneity	P Egger test	P difference
Global	7.0 (6.4-7.7)	1.0-17.7	226	729,434	8.9 (8.7-9.1)	98.7 (98.7-98.8)	< 0.0001	< 0.0001	-
- Low risk of bias	6.9 (6.0-7.9)	1.6-15.2	59	204,450	6.7 (6.3-7.1)	97.7 (97.4-98.0)	< 0.0001	< 0.0001	-
By Level of income									
- Low	11.1 (5.5-18.2)	0.0-42.2	9	1,496	3.8 (3.0-4.8)	93.1 (89.0-95.6)	< 0.0001	0.735	0.015
- Lower-middle	9.2 (6.3-12.6)	0.0-31.6	27	10,379	5.1 (4.6-5.7)	96.2 (95.3-96.9)	< 0.0001	0.960	
- Upper-middle	8.5 (6.5-10.8)	0.3-25.3	36	26,557	5.4 (2.9-5.9)	96.6 (95.9-97.1)	< 0.0001	0.392	
- High	6.2 (5.6-6.9)	0.9-15.3	154	691,002	9.5 (9.2-9.8)	98.9 (98.8-99.0)	< 0.0001	< 0.0001	
By WHO regions									
- Africa	12.6 (3.3-26.4)	0.0-72.5	8	3,001	9.1 (7.9-10.5)	98.8 (98.4-99.1)	< 0.0001	0.628	< 0.0001
- Western Pacific	9.6 (8.1-11.2)	2.3-20.8	43	30,822	3.8 (3.5-4.2)	93.2 (91.7-94.4)	< 0.0001	0.150	
- Eastern Mediterranean	8.2 (6.4-10.2)	1.7-18.6	23	7,779	2.6 (2.2-3.1)	85.3 (79.1-89.6)	< 0.0001	0.515	
- South-East Asia	7.6 (4.7-11.1)	0.0-24.6	16	5,782	3.8 (3.2-4.5)	93.0 (90.1-95.0)	< 0.0001	0.0001	
- Americas	5.9 (5.2-6.6)	1.9-11.7	67	401,931	7.5 (7.1-7.9)	98.2 (98.0-98.4)	< 0.0001	0.0004	
- Europe	5.8 (4.6-7.0)	0.0-19.1	68	276,793	10.4 (10.0-10.8)	99.1 (99.0-99.1)	< 0.0001	< 0.0001	
By type of surgical procedure									
- Laparoscopy with open surgery	4.6 (2.5-7.2)	0.0-15.6	10	4,892	3.2 (2.6-4.2)	90.7 (85.0-94.2)	< 0.0001	0.942	0.0002
- Laparoscopy	4.6 (3.4-5.9)	0.0-14.3	40	33,873	4.4 (4.0-4.8)	94.7 (93.6-95.7)	< 0.0001	0.0002	
- Open surgery	11.0 (7.9-14.4)	0.0-39.3	44	13,120	5.7 (5.2-6.1)	96.9 (96.4-97.3)	< 0.0001	0.077	


WHO: World Health Organization; CI: confidence interval; H: H statistics






Author, Year **Sample** **Incidence** **[95% C.I.]** **Per 100 surgical procedure**



Procedure = Laparoscopy



Procedure = Laparoscopy with Open Surgery



Procedure = Open Surgery



MOOSE Checklist for Meta-analyses of Observational Studies

Item No	Recommendation	Reported on Page No
Reporting of background should include		
1	Problem definition	5
2	Hypothesis statement	6
3	Description of study outcome(s)	5
4	Type of exposure or intervention used	NA
5	Type of study designs used	5-6
6	Study population	5-6
Reporting of search strategy should include		
7	Qualifications of searchers (eg, librarians and investigators)	13
8	Search strategy, including time period included in the synthesis and key words	6; Suppl. Table 1
9	Effort to include all available studies, including contact with authors	6
10	Databases and registries searched	6
11	Search software used, name and version, including special features used (eg, explosion)	6
12	Use of hand searching (eg, reference lists of obtained articles)	6
13	List of citations located and those excluded, including justification	8, Suppl. Fig 1, Suppl. References
14	Method of addressing articles published in languages other than English	7
15	Method of handling abstracts and unpublished studies	7
16	Description of any contact with authors	6
Reporting of methods should include		
17	Description of relevance or appropriateness of studies assembled for assessing the hypothesis to be tested	6
18	Rationale for the selection and coding of data (eg, sound clinical principles or convenience)	6-7
19	Documentation of how data were classified and coded (eg, multiple raters, blinding and interrater reliability)	6-7
20	Assessment of confounding (eg, comparability of cases and controls in studies where appropriate)	8
21	Assessment of study quality, including blinding of quality assessors, stratification or regression on possible predictors of study results	8
22	Assessment of heterogeneity	7
23	Description of statistical methods (eg, complete description of fixed or random effects models, justification of whether the chosen models account for predictors of study results, dose-response models, or cumulative meta-analysis) in sufficient detail to be replicated	7-8
24	Provision of appropriate tables and graphics	Table 1; Fig 1-3
Reporting of results should include		
25	Graphic summarizing individual study estimates and overall estimate	Fig 1-3
26	Table giving descriptive information for each study included	Suppl. Table 1
27	Results of sensitivity testing (eg, subgroup analysis)	9; Table 1

28	Indication of statistical uncertainty of findings	9; Table 1; Fig 1-3
Item No	Recommendation	Reported on Page No
Reporting of discussion should include		
29	Quantitative assessment of bias (eg, publication bias)	12-13
30	Justification for exclusion (eg, exclusion of non-English language citations)	12-13
31	Assessment of quality of included studies	12-13
Reporting of conclusions should include		
32	Consideration of alternative explanations for observed results	13
33	Generalization of the conclusions (ie, appropriate for the data presented and within the domain of the literature review)	13
34	Guidelines for future research	13
35	Disclosure of funding source	14


From: Stroup DF, Berlin JA, Morton SC, et al, for the Meta-analysis Of Observational Studies in Epidemiology (MOOSE) Group. Meta-analysis of Observational Studies in Epidemiology. A Proposal for Reporting. *JAMA*. 2000;283(15):2008-2012. doi: 10.1001/jama.283.15.2008.

Global incidence of surgical-site infection after appendectomy: a systematic review and meta-analysis

APPENDIX

Celestin **Danwang**, Jean Joel **Bigna**, Joel Noutakdie **Tochie**,
 Aime **Mbonda**, Clarence Mvalo **Mbanga**, Rolf Nyah Tuku **Nzalie**,
 Marc Leroy **Guifo**, Arthur **Essomba**


Supplementary Figure 1. Study flow	2
Supplementary Figure 2. Funnel plot for publication bias	3
Supplementary Table 1. Search strategy in EMBASE	4
Supplementary Table 2 : Characteristics of included studies	5
Supplementary Table 3. Individual characteristics of included studies	7
Reference list of included studies.....	23



Supplementary Figure 1. Study flow

review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Supplementary Figure 2. Funnel plot for publication bias

Supplementary Table 1. Search strategy in EMBASE

	Search terms
#1	'appendectomy'/exp OR appendectomy OR 'appendicectomy'/exp OR appendicectomy OR appendices OR 'appendix epiploica' OR 'omental appendix' OR 'appendicitis'/exp OR appendicitis
#2	'surgical site infection'/exp OR 'surgical site infection' OR 'surgical wound infection'/exp OR 'surgical wound infection' OR 'surgical wound infections'/exp OR 'surgical wound infections' OR 'surgical site infections' OR 'operative site infections' OR 'postoperative wound infections'/exp OR 'postoperative wound infections' OR 'postoperative wound infection'/exp OR 'postoperative wound infection'
#3	[2000-2018]/py
#4	#1 AND #2 AND #3

Supplementary Table 2 : Characteristics of included studies

Characteristics	N = 226
Year of publication, range	2000-2018
%Male, range	0-100 (n = 195)
Mean/median age, range	7-74 (n = 186)
%HIV	0-13.1 (n = 2)
%Diabetes	0-95.7 (n = 34)
%Obesity	0-7.4 (n = 18)
Design, n	
- Cross sectional	120
- Cohort study	99
- Case control	7
WHO regions, n	
- Africa	8
- Americas	67
- Eastern Mediterranean	23
- Europe	68
- Multiregional	1
- South-East Asia	16
- Western Pacific	43
Level of income, n	
- Low	9
- Lower-middle	27
- Upper-middle	36
- High	154
Timing of data collection	
- Retrospective	123
- Prospective	101
- Unclear	2
Sampling	
- Consecutive	131
- Systematic	37
- Random	32
- Exhaustive	11
- Unclear	15
Number of sites	
- Multisite	51
- One site	170
- Unclear	5
Pattern of appendicitis, range	
- %Catarrhal	0-100 (n = 84)
- %Perforated	0-100 (n = 110)
- %Suppurated	0-100 (n = 70)
- %Gangrenous	0-46.7 (n = 89)
%With administered antibiotics	24.1-100 (n = 109)
%With administered analgesics	64.5-100 (n = 20)
%With diet > 6 or 8 hours	50-100 (n = 3)
Type of surgery	
- %Open surgery	0-100 (n = 134)

1		
2		
3	- %Laparoscopy	0-100 (n = 187)
4	Mean/median time to complete the	0.1-2.2 (n = 106)
5	intervention (in hours), range	
6	Type of anesthesia, n	
7		
8	- General	118
9	- Spinal and general	2
10	- Unclear	106
11	SSI definition, n	
12	- CDC-NNIS criteria	50
13	- Other criteria	25
14	- Not reported/Unclear	151
15		
16		
17		
18		
19		
20		
21		
22		
23		
24		
25		
26		
27		
28		
29		
30		
31		
32		
33		
34		
35		
36		
37		
38		
39		
40		
41		
42		
43		
44		
45		
46		
47		
48		
49		
50		
51		
52		
53		
54		
55		
56		
57		
58		
59		
60		

For peer review only

Supplementary Table 3. Individual characteristics of included studies

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Adejumo	2015	Moderate	Cohort	Nigeria	Retrospective	Consecutive	One site	2007-2014	Adults	39	26	NR	Catarrhal, Perforated, Suppurated, Gangrenous	48.9	NR	NR	NR	100	NR	NR	NR	NR	141
Adhikar	2008	Moderate	Cohort	Nepal	Prospective	Consecutive	One site	2005-2006	Children, Adolescents, Adults, Elderly	NR	NR	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy	0.5	General	NR	50
Aguillo	2005	Moderate	Cohort	Spain	Prospective	Consecutive	Unclear	NR	Children, Adolescents, Adults, Elderly	63.1	NR	NR	Unclear	NR	NR	NR	NR	100	NR	NR	NR	NR	792
Ahmad	2014	Moderate	Clinical trial	Pakistan	Prospective	Consecutive	One site	2012	Adults	35.4	27.4	NR	Perforated	0	100	0	0	100	Open Surgery	NR	General	NR	158
Ahmed	2015	Moderate	Cross sectional	Pakistan	Retrospective	Consecutive	One site	2009-2010	Children, Adolescents	51.89	10.1	NR	Unclear	NR	NR	NR	NR	100	Open Surgery	NR	NR	NR	79
Akkoyun	2012	Moderate	Case control	Turkey	Retrospective	Consecutive	One site	1998-2011	Children	64.5	8.9	NR	Perforated	0	100	0	0	100	Open Surgery	0.6	General	NR	234
Ali	2015	Moderate	Cross sectional	Pakistan	Prospective	Consecutive	One site	2014	Adults	46.3	27.4	NR	Unclear	NR	NR	NR	NR	100	Open Surgery	NR	General	NR	121
Ali	2012	Moderate	Cohort	Nigeria	Prospective	Consecutive	One site	2002-2009	Children, Adolescents, Adults, Elderly	33.9	32	NR	Catarrhal, Perforated, Suppurated, Gangrenous	NR	23.47	NR	NR		Open Surgery	NR	NR	NR	1257
Almström	2016	Moderate	Cohort	Sweden	Retrospective	Systematic	One site	2006-2013	Children, Adolescents	59.5	NR	NR	Perforated, Non Perforated 76%	NR	24	NR	NR		Laparoscopy or Open Surgery	0.8	NR	NR	2756
Al-Saadi	2006	Moderate	Cohort	Yemen	Retrospective	Consecutive	One site	2003-2005	Children, Adolescents, Adults	75	NR	NR	Catarrhal, Perforated, Suppurated, Gangrenous	NR	60	NR	13	100	Laparoscopy	NR	NR	NR	160
Al-Temimi	2017	Low	Cohort	USA	Prospective	Systematic	One site	2016	Children, Adolescents, Adults, Elderly	40.2	30	NR	Catarrhal, Perforated, Suppurated, Gangrenous, Normal	73	17.4	3.3	3.3		Laparoscopy	0.6	NR	NR	92
Álvarez-Moreno	2014	Low	Cohort	Colombia	Prospective	Systematic	Multisite	2008-2010	Children, Adolescents, Adults, Elderly	NR	NR	NR	Unclear	NR	NR	NR	NR		NR	NR	NR	According to CDC-NNIS diagnostic criteria	947
Anderson	2014	Moderate	Cohort	Sweden	Retrospective	Exhaustive	Multisite	1992-2008	Adults	54	NR	NR	Perforated, Not perforated	NR	19.4	0	0		Laparoscopy or Open Surgery	NR	NR	NR	169896
Andert	2017	Moderate	Cohort	Germany	Retrospective	Consecutive	One site	2003-2014	Adults	48.6	30.5	NR	Catarrhal, Perforated, Suppurated, Gangrenous	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	NR	NR	Local signs of inflammation	2136
Aranda-Narvaez	2014	Moderate	Cohort	Spain	Not reported/Unclear	Not clear	One site	2007-2010	Adults	57	29	NR	Catarrhal, Perforated, Suppurated, Gangrenous	65.8	NR	NR	NR	62.00	Laparoscopy or Open Surgery	0.92	NR	According to CDC-NNIS diagnostic Criteria	868
Aranda-Narváez	2010	Low	Cohort	Spain	Retrospective	Random	One site	1997-2009	Children, Adolescents, Adults, Elderly	63.3	35	NR	Suppurated, Gangrenous	0	0	53.3	46.7	100	Laparoscopy or Open Surgery	NR	General	According to CDC-NNIS diagnostic Criteria	30

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Arthur	2017	Low	Cross sectional	Australia	Prospective	Systematic	Multisite	2016	Children, Adolescents, Adults, Elderly	49.5	31.4	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	1	NR	NR	1189
Asefa	2014	Moderate	Cross sectional	Ethiopia	Retrospective	Consecutive	One site	2006-2010	Children	62.1	10	NR	Catarrhal, Perforated, Suppurated, Gangrenous	32.3	59.6	6.2	1.9		NR	NR	NR	NR	322
Asefa	2002	High	Cross sectional	Ethiopia	Retrospective	Consecutive	One site	1997-1999	Adults	79.5	25.6	NR	Catarrhal, Perforated	45.4	44	0	0		NR	NR	NR	NR	200
Bae	2016	Low	Cross sectional	Korea	Prospective	Systematic	One site	2014-2016	Adults	52	62	NR	Perforated, Suppurated, Gangrenous	NR	4	72	24	100	Laparoscopy	1.2	General	According to CDC-NNIS diagnostic criteria	25
Bae	2016	Moderate	Cross sectional	USA	Retrospective	Systematic	One site	2010-2013	Children, Adolescents, Adults	NR	32	NR	Unclear	NR	NR	NR	NR	36.4	Laparoscopy	NR	NR	According to CDC-NNIS diagnostic criteria	143
Baek	2011	Moderate	Cross sectional	Korea	Retrospective	Exhaustive	One site	2007-2009	Elderly	45.5	68.2	NR	Catarrhal, Perforated, Suppurated, Gangrenous	22.1	29.9	32.5	15.6	100	Laparoscopy or Open Surgery	1.05	General	NR	77
Bali	2015	Moderate	Cohort	Turkey	Prospective	Consecutive	One site	2009-2013	Adults	35.7	32.33	NR	Unclear	NR	NR	NR	NR		Laparoscopy	1	NR	NR	126
Banghu	2013	Low	Cohort	UK, Spain, Japan, Hong Kong, Australia, New Zealand	Prospective	Consecutive	Multisite	2012	Children, Adolescents, Adults, Elderly	51.1	NR	NR	Unclear	NR	NR	NR	NR	96.9	Laparoscopy or Open Surgery	NR	NR	According to CDC-NNIS diagnostic criteria	3326
Bansal	2012	Low	Cohort	Switzerland	Prospective	Consecutive	One site	NR	Children	62	9.8	NR	Catarrhal, Perforated	74.3	25.7	NR	NR	49.2	Laparoscopy or Open Surgery	1.0	NR	According to CDC-NNIS diagnostic Criteria	187
Batajoo	2012	Moderate	Cross sectional	Nepal	Retrospective	Consecutive	One site	2009-2012	Children, Adolescents, Adults, Elderly	45.6	29.6	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	0.8	NR	NR	226
Bharathi	2011	Moderate	Cohort	Nepal	Prospective	Consecutive	One site	2008-2009	Children, Adolescents, Adults, Elderly	50	22.9	NR	Catarrhal, Perforated, Suppurated, Gangrenous	80	NR	NR	NR	100	Laparoscopy with Open Surgery	0.5	NR	NR	26
Bıçakcı	2011	High	Cross sectional	Turkey	Retrospective	Systematic	Unclear	2006-2009	Children, Adolescents	64.5	10	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	NR	NR	279
Bickel	2011	Moderate	Clinical trial	Israel	Prospective	Random	One site	2006-2009	Adults	73	28	NR	Catarrhal, Gangrenous, Phlegmonous 58.6 , Normal 4.3%	17	NR	NR	20.5	100	Open Surgery	0.5	General	NR	210
Blackwood	2017	Moderate	Cross sectional	USA	Retrospective	Random	One site	2010-2015	Children	55.6	10.4	29.6	Unclear	NR	NR	NR	NR		Open Surgery	2	General	According to CDC-NNIS diagnostic Criteria	115
Blakely	2011	Low	Clinical trial	USA	Prospective	Random	One site	2006-2009	Children, Adolescents	55.7	10.2	NR	Perforated	0	100	0	0	100	Laparoscopy or Open Surgery	1.9	NR	NR	131

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Bonadjo	2017	Moderate	Cross sectional	USA	Retrospective	Consecutive	One site	2008-2015	Children, Adolescents	56.2	8.4	NR	Perforated	0	100	0	0	100	Laparoscopy	NR	General	NR	203
Boomer	2016	Low	Cross sectional	USA	Retrospective	Systematic	Multisite	2010-2012	Children, Adolescents	60.3	11.0	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	NR	According to CDC-NNIS diagnostic criteria	1338
Boomer	2014	Low	Cohort	USA	Retrospective	Consecutive	One site	2010-2012	Children, Adolescents	61.1	10.9	NR	Catarrhal, Perforated, Gangrenous	66.2	NR	NR	NR	97.8	Laparoscopy or Open Surgery	NR	General	Wound infection or abdominal/pelvic abscess	1388
Bozkurt	2014	Moderate	Case control	Turkey	Retrospective	Consecutive	One site	2008-2012	Children, Adolescents, Adults, Elderly	54	30.4	NR	Catarrhal	100	0	0	0		Laparoscopy	0.8	General	NR	1849
Brandt	2008	Moderate	Cross sectional	Germany	Retrospective	Systematic	Multisite	2000-2004	Children, Adolescents, Adults, Elderly	NR	NR	NR	NR	NR	NR	NR	NR		NR	NR	NR	According to CDC-NNIS diagnostic criteria	10969
Brummer	2009	Moderate	Cohort	Germany	Retrospective	Consecutive	Multisite	2004-2007	Children, Adolescents, Adults, Elderly	NR	NR	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	NR	According to CDC-NNIS diagnostic Criteria	14209
Cairo	2017	Moderate	Cohort	USA	Retrospective	Consecutive	Multisite	2012-2015	Children, Adolescents	61.3	11.0	29.9	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	General	NR	20981
Cameron	2017	Low	Cohort	USA	Retrospective	Systematic	Multisite	2012-2015	Children, Adolescents	60.4	11	11.7	Unclear	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	NR	NR	According to CDC-NNIS diagnostic criteria	1389
Cao	2015	Moderate	Cohort	China	Retrospective	Consecutive	Multisite	2011-2013	Adults	54.2	37.3	12.4	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	0.8	General	NR	12586
Cervantes-sánchez	2000	Low	Clinical trial	Mexico	Prospective	Random	One site	1994-1995	Children, Adults	53.4	28	NR	Unclear	NR	NR	NR	NR		Open Surgery	NR	General	Pus or a positive bacteriologic culture from a wound discharge	350
Chamisa	2009	High	Cross sectional	South Africa	Retrospective	Exhaustive	One site	2002-2004	Children, Adolescents, Adults, Elderly	78.4	NR	NR	Catarrhal, Perforated, Gangrenous, Normal	53	30.5	NR	10.2		Laparoscopy or Open Surgery	NR	NR	NR	324
Chaudhary	2005	Moderate	Clinical trial	Pakistan	Prospective	Random	One site	1999-2003	Children, Adolescents, Adults, Elderly	45.4	NR	NR	Catarrhal	NR	0	0	0		Open Surgery	NR	NR	NR	677
Chen	2015	Moderate	Cohort	Taiwan	Retrospective	Consecutive	One site	2010-2012	Adults	43.6	42.5	NR	Catarrhal, Perforated	87.3	12.7	0	0		Laparoscopy or Open Surgery	NR	NR	NR	236
Chen	2012	High	Cross sectional	Taiwan	Prospective	Consecutive	One site	2010	Adults	60	38	NR	Unclear	NR	NR	NR	NR	73	Laparoscopy or Open Surgery	NR	General	NR	117
Chen	2010	High	Cross sectional	China	Prospective	Systematic	One site	2008-2009	Adults	NR	NR	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	0.8	Spinal and General	NR	69
Chiang	2006	Moderate	Cross sectional	Taiwan	Retrospective	Exhaustive	One site	2002-2004	Adults	59.7	35	NR	Catarrhal, Perforated	68	17	0	0	100	Open Surgery	1.1	General	NR	390

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Chiang	2012	Moderate	Cohort	Taiwan	Prospective	Consecutive	One site	2008-2009	Adults	58.6	37.8	10	Unclear	NR	NR	NR	NR	100	Open Surgery	NR	General	Presence of gross or purulent discharge at the incision site with or without a positive bacterial culture	70
Cho	2014	Low	Cross sectional	Korea	Prospective	Consecutive	One site	2011-2012	Adults	53	38.7	18.8	Unclear	NR	NR	NR	NR		Laparoscopy	NR	General	According to CDC-NNIS diagnostic Criteria	101
Choudhary	2014	Moderate	Cross sectional	India	Prospective	Random	One site	2010-2013	Adults	67	NR	NR	Appendicular mass	0	0	0	0		NR	NR	NR	NR	92
Clyde	2008	High	Cross sectional	USA	Retrospective	Systematic	One site	2002-2007	Children, Adolescents, Adults, Elderly	52	35	NR	Catarrhal, Perforated, Unclear	77	14	0	0		Laparoscopy or Open Surgery	NR	NR	NR	1198
Coakley	2011	Low	Cohort	USA	Retrospective	Exhaustive	One site	2005-2010	Adults	47.3	28.7	NR	Catarrhal, Perforated, Suppurated, Gangrenous	38.3	1.2	47.1	9.8	100	Laparoscopy or Open Surgery	1.0	NR	According to CDC-NNIS diagnostic Criteria	728
Crandall	2009	High	Cross sectional	USA	Retrospective	Not clear	One site	2004-2005	Adults	54	32.5	NR	NR	NR	74	NR	NR		Laparoscopy or Open Surgery	0.9	General	NR	176
Dede	2008	Moderate	Cohort	Hungary	Prospective	Consecutive	One site	2005-2007	Children, Adolescents, Adults, Elderly	NR	NR	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	NR	NR	273
Dhiman	2013	High	Cross sectional	USA	Retrospective	Not clear	Multisite	2003-2009	Adults	58	30.1	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	General	NR	1250
Dimitriou	2013	Moderate	Cohort	Germany	Retrospective	Consecutive	One site	2007-2010	Children, Adolescents, Adults, Elderly	53.5	34.9	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	1	NR	NR	404
Durkin	2015	Moderate	Cohort	USA	Retrospective	Consecutive	Multisite	2007-2012	Adults	NR	NR	NR	Unclear	NR	NR	NR	NR		Laparoscopy	NR	NR	NR	14763
Ein	2013	Moderate	Cross sectional	Canada	Retrospective	Consecutive	One site	1969-2003	Children	70	7	NR	Perforated	0	100	0	0	78.8	Open Surgery	NR	General	1.Wound infection=pus draining from between the stitches or staples 2.Intra-abdominal abscess=presence of fever, abdominal pain and or gastrointestinal dysfunction and confirmed by radiologic evidence of intra-abdominal fluid collection	496

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Fukuda	2016	Moderate	Cohort	Japan	Retrospective	Consecutive	Multisite	2007-2011	Children, Adolescents, Adults	54.4	64.5	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	1.3	General	NR	2074
Gandaglia	2014	Low	Cohort	USA	Prospective	Consecutive	Multisite	2005-2011	Adolescents, Adults, Elderly	NR	NR	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	NR	General	According to CDC-NNIS diagnostic Criteria	97780
Garcell	2016	Low	Cohort	Cuba	Prospective	Consecutive	One site	2013-2015	Children, Adolescents, Adults, Elderly	95.3	30.7	2.1	Unclear	NR	NR	NR	NR		Open Surgery	NR	NR	According to CDC-NNIS diagnostic Criteria	603
Ghnnam	2011	Moderate	Cross sectional	Saudi Arabia	Retrospective	Not clear	One site	2007-2010	Adults	63.4	49.0	NR	Perforated, Unclear	NR	38.1	NR	NR		NR	NR	NR	NR	63
Giesen	2017	Moderate	Cohort	Netherlands	Retrospective	Consecutive	Multisite	2014-2015	Children, Adults	54.3	31	NR	Catarrhal, Perforated, Suppurated, Gangrenous	48.2	17.3	23.2	11.3	100	Laparoscopy or Open Surgery	0.52	NR	According to CDC-NNIS diagnostic criteria	637
Giiti	2010	Moderate	Cross sectional	Tanzania	Prospective	Systematic	One site	2008-2009	Children, Adolescents, Adults, Elderly	44.7	27	NR	Catarrhal, Perforated, Suppurated, Mass	87.4	7.0	1.5	0		NR	NR	NR	NR	199
Golub	2016	Moderate	Cohort	Russia	Retrospective	Consecutive	Multisite	2012	Adolescents, Adults	NR	34.8	NR	Unclear	NR	NR	NR	NR	100	Open Surgery	NR	General	NR	332
Gross	2016	Moderate	Cross sectional	USA	Retrospective	Consecutive	Multisite	2012-2013	Children, Adolescents	60.1	NR	17.8	Perforated	0	100	0	0		Laparoscopy or Open Surgery	NT	General	NR	2585
Gurien	2016	Moderate	Cohort	USA	Retrospective	Consecutive	One site	2009-2012	Children, Adolescents	62	10.5	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	NR	General	Wound infections or intra abdominal abscesses	484
Hamzaoglu	2004	Low	Cross sectional	Turkey	Prospective	Consecutive	One site	1999-2001	Adults	57	46.7	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	General	NR	100
Harmon	2016	Low	Cohort	USA	Retrospective	Systematic	One site	2007-2012	Children, Adolescents, Adults, Elderly	47.4	39.7	NR	Non perforated	0	0	0	0		Laparoscopy or Open Surgery	NR	NR	According to CDC-NNIS diagnostic criteria	411
Helling	2017	Low	Cross sectional	USA	Retrospective	Systematic	One site	2009-2014	Adults	64.3	34.4	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	NR	NR	According to CDC-NNIS diagnostic criteria	611
Helmer	2002	Low	Cross sectional	USA	Retrospective	Systematic	One site	1998-1999	Children, Adolescents, Adults, Elderly	NR	NR	NR	Perforated, Non perforated	NR	19.4	NR	NR	100	NR	NR	NR	A surgical wound infection was defined as purulent drainage from the wound, cellulitis requiring antibiotics, or the opening of a closed wound. An intra-abdominal abscess was defined as an intraabdominal fluid collection that contained purulent material.	438
Hesami	2014	Low	Clinical trial	Iran	Prospective	Random	Unclear	2010-2011	Children, Adolescents, Adults	58.9	27	NR	Unclear	NR	NR	NR	NR	100	Open Surgery	NR	NR	1..wound unfection= Purulent discharge, redness, inflammation, and the need to reooen	90

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
																						the wound 2...intra-abdominal abscess=abdominal pain, fullness, fever and confirmed by ecography	
Hissain	2012	Moderate	Clinical trial	Saudi Arabia	Prospective	Consecutive	One site	2010-2011	Adults	NR	32.2	NR	Catarrhal	100	0	0	0	100	Open Surgery	NR	NR	1..SSI=Pus discharge from wound needing its opening and drainage 2..Intra-abdominal collection=fluid collection inside the peritoneal cavity confirmed by ultrasound or CT scan that required drainage	377
Horvath	2016	Moderate	Cross sectional	Germany	Retrospective	Consecutive	One site	2005-2013	Adults	47	28.6	NR	Perforated, phlegmonous	NR	52	NR	NR		Laparoscopy or Open Surgery	1.0	General	According to CDC-NNIS diagnostic Criteria	1516
Hughes	2013	Moderate	Cross sectional	United Kingdom	Retrospective	Systematic	One site	2009-2010	Adults	55.6	30	NR	Unclear, simple and complicated	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	NR	General	NR	266
Hung	2016	Moderate	Cross sectional	Vietnam	Prospective	Systematic	Multisite	2008-2010	Adults	45	41.6	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	1	General	NR	752
Inigo	2006	Low	Cohort	Spain	Prospective	Consecutive	One site	1998-2002	Adults	NR	NR	NR	Unclear	NR	NR	NR	NR		NR	0.7	NR	According to CDC-NNIS diagnostic Criteria	721
Iqbal	2015	Low	Clinical trial	Pakistan	Prospective	Random	One site	2011	Adolescents, Adults, Elderly	66.3	26	NR	Catarrhal	100	0	0	0	100	Open Surgery	NR	General	According to Southampton criteria. Southampton grade 2 and above was considered as surgical site infection.	166
Javadi	2017	Moderate	Clinical trial	Iran	Prospective	Random	One site	2016	Children, Adolescents, Adults	65	19.3	NR	Catarrhal, Suppurated, Gangrenous	NR	0	NR	NR		Open Surgery	0.5	General	NR	69
Jenkins	2016	Low	Cohort	USA	Prospective	Systematic	Multisite	2006-2011	Children, Adolescents, Adults, Elderly	51.3	40.1	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	NR	According to CDC-NNIS diagnostic criteria	12410
Kang	2012	Moderate	Case control	Korea	Retrospective	Random	One site	2010-2012	Adults	54.4	31.7	NR	Catarrhal, Perforated, Suppurated, Gangrenous	NR	5.1	NR	NR	100	Laparoscopy	1.1	General	NR	217
Kapischke	2013	Low	Case control	Germany	Retrospective	Consecutive	One site	1999-2001	Children, Adolescents	47.8	11.5	NR	Catarrhal, Perforated	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	0.6	General	According to CDC-NNIS diagnostic criteria	159
Karam	2016	Moderate	Cross sectional	USA	Retrospective	Not clear	One site	2010-2015	Children	62	12	NR	Perforated, Gangrenous	NR	20.6	NR	6.2		Laparoscopy	NR	General	NR	625
Karam	2016	Moderate	Cross sectional	USA	Retrospective	Consecutive	One site	2010-2015	Children, Adolescents	63	NR	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy	1	General	NR	101
Kasatpibal	2005	Low	Cross sectional	Thailand	Prospective	Systematic	Multisite	2003-2004	Children, Adolescents, Adults, Elderly	26.6	37.2	NR	Unclear	NR	NR	NR	NR	24.1	NR	0.8	NR	According to CDC-NNIS diagnostic criteria	1487
Kasatpibal	2006	Moderate	Cohort	Thailand	Prospective	Not clear	Multisite	2003-2004	Children, Adolescents, Adults, Elderly	46.9	26	NR	Catarrhal	100	NR	NR	NR	92.2	Open Surgery	0.97	NR	According to CDC-NNIS diagnostic criteria	2139

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Kato	2008	Low	Cohort	Japan	Prospective	Systematic	One site	2004-2006	Children	NR	9.4	NR	Perforated, Non perforated 75%	NR	25	NR	NR	100	Open Surgery	NR	NR	NR	64
Kell	2003	Moderate	Cohort	Ireland	Prospective	Consecutive	Unclear	NR	Children, Adolescents, Adults, Elderly	75.2	20.7	NR	Unclear	NR	NR	NR	NR	100	Open Surgery	NR	General	Defined based on clinical and microbiological criteria	149
Khan	2007	Low	Cohort	United Kingdom	Prospective	Consecutive	One site	2006	Children, Adolescents, Adults, Elderly	47.0	24	NR	Catarrhal, Perforated	63.4	20.1	0	0	100	Laparoscopy or Open Surgery	0.9	General	According to CDC-NNIS diagnostic Criteria	134
Khan	2012	Moderate	Clinical trial	Pakistan	Prospective	Random	Multisite	2006-2009	Adults	69	33.3	NR	Unclear	NR	NR	NR	NR	100	Open Surgery	0.6	General	NR	100
Khan	2014	Moderate	Clinical trial	Pakistan	Prospective	Random	Multisite	2013-2014	Children, Adolescents, Adults, Elderly	56.7	24	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	NR	NR	Observation of pain, redness, tenderness and purulent discharge	270
Khiria	2011	Moderate	Cross sectional	India	Retrospective	Consecutive	One site	1999-2009	Adults	66	33.4	NR	Perforated, Gangrenous	NR	14.3	NR	5.2	100	Laparoscopy or Open Surgery	1.2	General	Any evidence of infection(erythema, purulent discharge, induration...) and requiring suture removal, antibiotic treatment, or evidence of dehiscence	497
Kilic	2016	Moderate	Cross sectional	Turkey	Retrospective	Consecutive	One site	2004-2010	Children	62.1	9.5	NR	Perforated	0	100	0	0	100	Open Surgery	NR	NR	According to CDC-NNIS diagnostic Criteria	110
Kim	2015	Low	Cross sectional	Korea	Retrospective	Systematic	One site	2008-2013	Children, Adolescents, Adults, Elderly	47.8	32.6	NR	Perforated, Suppurated, Gangrenous, Normal	6	13.8	64.5	7.1	100	Laparoscopy	0.7	NR	According to CDC-NNIS diagnostic criteria	2587
Kim	2011	Moderate	Cross sectional	USA	Prospective	Consecutive	One site	2005-2008	Elderly	48.1	73.4	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	NR	General	According to CDC-NNIS diagnostic Criteria	3335
Kim	2016	Moderate	Cohort	Korea	Retrospective	Consecutive	One site	2005-2012	Adults	59	NR	NR	Perforated, Gangrenous	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	1.9	General	NR	207
Kiriakopoulos	2006	Moderate	Cross sectional	Greece	Retrospective	Consecutive	One site	2000-2004	Adults	73.8	42.3	NR	Perforated, Suppurated, Generalized peritonitis	0	61.9	9.5	0	100	Laparoscopy with Open Surgery	1.1	General	NRR	42
Kirshtein	2009	Moderate	Cross sectional	Israel	Retrospective	Consecutive	One site	2000-2007	Adults	31.9	70.1	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	0.7	General	NR	1435
Kiudelis	2013	Moderate	Cross sectional	Lithuania	Prospective	Consecutive	One site	2004-2009	Adults	46.3	32.3	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy	1.1	General	NR	152
Kleif	2017	Moderate	Cross sectional	Denmark	Retrospective	Not clear	Multisite	2012-2014	Adults	53	47	NR	Suppurated, Gangrenous	NR	NR	NR	NR	98	Laparoscopy or Open Surgery	NR	General	NR	1151
Koizumi	2014	Moderate	Cross sectional	Japan	Prospective	Consecutive	One site	2010	Adults	57.9	39.8	NR	Catarrhal, Perforated, Gangrenous, phelmong	6.4	6.4	NR	25.4	100	Laparoscopy or Open Surgery	0.9	General	NR	185
Kubota	2014	Low	Clinical trial	Japan	Prospective	Random	One site	2008-2012	Children	63.6	NR	NR	Unclear	NR	NR	NR	NR	100	Open Surgery	NR	General	According to CDC-NNIS diagnostic Criteria	10

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Kumamoto	2014	Low	Cohort	Japan	Prospective	Consecutive	One site	1997-2011	Adults	0	28	NR	Catarrhal, Gangrenous, Phlegmonous	21.8	NR	NR	33.4	100	Open Surgery	0.7	General	NR	124
Kumar	2016	Moderate	Cohort	Nepal	Prospective	Consecutive	One site	2015-2016	Adolescents, Adults	49	33.9	NR	Catarrhal, Perforated, Suppurated, Gangrenous, Normal	88.6	4.7	1.9	2.4		Laparoscopy or Open Surgery	0.7	General	According to CDC-NNIS diagnostic Criteria	212
Kumar	2008	Moderate	Cohort	Pakistan	Prospective	Consecutive	One site	1997-2000	Children, Adolescents, Adults, Elderly	NR	NR	NR	Unclear	NR	NR	NR	NR	50	Laparoscopy or Open Surgery	0.7	NR	NR	100
Lacher	2012	Moderate	Cohort	USA	Prospective	Consecutive	One site	2009-2011	Children, Adolescents	64.1	10.9	22.4	Catarrhal, Perforated	71.8	19	NR	NR	100	Laparoscopy	0.7	General	NR	415
Le	2009	Moderate	Cross sectional	USA	Retrospective	Systematic	One site	1997-2007	Children, Adolescents, Adults, Elderly	52.1	31.8	NR	Catarrhal, Perforated, Gangrenous, Normal Appendix	92.9	2	0	2.2	86	Laparoscopy or Open Surgery	0.9	NR	According to CDC-NNIS diagnostic criteria	507
Lee	2009	Low	Clinical trial	USA	Prospective	Random	One site	2006-2008	Children, Adolescents, Adults, Elderly	64.2	34.2	NR	Catarrhal, Perforated, Suppurated, Gangrenous	46.8	26.6	16.5	10.1	100	Open Surgery	NR	NR	Any significant subcutaneous SSI necessitating wound opening or treatment with antibiotics. This also included any subject who was prescribed a separate course of antibiotics after discharge from the hospital.	109
Lee	2010	Moderate	Cross sectional	Taiwan	Prospective	Consecutive	One site	2006-2008	Children	58	11.1	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	2	General	NR	264
Lee	2010	Moderate	Cross sectional	Korea	Retrospective	Consecutive	One site	2008-2009	Adults	49.3	26.7	NR	Perforated, Suppurated	0	26.7	73.3	0		Laparoscopy	1.0	General	NR	75
Lee	2010	Moderate	Cohort	USA	Retrospective	Consecutive	Multisite	1998-2007	Children, Adolescents	61.5	11	NR	Perforated, Non perforated	NR	25.7	NR	NR		Laparoscopy or Open Surgery	NR	General	NR	2462
Lee	2011	Moderate	Cohort	USA	Retrospective	Systematic	Multisite	1998-2007	Children, Adolescents	61	11.6	NR	Perforated, Non perforated 70.8%	NR	29.2	NR	NR		Laparoscopy or Open Surgery	NR	NR	NR	7650
Lemieux	2008	Moderate	Cohort	Canada	Retrospective	Consecutive	One site	1997-2007	Adults	0	28.8	NR	Perforated	NR	NR	NR	NR		Laparoscopy with Open Surgery	0.8	NR	NR	45
Levy	2013	Moderate	Cohort	USA	Retrospective	Consecutive	One site	2010-2011	Children	NR	NR	NR	Catarrhal, Perforated, Suppurated, Gangrenous	56.4	32.7	4.2	6.7		Laparoscopy or Open Surgery	NR	NR	NSQIP criteria	312

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Li	2005	Moderate	Cohort	China	Prospective	Consecutive	One site	2002-2004	Children, Adolescents	71.3	7.9	NR	Catarrhal, Suppurated, Gangrenous	11.0	0	69.4	19.7	100	Laparoscopy or Open Surgery	0.65	General	NR	160
Li	2017	Moderate	Cohort	China	Retrospective	Consecutive	One site	2005-2016	Children	58.8	5.2	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	1	NR	Erythema, swelling and pus at the site of operation	398
Lim	2011	Low	Cohort	Korea	Retrospective	Consecutive	One site	2009-2011	Adults	47.8	50.8	NR	Perforated, Gangrenous	0	61.6	NR	18.3	100	Laparoscopy or Open Surgery	1.3	General	Any evidence of infection (e.g., erythema, purulent discharge, induration, etc) requiring suture removal, antibiotics or dehiscence.	60
Lima	2007	Moderate	Cross sectional	Spain	Retrospective	Consecutive	One site	2001-2006	Children, Adolescents, Adults, Elderly	NR	NR	NR	Catarrhal, Perforated, Suppurated, Gangrenous	53	2	26.3	9.7		NR	NR	NR	NR	300
Lin	2006	Moderate	Cross sectional	Taiwan	Retrospective	Consecutive	One site	2001-2003	Adults	57.6	37.5	NR	Perforated	NR	100	NR	NR	100	Laparoscopy or Open Surgery	1.4	Not described	NR	229
Litz	2016	Moderate	Cohort	USA	Retrospective	Consecutive	One site	2012-2015	Children, Adolescents	NR	11.4	17.7	Catarrhal, Perforated, Suppurated, Gangrenous, Interval, Normal	54	11.4	15.0	11.9		Laparoscopy	0.5	General	NR	413
Liu	2017	High	Cross sectional	China	Retrospective	Consecutive	Unclear	2015-2016	Children	53.6	6.6	NR	Catarrhal, Suppurated, Gangrenous	34.8	0	38.4	26.8		Laparoscopy or Open Surgery	1.0	General	NR	112
Mahmood	2016	Moderate	Clinical trial	Pakistan	Prospective	Random	One site	2012	Children, Adolescents, Adults	55.5	22.3	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	NR	General	Based on Wound Asepsis Score	200
Martinez	2012	Moderate	Cross sectional	Spain	Retrospective	Random	One site	2011	Adults	60	35.8	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	General	NR	75
Mbah	2006	Moderate	Cohort	Nigeria	Prospective	Consecutive	One site	2005	Children, Adolescents, Adults, Elderly	70	25	NR	Unclear	NR	NR	NR	NR		Open Surgery	NR	General	NR	136
Memon	2017	Moderate	Clinical trial	Pakistan	Prospective	Random	One site	2014-2016	Adults	53.3	26	NR	Catarrhal	100	0	0	0	100	Laparoscopy or Open Surgery	NR	NR	NR	227
Menezes	2008	Moderate	Cross sectional	Ireland	Retrospective	Consecutive	One site	2000-2006	Children, Adolescents	62.7	10.5	NR	Perforated, Gangrenous	0	81.4	0	17.8		Laparoscopy or Open Surgery	NR	NR	NR	118
Merenda	2013	Moderate	Cross sectional	Poland	Retrospective	Consecutive	One site	2006-2012	Children, Adolescents, Adults, Elderly	NR	NR	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	NR	NR	317
Michailidou	2015	Low	Cross sectional	USA	Retrospective	Systematic	One site	2007-2013	Children, Adolescents	56.1	9.6	NR	Perforated, Negative appendectomy	NR	26.5	NR	NR		Laparoscopy or Open Surgery	1.3	NR	According to CDC-NNIS diagnostic criteria	264
Michailidou	2015	Moderate	Cross sectional	USA	Retrospective	Consecutive	Multisite	2012	Children, Adolescents	60.1	11.2	22.5	Catarrhal, Perforated, Suppurated, Gangrenous	NR	NR	NR	NR	100	Laparoscopy	0	General	NR	2812

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Mickovic	2015	Moderate	Cross sectional	Serbia	Retrospective	Not clear	One site	2010	Children	46.4	11.7	NR	Catarrhal, Perforated, Gangrenous	45.9	2.2	NR	19.5	100	Laparoscopy or Open Surgery	0.7	General	NR	218
Ming	2009	Moderate	Cross sectional	China	Retrospective	Consecutive	One site	2003-2005	Adults	57.2	48.8	NR	Perforated, Gangrenous, Appendicular abscess	NR	72.3	NR	38.2	100	Laparoscopy or Open Surgery	NR	General	NR	173
Mingmalairak	2009	Low	Clinical trial	Thailand	Prospective	Random	One site	2006-2007	Adults	61	29.5	0	Catarrhal, Perforated, Suppurated, Gangrenous	24	16	52	8.0	100	Open Surgery	43	General	NR	100
Miyano	2010	Low	Cohort	Japan	Prospective	Consecutive	One site	2004-2008	Children, Adolescents	56.5	7.7	NR	Peritonitis complicating appendicitis	0	100	0	0	100	Laparoscopy or Open Surgery	1.9	General	NR	23
Moazzez	2013	Low	Cohort	USA	Retrospective	Not clear	One site	2005-2009	Elderly	49.3	74	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	General	NR	2060
Monge Jodra	2003	Moderate	Cohort	Spain	Prospective	Consecutive	Multisite	1997-2000	Children, Adolescents, Adults, Elderly	NR	NR	NR	Unclear	NR	NR	NR	NR		NR	NR	NR	According to CDC-NNIS diagnostic Criteria	5780
Mueck	2017	Moderate	Cohort	USA	Prospective	Consecutive	One site	2012-2015	Children, Adolescents	62.5	11.0	NR	Catarrhal, Suppurated	NR	NR	NR	NR	95	Laparoscopy with Open Surgery	NR	General	NR	697
Muensterer	2011	Low	Cohort	USA	Prospective	Consecutive	One site	2009-2010	Children, Adolescents	NR	11.2	NR	Catarrhal, Perforated	78.1	10.4	0	0	100	Laparoscopy	0.6	General	Infected umbilicus requiring antibiotics, or incision and drainage	183
Muensterer	2009	Moderate	Cross sectional	USA	Prospective	Consecutive	One site	2009	Children	61.3	11	NR	Perforated	NR	21.4	NR	NR		Laparoscopy	0.73	General	NR	75
Mustafa	2016	Low	Clinical trial	Pakistan	Prospective	Random	One site	2015-2016	Adults	52.9	26.6	NR	Perforated	0	100	0	0	100	Open Surgery	NR	NR	Redness around the wound, serosanguinous discharge, fever > 100°F	68
Nadler	2003	High	Cross sectional	USA	Retrospective	Systematic	One site	1998-2001	Children	62.2	9.35	NR	Perforated	NR	100	NR	NR	100	Open Surgery	NR	General	NR	94
Nataraja	2010	Moderate	Cohort	United Kingdom	Retrospective	Consecutive	One site	2008-2010	Children, Adolescents	59.1	11	NR	Catarrhal, Perforated, Suppurated	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	General	NR	191
Nataraja	2012	Moderate	Case control	United Kingdom	Retrospective	Consecutive	Multisite	2003-2010	Children, Adolescents	58.2	11.3	NR	Catarrhal, Perforated, Suppurated, Gangrenous	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	General	Post op intra abdominal abscess	1205
Obayashi	2015	Moderate	Cross sectional	Japan	Retrospective	Consecutive	One site	2006-2014	Children, Adolescents	60	11	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	NR	NR	485
Obinwa	2015	Moderate	Cohort	Ireland	Retrospective	Consecutive	One site	1995-2008	Children	54.5	9.6	NR	Catarrhal, Perforated, Suppurated, Gangrenous	62.7	NR	NR	4.2	100	NR	NR	NR	NR	1037
Ohene	2006	Moderate	Cross sectional	Ghana	Prospective	Consecutive	One site	1998-2004	Adults	63.9	32.4	NR	Unclear	NR	NR	NR	NR	100	Open Surgery	NR	General	NR	638
Okkyung	2002	Moderate	Clinical trial	Korea	Prospective	Random	One site	2002	Children, Adolescents, Adults, Elderly	54.7	30.5	NR	Catarrhal, Suppurated, Gangrenous	NR	0	50	27.3	100	Open Surgery	NR	General	NR	84

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Onieva	2017	Moderate	Cross sectional	Spain	Retrospective	Consecutive	One site	2012-2014	Children, Adolescents, Adults, Elderly	53.7	32	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	NR	NR	294
Page	2010	Moderate	Cross sectional	USA	Retrospective	Exhaustive	Multisite	2008	Adults	51.4	39.2	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	0.94	General	NR	17199
Palesty	2004	Moderate	Cross sectional	USA	Retrospective	Consecutive	One site	2000-2002	Adults	47	25.2	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy	1.2	General	NR	50
Pandit	2016	High	Cohort	Nepal	Retrospective	Consecutive	Multisite	2009-2014	Children, Adolescents, Adults	51	24.3	NR	Perforated, Suppurated	NR	2.6	97.4	0		Open Surgery	0.6	Spinal and General	NR	101
Parcells	2009	Low	Cohort	USA	Retrospective	Systematic	One site	1997-2007	Adults	NR	39.3	NR	Perforated, Not perforated	NR	33.1	NR	NR		Laparoscopy or Open Surgery	NR	NR	According to CDC-NNIS diagnostic criteria	1063
Park	2017	Low	Cohort	Korea	Prospective	Systematic	One site	2012-2014	Adults	53.3	37.45	NR	Perforated, Gangrenous	NR	38.7	NR	45.9		Laparoscopy	1.1	General	According to CDC-NNIS diagnostic Criteria	1343
Park	2018	Moderate	Cohort	Korea	Retrospective	Consecutive	One site	2009-2013	Adults	53.7	34.2	NR	Perforated	NR	13.2	NR	NR		Laparoscopy with Open Surgery	1.1	General	NR	986
Pascual	2017	Moderate	Cohort	Spain	Prospective	Consecutive	One site	2013-2017	Adults	49	41	NR	Unclear	NR	NR	NR	NR		Laparoscopy	0.1	General	NR	100
Patel	2003	High	Cohort	Kenya	Retrospective	Consecutive	One site	1996-2002	Children, Adolescents, Adults	30.2	30.6	NR	Catarrhal, Suppurated, Gangrenous, Carcinoid tumor	94.3	0	0.9	2.8	100	Laparoscopy with Open Surgery	1.5	General	NR	106
Pearcy	2017	Moderate	Case control	USA	Retrospective	Random	Multisite	2010-2014	Adults	54	36	NR	Unclear	NR	NR	NR	NR		NR	1.1	NR	NR	2903
Pishori	2003	Low	Cross sectional	Pakistan	Prospective	Systematic	One site	1997-1999	Children, Adolescents, Adults, Elderly	NR	NR	NR	Unclear	NR	NR	NR	NR		NR	NR	NR	According to CDC-NNIS diagnostic criteria	3304
Putnam	2016	Moderate	Cross sectional	USA	Prospective	Consecutive	One site	2012-2015	Children, Adolescents	61	9.4	NR	Perforated, Suppurated, Gangrenous	0	100	0	0	100	Laparoscopy or Open Surgery	0.9	General	NR	410
Qahtani	2014	Moderate	Cohort	Saudi Arabia	Prospective	Random	One site	2012	Adolescents, Adults	68	23.6	NR	Catarrhal, Perforated, Gangrenous	9.6	19.9	NR	22.4	100	Open Surgery	1.5	General	NR	91
Quezada	2015	Moderate	Cohort	Chile	Retrospective	Consecutive	One site	2003-2013	Adults	43	39	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	2.2	NR	NR	227
Raakow	2014	High	Cohort	Germany	Prospective	Not clear	One site	2009-2013	Adolescents, Adults	28.8	27.2	NR	Catarrhal, Suppurated, Gangrenous	12.8	0	16	4		Laparoscopy	0.8	General	NR	156
Rafiq	2015	Low	Clinical trial	Pakistan	Prospective	Random	One site	2012-2014	Adolescents, Adults, Elderly	48.5	22.6	0	Unclear	NR	NR	NR	NR	100	Open Surgery	0.7	General	NR	390
Rajabi-Mashhadi	2012	Moderate	Clinical trial	Iran	Prospective	Random	One site	2006-2007	Adults	62.5	26.2	NR	Unclear, Non perforated	NR	NA	NR	NR	100	Open Surgery	NR	NR	NR	291

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Reinisch	2017	Moderate	Cross sectional	Germany	Retrospective	Consecutive	One site	2008-2015	Adults	56	32	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy with Open Surgery	NR	General	NR	680
Rios	2004	High	Cross sectional	Peru	Not reported/Unclear	Consecutive	One site	2001-2002	Children, Adolescents, Adults, Elderly	NR	30.6	NR	Catarrhal, Perforated, Suppurated, Gangrenous	15.38	16.35	44.23	24.04	100	NR	0.98	Unclear	NR	104
Romano	2014	Moderate	Cross sectional	USA	Retrospective	Systematic	One site	2010-2012	Adults	66	35.7	NR	Gangrenous	NR	NR	NR	9.7	86	Laparoscopy	NR	General	NR	372
Romel Hilaire	2014	Moderate	Cross sectional	Cuba	Retrospective	Consecutive	One site	2007-2009	Adults	100	NR	NR	Suppurated	0	0	100	0		Laparoscopy or Open Surgery	NR	NR	NR	720
Romeo	2009	Moderate	Cross sectional	Colombia	Retrospective	Consecutive	One site	1997	Children, Adolescents, Adults, Elderly	NR	NR	NR	Unclear	NR	NR	NR	NR		Laparoscopy	NR	NR	NR	310
Romy	2008	Low	Cross sectional	Switzerland	Prospective	Systematic	Multisite	1998-2004	Children, Adolescents, Adults, Elderly	53.9	32.7	NR	Unclear	NR	NR	NR	NR	59.5	Laparoscopy or Open Surgery	NR	NR	According to CDC-NNIS diagnostic criteria	2468
Rooh-ul-Muqim	2010	Moderate	Cohort	Pakistan	Prospective	Consecutive	One site	2008-2009	Adolescents, Adults, Elderly	48.5	24	NR	Catarrhal, Perforated, Suppurated, Gangrenous	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	0.5	General	NR	165
Rossem	2015	Moderate	Cohort	Netherlands	Prospective	Consecutive	Multisite	2014	Adults	47.5	44	NR	Perforated, Gangrenous	NR	68	10.4	21.7	100	Laparoscopy with Open Surgery	0.9	General	NR	415
Rossem	2016	Low	Cohort	Netherlands	Prospective	Not clear	Multisite	2014	Children, Adolescents, Adults	46.2	28.0	NR	Unclear	NR	NR	NR	NR		Laparoscopy	0.8	General	Superficial surgical site infection: recorded when administration of antibiotics, opening of the incision or both was necessary. An intra-abdominal abscess was defined as a postoperative intra-abdominal fluid collection diagnosed by cross-sectional imaging for which administration of antibiotics or a radiological or surgical intervention was needed.	1995
Rotermann	2004	Moderate	Cohort	Canada	Retrospective	Consecutive	Multisite	1997-2000	Children, Adolescents, Adults, Elderly	55.2	NR	NR	Unclear	NR	NR	NR	NR		NR	NR	NR	According to CDC-NNIS diagnostic criteria	80867

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Saar	2016	Low	Cross sectional	Estonia	Prospective	Consecutive	One site	2013-2014	Adults	48.9	35.4	NR	Perforated, Gangrenous	NR	15.4	NR	59.4	95.1	Laparoscopy or Open Surgery	0.7	General	According to CDC-NNIS diagnostic Criteria	266
Saber	2010	Moderate	Clinical trial	USA	Prospective	Consecutive	One site	2008-2009	Adults	42.3	33	NR	Catarrhal	100	0	0	0		Laparoscopy	0.8	NR	NR	26
Sadraei-Mosavi	2017	Moderate	Clinical trial	Iran	Prospective	Random	One site	2013-2014	Adults	NR	28.4	NR	Catarrhal	100	0	0	0	100	Open Surgery	NR	NR	SSI=pus discharge from wound, redness, tenderness, edema	152
Saha	2010	Moderate	Cohort	Bangladesh	Prospective	Consecutive	One site	2007-2008	Children	NR	NR	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	NR	NR	NR	60
Sahm	2010	Moderate	Cross sectional	Germany	Prospective	Systematic	One site	1998-2006	Adults	54	39	NR	Catarrhal, Perforated, Gangrenous	50.7	17.0	NR	6.9	100	Laparoscopy or Open Surgery	1.0	General	NR	1710
Sahm	2015	Moderate	Cross sectional	Germany	Prospective	Exhaustive	Multisite	1988-2009	Children, Adolescents, Adults, Elderly	43	31	NR	Perforated, Non Perforated	91.5	NR	8.5	NR	NR	Laparoscopy or Open Surgery	NR	NR	NR	12570
Saló	2016	High	Cohort	Sweden	Retrospective	Consecutive	One site	2006-2014	Children	55.6	10.4	NR	Perforated, Gangrenous, Phlegmonous	NR	7.3	NR	11.6	100	Laparoscopy	0.94	NR	NR	259
Sanchez-Santana	2017	Low	Cohort	Spain	Prospective	Consecutive	One site	2007-2015	Adults	55.2	32.9	2.6	Unclear	NR	NR	NR	NR	71.3	Laparoscopy or Open Surgery	NR	NR	According to CDC-NNIS diagnostic criteria	930
Sauvain	2016	Moderate	Cohort	Switzerland	Retrospective	Consecutive	Multisite	2007-2011	Adults	53.2	34	NR	Catarrhal, Perforated, Suppurated, Gangrenous	NR	19	NR	NR		Laparoscopy or Open Surgery	NR	NR	NR	2559
Scarborough	2012	Low	Cross sectional	USA	Retrospective	Systematic	Multisite	2005-2009	Children, Adolescents, Adults, Elderly	52	38.9	NR	Perforated, Non rupture	NR	11.2	NR	NR		Laparoscopy or Open Surgery	0.9	NR	According to CDC-NNIS diagnostic criteria	39122
Scarless	2013	Moderate	Clinical trial	Scotland	Prospective	Random	One site	2011	Adults	53	32	NR	Unclear	NR	NR	NR	NR		Laparoscopy	1.4	General	NR	38
Seifarth	2016	Moderate	Cohort	USA	Retrospective	Consecutive	Multisite	2007-2012	Children, Adults	60	12	NR	Catarrhal	100	0	0	0		Laparoscopy	NR	General	NR	1283
Seifarth	2016	Low	Cohort	USA	Retrospective	Consecutive	Multisite	2007-2012	Children, Adolescents, Adults	60	12	NR	Perforated, Suppurated, Gangrenous	100	0	0	0		Laparoscopy	NR	NR	NR	1283
Senekjan	2013	Moderate	Cohort	USA	Retrospective	Consecutive	Multisite	2005-2009	Adolescents	56.5	40.3	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	0.9	NR	1) SSI (superficial and deep incisional)...infection within 30 days of operation and involved skin, subcutaneous tissue or deep soft tissue 2) Organ space infection (OSI)...infection within 30 days of operation when the infection appeared to be related to the operation and involved any part of the anatomy other than the incision	61830

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Seo	2002	Moderate	Cross sectional	Korea	Retrospective	Systematic	One site	2000	Adults	0	NR	NR	Catarrhal, Perforated, Suppurated, Gangrenous	14.7	15.6	49.5	20.2		NR	NR	NR	NR	129
Sesia	2011	Moderate	Cohort	Germany	Prospective	Consecutive	One site	2006-2008	Children, Adolescents, Adults, Elderly	NR	NR	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	1	NR	NR	265
Shaikg	2011	Moderate	Cross sectional	Pakistan	Prospective	Consecutive	One site	2007-2009	Adults	51.4	29	NR	Catarrhal, Perforated, Suppurated	82.86	8.67	1.51	0		Open Surgery	NR	General	NR	461
Shang	2017	Moderate	Cohort	China	Retrospective	Consecutive	One site	2013-2016	Adults	54.3	2.2	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	NR	Erythema, swelling and purulent discharge	398
Shimizu	2014	Low	Cross sectional	Japan	Retrospective	Not clear	One site	2000-2012	Adults	44	35	NR	Catarrhal, Gangrenous	19	NR	NR	37		Open Surgery	NR	General	According to CDC-NNIS diagnostic Criteria	300
Shindholimath	2011	Moderate	Cross sectional	India	Retrospective	Consecutive	One site	2007-2009	Adults	68.4	NR	NR	Perforated, Suppurated, Gangrenous, Appendicular abscess	0	36.8	5.3	26.3	100	Laparoscopy	1.6	General	NR	19
Siam	2017	Moderate	Cohort	Israel	Retrospective	Consecutive	One site	2008-2015	Adults	62.8	34.1	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy with Open Surgery	0.7	General	NR	1649
Silva	2008	Moderate	Cohort	Chile	Prospective	Random	One site	2005-2006	Adults	58.9	NR	NR	Unclear	NR	NR	NR	NR		Open Surgery	NR	General	NR	433
Singh	2017	Moderate	Clinical trial	India	Prospective	Consecutive	One site	2014-2015	Adults	43.2	28.7	11.4	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	General	NR	44
Siribumrungwong	2013	Low	Cohort	Thailand	Retrospective	Systematic	One site	2006	Adults	65	37	NR	Perforated	NR	100	NR	NR	100	Open Surgery	1.2	NR	According to CDC-NNIS diagnostic criteria	128
Sivrikoz	2015	Moderate	Cohort	USA	Retrospective	Exhaustive	Multisite	2004-2010	Children, Adolescents, Adults, Elderly	52.1	48	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	0.9	NR	NR	4844
Soll	2016	Low	Cohort	Switzerland	Retrospective	Consecutive	One site	2009-2013	Children, Adolescents, Adults, Elderly	54.7	26.5	NR	Catarrhal, Perforated, Suppurated, Gangrenous	NR	46	NR	NR	100	Laparoscopy	1	NR	According to CDC-NNIS diagnostic Criteria	813
Sozutek	2013	Low	Clinical trial	Turkey	Retrospective	Consecutive	One site	2010-2011	Adults	44	30.9	NR	Catarrhal, Perforated	57	20	NR	NR		Laparoscopy or Open Surgery	0.5	General	NR	75
Shishewachart	2016	Moderate	Cross sectional	Thailand	Retrospective	Consecutive	One site	2012-2014	Children, Adolescents, Adults, Elderly	52	43.7	7.4	Unclear	NR	NR	NR	NR		Open Surgery	NR	General	NR	450
Staszewicz	2014	Moderate	Cohort	Switzerland	Prospective	Systematic	Multisite	1998-2011	Children, Adolescents, Adults, Elderly	54	34.2	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	1	NR	According to CDC-NNIS diagnostic Criteria	6383
Suttie	2004	High	Case control	Scotland	Retrospective	Not clear	One site	1997-2002	Children	50	10.8	NR	Perforated, Suppurated, Gangrenous	0	2	50	14		Laparoscopy	1	General	NR	60
Svensson	2016	Moderate	Cohort	Sweden	Prospective	Consecutive	One site	2006-2010	Children, Adolescents	60.2	11.3	NR	Catarrhal, Perforated, Suppurated, Gangrenous, not described	6.6	21.8	44.6	29.8	100	Laparoscopy or Open Surgery	0.7	General	NR	1745

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Taguchi	2015	Moderate	Clinical trial	Japan	Prospective	Random	One site	2009-2014	Adults	65.43	47.5	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	1.2	General	According to CDC-NNIS diagnostic Criteria	81
Tanaka	2007	Moderate	Cohort	Japan	Retrospective	Consecutive	One site	2002-2005	Children	54.3	2.2	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	NR	NR	95
Tijerina	2010	Low	Clinical trial	Mexico	Prospective	Exhaustive	One site	2005-2007	Children, Adolescents, Adults, Elderly	46	NR	NR	Unclear	NR	NR	NR	NR	100	Open Surgery	NR	General	NR	529
Toro Pablo	2017	Moderate	Cohort	Spain	Retrospective	Consecutive	One site	2012-2016	Children, Adolescents, Adults	NR	26	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	NR	NR	600
Towfigh	2011	Low	Clinical trial	USA	Prospective	Random	One site	2007-2009	Adults	77.3	33	NR	Perforated	0	100	0	0		Open Surgery	NR	NR	NR	75
Troillet	2017	Low	Cohort	Switzerland	Prospective	Consecutive	Multisite	2011-2015	Children, Adolescents, Adults, Elderly	NR	NR	NR	Unclear	NR	NR	NR	NR	92.2	Laparoscopy or Open Surgery	NR	NR	According to CDC-NNIS diagnostic Criteria	15439
Tsioplis	2013	Moderate	Cross sectional	Germany	Retrospective	Consecutive	One site	1999-2008	Children, Adolescents, Adults, Elderly	51	23	9	Catarrhal, Perforated, Suppurated, Gangrenous	19	NR	50	25	75	Laparoscopy or Open Surgery	NR	Not reported	NR	1439
Vahdad	2016	Moderate	Cross sectional	Germany	Retrospective	Systematic	One site	2008-2012	Children, Adolescents	52.4	NR	NR	Catarrhal, Perforated, Phelgmonous in 43% of cases	48.2	8.7	NR	NR		Laparoscopy	1.1	NR	NR	309
Van Rossem	2016	High	Cohort	Netherlands	Prospective	Consecutive	Multisite	2014	Adults	49.7	39.0	NR	Catarrhal, Perforated, Gangrenous	73.7	11.0	NR	9.9	96.6	Laparoscopy or Open Surgery	0.72	NR	NR	1378
Van Rossem	2014	High	Cohort	Netherlands	Retrospective	Consecutive	Multisite	2004-2010	Adults	53.2	49	NR	Perforated	0	100	0	0	100	Laparoscopy or Open Surgery	0.85	NR	NR	267
Wang-Chan	2017	Low	Cross sectional	Switzerland	Retrospective	Consecutive	One site	2013-2014	Children, Adolescents, Adults, Elderly	55.3	47	13.8	Unclear	NR	NR	NR	NR		Laparoscopy with Open Surgery	NR	General	According to CDC-NNIS diagnostic Criteria	246
Watanabe	2011	Low	Cross sectional	Japan	Prospective	Consecutive	Multisite	2005-2006	Adults	59.4	63.8	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	General	According to CDC-NNIS diagnostic Criteria	903
Willis	2016	Moderate	Cohort	USA	Prospective	Consecutive	One site	2013-2014	Children, Adolescents	58.5	8.8	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	NR	General	NR	313
Wong	2015	High	Cohort	Peru	Prospective	Not clear	Multisite	2005-2010	Adults	NR	NR	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	General	NR	352
Wu	2006	Low	Cross sectional	Taiwan	Retrospective	Not clear	One site	2001-2005	Adults	75	42	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	0.95	General	NR	1795

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Wu	2011	Moderate	Cohort	Taiwan	Retrospective	Exhaustive	Multisite	2004-2009	Children, Adolescents, Adults, Elderly	58.1	36.4	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	General	NR	1366
Wu	2017	Moderate	Cross sectional	China	Retrospective	Consecutive	One site	2014-2016	Elderly	59	71	NR	Perforated, Suppurated, Gangrenous	0	61.7	10	28.7	100	Laparoscopy or Open Surgery	1	General	NR	115
Wu	2014	Moderate	Clinical trial	China	Prospective	Random	One site	2011-2013	Children, Adolescents	60	8.5	NR	Catarrhal	100	0	0	0		Laparoscopy	1	General	NR	60
Yaghoubian	2010	High	Cross sectional	USA	Retrospective	Exhaustive	Multisite	1998-2007	Children, Adolescents, Adults, Elderly	61.5	29.2	NR	Catarrhal, Perforated	73.4	26.6	0	0		Laparoscopy or Open Surgery	NR	NR	NR	4325
Yagnik	2010	Moderate	Cross sectional	India	Retrospective	Consecutive	One site	2007-2009	Children, Adolescents, Adults	32.5	23.41	NR	Catarrhal	100	0	0	0	100	Laparoscopy or Open Surgery	1	General	NR	151
Yousef	2017	Moderate	Cohort	Canada	Prospective	Consecutive	One site	2015-2016	Children, Adolescents	63.1	9.3	NR	Perforated	0	100	0	0	100	Laparoscopy	NR	General	NR	122
Zhang	2015	Moderate	Clinical trial	China	Prospective	Random	One site	2012-2013	Adults	47.2	30.8	NR	Unclear	10.2	7.4	54.6	9.3		Laparoscopy	0.9	General	NR	108

NR: not reported

References

1. Adejumo A.A., N.M., Mshelia, Y.M., Saleh. Clinicopathological presentation and management outcome of appendicitis in gombe, north-east nigeria: a 7-year retrospective audit. *Niger J Med.* 2015;24(4):337-43.
2. Aguiló J., S., Peiró, C., Muñoz, J., García del Caño, M., Garay, V., Viciano, et al. Adverse outcomes in the surgical treatment of acute appendicitis. *Efectos adversos en la cirugía de la apendicitis aguda.* 2005;78(5):312-7.
3. Adhikary S., S., Tyagi, G., Sapkota, A., Afaq, B.K., Bhattarai, C.S., Agrawal. Port exteriorization appendectomy: is it the future? *Nepal Med Coll J.* 2008;10(1):30-4.
4. Ahmad M., K., Ali, H., Latif, S., Naz, K., Said. Comparison of primary wound closure with delayed primary closure in perforated appendicitis. *J Ayub Med Coll Abbottabad.* 2014;26(2):153-7.
5. Ahmed I., J., Burr, M., Castillo, D., Collins, J.A., Cook, M., Campbell, et al. Single port/incision laparoscopic surgery compared with standard three-port laparoscopic surgery for appendectomy: A randomized controlled trial. *Surg Endosc Interv Tech.* 2015;29(1):77-85.
6. Akkoyun I., A., Taş Tuna. Advantages of abandoning abdominal cavity irrigation and drainage in operations performed on children with perforated appendicitis. *J Pediatr Surg.* 2012;47(10):1886-90.
7. Al-Saadi A.S., A.H., Al-Wadan, S.A., Hamarnah, H., Amin. Is abandoning routine peritoneal cultures during appendectomy justified? *Saudi Med J.* 2007;28(12):1827-9.
8. Al-Temimi M.H., M.A., Berglin, E.G., Kim, D.J., Tessier, S.D., Johna. Endostapler versus Hem-O-Lok clip to secure the appendiceal stump and mesoappendix during laparoscopic appendectomy. *Am J Surg.* 2017;214(6):1143-8.

- 1
2
3 9. Ali N., S., Aliyu. Appendicitis and its surgical management experience at the University of
4 Maiduguri Teaching Hospital Nigeria. *Niger J Med.* 2012;21(2):223-6.
5
6
7
- 8
9 10. Ali K., H., Latif, S., Ahmad. Frequency of wound infection in non-perforated appendicitis
10 with use of single dose preoperative antibiotics. *J Ayub Med Coll Abbottabad.*
11 2015;27(2):378-80.
12
13
- 14
15 11. Almström M., J.F., Svensson, B., Patkova, A., Svenningsson, T., Wester. In-hospital
16 surgical delay does not increase the risk for perforated appendicitis in children. *Ann Surg.*
17 2017;265(3):616-21.
18
19
- 20
21 12. Álvarez-Moreno C., A.M., Pérez-Fernández, V.D., Rosenthal, J., Quintero, E., Chapeta-
22 Parada, C., Linares, et al. Surgical site infection rates in 4 cities in Colombia: Findings of
23 the International Nosocomial Infection Control Consortium (INICC). *Am J Infect Control.*
24 2014;42(10):1089-92.
25
26
- 27
28 13. Andert A., H.P., Alizai, C.D., Klink, N., Neitzke, C., Fitzner, C., Heidenhain, et al. Risk
29 factors for morbidity after appendectomy. *Langenbeck's Arch Surg.* 2017;402(6):987-93.
30
31
- 32
33 14. Andersson R.E. Short-term complications and long-term morbidity of laparoscopic and
34 open appendectomy in a national cohort. *Br J Surg.* 2014;101(9):1135-42.
35
36
- 37
38 15. Aranda-Narváez J.M., A.J., González-Sánchez, N., Marín-Camero, C., Montiel-Casado, P.,
39 López-Ruiz, B., Sánchez-Pérez, et al. Conservative approach versus urgent appendectomy
40 in surgical management of acute appendicitis with abscess or phlegmon. *Resultados del*
41 *tratamiento conservador inicial y de la cirugía urgente en la apendicitis aguda evolucionada.*
42 2010;102(11):648-52.
43
44
- 45
46 16. Aranda-Narváez J.M., T., Prieto-Puga Arjona, B., García-Albiach, M.C., Montiel-Casado,
47 A.J., González-Sánchez, B., Sánchez-Pérez, et al. Postappendectomy surgical site infection:
48 Overall rate and type according to open/laparoscopic approach. *Infección de sitio quirúrgico*
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 tras apendicectomía urgente: tasa global y tipo según la vía de abordaje
4 (abierta/laparoscópica). 2014;32(2):76-81.
5
6
7
8

- 9 17. Arthur T., R., Gartrell, B., Manoharan, D., Parker. Emergency appendectomy in Australia:
10 findings from a multicentre, prospective study. ANZ J Surg. 2017;87(9):656-60.
11
12
13
14 18. Asefa Z. Acute appendicitis in Yirgalem Hospital, southern Ethiopia. Ethiop Med J.
15 2002;40(2):155-62.
16
17
18
19 19. Assefa Z., A., G/yesuse. Acute appendicitis in children admitted to zewditu memorial
20 hospital. Ethiop Med J. 2014;52(4):189-95.
21
22
23
24 20. Atif M.L., F., Sadaoui, A., Bezzaoucha, C.A., BezzaouchaKaddache, R., Boukari, S.,
25 Djelato, et al. Intra-abdominal abscesses and laparoscopic versus open appendectomies.
26 Infect Control Hosp Epidemiol. 2009;30(7):713-5
27
28
29
30
31
32 21. Bae E., A., Dehal, V., Franz, M., Joannides, N., Sakis, J., Scurlock, et al. Postoperative
33 antibiotic use and the incidence of intra-abdominal abscess in the setting of suppurative
34 appendicitis: a retrospective analysis. Am J Surg.2016;212(6):1121-5.
35
36
37
38
39
40 22. Bae S.U., W.K., Jeong, S.K., Baek. Single-port laparoscopic interval appendectomy for
41 perforated appendicitis with a periappendiceal abscess. Ann Coloproctol. 2016;32(3):105-
42 10.
43
44
45
46
47 23. Baek H.N., Y.H., Jung, Y.H., Hwang. Laparoscopic versus open appendectomy for
48 appendicitis in elderly patients. J Korean Soc Coloproctology. 2011;27(5):241-5.
49
50
51
52
53 24. Bali İ., F., Karateke, S., Özyazıcı, A., Kuvvetli, C., Oruç, E., Menekşe, et al. Comparison
54 of intracorporeal knotting and endoloop for stump closure in laparoscopic appendectomy.
55 Laparoskopik appendektomide intrakorporal düğüm ve endoloop ile güdük kapama
56 yöntemlerinin karşılaştırılması. 2015;21(6):446-9.
57
58
59
60

- 1
2
3
4
5 25. Bhangu A., Richardson, C., Torrance, A., Pinkney, T., Collaborative, Natl Surg Res.
6 Multicentre observational study of performance variation in provision and outcome of
7 emergency appendicectomy. *British Journal Of Surgery*. 2013;100(9):1240-52.
8
9
10
11
12 26. Bansal V., S., Altermatt, D., Nadal, C., Berger. Lack of benefit of preoperative antimicrobial
13 prophylaxis in children with acute appendicitis: A prospective cohort study. *Infection*.
14 2012;40(6):635-41.
15
16
17
18
19 27. Batajoo H., N.K., Hazra. Laparoscopic versus open appendectomy in acute appendicitis. *J*
20 *Nepal Health Res Counc*. 2012;10(22):239-42.
21
22
23
24
25 28. Saranga Bharathi R., V., Sharma, A., Chakladar, P., Kumari. Port exteriorisation
26 appendectomy-our experience. *Med J Armed Forces India*. 2011;67(2):147-51.
27
28
29
30
31 29. Biçakci U., B., Tander, M., Günaydin, R., Rizalar, E., Aritürk, S.H., Ayyildiz, et al. The
32 comparison of open and laparoscopic appendectomy: Is there any outcome difference
33 between non-complicated and complicated appendicitis? *Balkan Med J*. 2011;28(3):304-6.
34
35
36
37
38 30. Bickel A., M., Gurevits, R., Vamos, S., Ivry, A., Eitan. Perioperative hyperoxygenation and
39 wound site infection following surgery for acute appendicitis : A randomized, prospective,
40 controlled trial. *Arch Surg*. 2011;146(4):464-70.
41
42
43
44
45
46 31. Blackwood B.P., C.D., Gause, J.C., Harris, C.M., Theodorou, I., Helenowski, T.B., Lautz,
47 et al. Overweight and obese pediatric patients have an increased risk of developing a
48 surgical site infection. *Surg Infect*. 2017;18(4):491-7.
49
50
51
52
53 32. Blakely M.L., R., Williams, M.S., Dassinger, J.W., Eubanks III, P., Fischer, E.Y., Huang,
54 et al. Early vs interval appendectomy for children with perforated appendicitis. *Arch Surg*.
55 2011;146(6):660-5.
56
57
58
59
60

- 1
2
3 33. Bonadio W., K., Rebillot, O., Ukwuoma, C., Saracino, A., Iskhakov. Management of
4 Pediatric Perforated Appendicitis: Comparing Outcomes Using Early Appendectomy
5 versus Solely Medical Management. *Pediatr Infect Dis J.* 2017;36(10):937-41.
6
7
8
9
10
11 34. Boomer L.A., J.N., Cooper, K.J., Deans, P.C., Minneci, K., Leonhart, K.A., Diefenbach, et
12 al. Does delay in appendectomy affect surgical site infection in children with appendicitis?
13 *J Pediatr Surg.* 2014;49(6):1026-9.
14
15
16
17
18 35. Boomer L.A., J.N., Cooper, S., Anandalwar, S.C., Fallon, D., Ostlie, C.M., Leys, et al.
19 Delaying appendectomy does not lead to higher rates of surgical site infections. *Ann Surg.*
20 2016;264(1):164-8.
21
22
23
24
25
26 36. Bozkurt M.A., M.G., Ünsal, S., Kapan, B., Kankaya, M.U., Kalaycii, H., Aliiş. Two
27 different methods for appendiceal stump closure: Metal clip and Hem-o-lok clip. *J*
28 *Laparoendosc Adv Surg Techn.* 2014;24(8):571-3.
29
30
31
32
33 37. Brandt C., U., Hott, D., Sohr, F., Daschner, P., Gastmeier, H., Rüden. Operating room
34 ventilation with laminar airflow shows no protective effect on the surgical site infection rate
35 in orthopedic and abdominal surgery. *Ann Surg.* 2008;248(5):695-700.
36
37
38
39
40
41 38. Cairo S.B., M.V., Raval, M., Browne, H., Meyers, D.H., Rothstein. Association of same-
42 day discharge with hospital readmission after appendectomy in pediatric patients. *JAMA*
43 *Surg.* 2017;152(12):1106-12.
44
45
46
47
48 39. Cameron D.B., P., Melvin, D.A., Graham, C.C., Glass, S.K., Serres, M.P., Kronman, et al.
49 Extended Versus Narrow-spectrum Antibiotics in the Management of Uncomplicated
50 Appendicitis in Children: A Propensity-matched Comparative Effectiveness Study. *Ann*
51 *Surg.* 2017.
52
53
54
55
56
57 40. Cao J.-G., F., Tao, X.-J., Zhou, X.-G., Wang, S.-S., Wang, H., Zhang, et al. Trends and
58 outcomes of laparoscopic appendectomy in China: A multicenter, retrospective cohort
59
60

- 1
2
3 study. *Surg Pract.* 2015;19(4):166-72.
4
5
6
7
8 41. Cervantes-Sánchez C.R., R., Gutiérrez-Vega, J.A., Vázquez-Carpizo, P., Clark, C., Athié-
9 Gutiérrez. Syringe pressure irrigation of subdermic tissue after appendectomy to decrease
10 the incidence of postoperative wound infection. *World J Surg.* 2000;24(1):38-42.
11
12
13
14
15 42. Chamisa I. A clinicopathological review of 324 appendices removed for acute appendicitis
16 in Durban, South Africa: a retrospective analysis. *Ann R Coll Surg Engl.* 2009;91(8):688-
17 92.
18
19
20
21
22 43. Chaudhary I.A., Samiullah, A.A., Mallhi, Z., Afridi, A., Bano. Is it necessary to invaginate
23 the stump after appendectomy? *Pak J Med Sci.* 2005;21(1):35-8.
24
25
26
27
28 44. Chen D., H., Shi, H., Dong, K., Liu, K., Ding. Gasless single-incision laparoscopic
29 appendectomy. *Surg Endosc Interv Tech.* 2011;25(5):1472-6.
30
31
32
33
34 45. Chen C.-Y., Y.-C., Chen, H.-N., Pu, C.-H., Tsai, W.-T., Chen, C.-H., Lin. Bacteriology of
35 acute appendicitis and its implication for the use of prophylactic antibiotics. *Surg Infect.*
36 2012;13(6):383-90.
37
38
39
40
41 46. Chen C.-C., C.-T., Ting, M.-J., Tsai, W.-C., Hsu, P.-C., Chen, M.-D., Lee, et al.
42 Appendectomy timing: Will delayed surgery increase the complications? *J Chin Med*
43 *Assoc.* 2015;78(7):395-9.
44
45
46
47
48 47. Chiang R.-A., S.-L., Chen, Y.-C., Tsai, M.-J., Bair. Comparison of primary wound closure
49 versus open wound management in perforated appendicitis. *J Formos Med Assoc.*
50 2006;105(10):791-5.
51
52
53
54
55 48. Chiang R.-A., S.-L., Chen, Y.-C., Tsai. Delayed primary closure versus primary closure for
56 wound management in perforated appendicitis: A prospective randomized controlled trial.
57 *J Chin Med Assoc.* 2012;75(4):156-9.
58
59
60

- 1
2
3
4
5
6 49. Cho M., J., Kang, I.-K., Kim, K.Y., Lee, S.-K., Sohn. Underweight body mass index as a
7 predictive factor for surgical site infections after laparoscopic appendectomy. *Yonsei Med*
8 *J.* 2014;55(6):1611-6.
9
10
11
12
13 50. Choudhary S.K., S.K., Dhakaita. Appendicular mass-early appendicectomy vs interval
14 appendicectomy. *Intl J Pharma Bio Sci.* 2014;5(1):B400-B4.
15
16
17
18
19 51. Clyde C., T., Bax, A., Merg, M., MacFarlane, P., Lin, S., Beyersdorf, et al. Timing of
20 intervention does not affect outcome in acute appendicitis in a large community practice.
21 *Am J Surg.* 2008;195(5):590-3.
22
23
24
25
26 52. Coakley B.A., E.S., Sussman, T.S., Wolfson, A.S., Bhagavath, J.J., Choi, N.E., Ranasinghe,
27 et al. Postoperative antibiotics correlate with worse outcomes after appendectomy for
28 nonperforated appendicitis. *J Am Coll Surg.* 2011;213(6):778-83.
29
30
31
32
33
34 53. Crandall M., M.B., Shapiro, M., Worley, M.A., West. Acute uncomplicated appendicitis:
35 case time of day influences hospital length of stay. *Surg Infect (Larchmt).* 2009;10(1):65-
36 9.
37
38
39
40
41
42 54. Dede K., T., Mersich, A., Zaránd, I., Besznyák, Z., Baranyai, B., Atkári, et al. Laparoscopic
43 or open appendectomy? *Laparoszkópos vagy nyílt appendectomia?* 2008;149(50):2357-61.
44
45
46
47
48 55. Dhiman N., A., Chi, T.M., Pawlik, D.T., Efron, E.R., Haut, E.B., Schneider, et al. Increased
49 complications after appendectomy in patients with cerebral palsy: Are special needs patients
50 at risk for disparities in outcomes? *Surgery.* 2013;154(3):479-85.
51
52
53
54
55 56. Dimitriou I., B., Reckmann, O., Nephuth, M., Betzler. Single institution's experience in
56 laparoscopic appendectomy as a suitable therapy for complicated appendicitis.
57 *Langenbeck's Arch Surg.* 2013;398(1):147-52.
58
59
60

- 1
2
3 57. Durkin M.J., K.V., Dicks, A.W., Baker, S.S., Lewis, R.W., Moehring, L.F., Chen, et al.
4 Seasonal variation of common surgical site infections: Does season matter? *Infect Control*
5 *Hosp Epidemiol.* 2015;36(9):1011-6.
6
7
8
9
10
11 58. Ein S.H., A., Nasr, A., Ein. Open appendectomy for pediatric ruptured appendicitis: a
12 historical clinical review of the prophylaxis of wound infection and postoperative intra-
13 abdominal abscess. *Can J Surg.* 2013;56(3):E7-E12.
14
15
16
17
18 59. Fukuda H. Patient-related risk factors for surgical site infection following eight types of
19 gastrointestinal surgery. *J Hosp Infect.* 2016;93(4):347-54.
20
21
22
23
24 60. Gandaglia G., K.R., Ghani, A., Sood, J.R., Meyers, J.D., Sammon, M., Schmid, et al. Effect
25 of minimally invasive surgery on the risk for surgical site infections results from the national
26 surgical quality improvement program (nsqip) database. *JAMA Surg.* 2014;149(10):1039-
27 44.
28
29
30
31
32
33 61. Garcell H.G., A.V., Arias, C.A., Pancorbo Sandoval, E.G., García, M.E., Valle Gamboa,
34 A.B., Sado, et al. Incidence and etiology of surgical site infections in appendectomies: A 3-
35 year prospective study. *Oman Med J.* 2017;32(1):31-5.
36
37
38
39
40
41 62. Ghnnam W.M. Elderly versus young patients with appendicitis 3 years experience. *Alex J*
42 *Med.* 2012;48(1):9-12.
43
44
45
46
47 63. Giesen L.J., A.L., van den Boom, C.C., van Rossem, P.T., den Hoed, B.P., Wijnhoven.
48 Retrospective Multicenter Study on Risk Factors for Surgical Site Infections after
49 Appendectomy for Acute Appendicitis. *Dig Surg.* 2017;34(2):103-7.
50
51
52
53
54 64. Giiti G.C., H.D., Mazigo, J., Heukelbach, W., Mahalu. HIV, appendectomy and
55 postoperative complications at a reference hospital in Northwest Tanzania: Cross-sectional
56 study. *AIDS Res Ther.* 2010;7.
57
58
59
60

- 1
2
3 65. Golub A.V., R.S., Kozlov, V.G., Pleshkov, A.P., Moskalev, R.A., Alibegov, M.A.,
4 Chelombitko. Surgical Site Infections after Open Appendectomy and Effectiveness of
5 Complex Approach to Their Prevention. *Khirurgiia (Mosk)*. 2016(6):68-76.
6
7
8
9
10
11 66. Gross T.S., C., McCracken, K.F., Heiss, M.L., Wulkan, M.V., Raval. The contribution of
12 practice variation to length of stay for children with perforated appendicitis. *J Pediatr Surg*.
13 2016;51(8):1292-7.
14
15
16
17
18 67. Gurien L.A., D.L., Wyrick, S.D., Smith, M.S., Dassinger. Optimal timing of appendectomy
19 in the pediatric population. *J Surg Res*. 2016;202(1):126-31.
20
21
22
23
24 68. Hamzaoglu I, B., Baca, D.E., Böler, E., Polat, Y., Özer. Is umbilical flora responsible for
25 wound infection after laparoscopic surgery? *Surg Laparoscopy Endosc Percutaneous Tech*.
26 2004;14(5):263-7.
27
28
29
30
31
32 69. Harmon L.A., M.L., Davis, D.C., Jupiter, R.C., Frazee, J.L., Regner. Computed tomography
33 to operating room in less than 3 hours minimizes complications from appendicitis. *Am J*
34 *Surg*. 2016;212(2):246-50.
35
36
37
38
39 70. Helling T.S., D.F., Soltys, S., Seals. Operative versus non-operative management in the care
40 of patients with complicated appendicitis. *Am J Surg*. 2017;214(6):1195-200.
41
42
43
44
45 71. Helmer K.S., E.K., Robinson, K.P., Lally, J.C., Vasquez, K.L., Kwong, T.H., Liu, et al.
46 Standardized patient care guidelines reduce infectious morbidity in appendectomy patients.
47 *Am J Surg*. 2002;183(6):608-13.
48
49
50
51
52 72. Hesami M.A., H., Alipour, H., Nikoupour Daylami, B., Alipour, S., Bazargan-Hejazi, A.,
53 Ahmadi. Irrigation of abdomen with imipenem solution decreases surgical site infections in
54 patients with perforated appendicitis: A randomized clinical trial. *Iran Red Crescent MedJ*.
55 2014;16(4).
56
57
58
59
60

- 1
2
3 73. Horvath P., J., Lange, R., Bachmann, F., Struller, A., Königsrainer, M., Zdichavsky.
4 Comparison of clinical outcome of laparoscopic versus open appendectomy for complicated
5 appendicitis. *Surg Endosc Interv Tech.* 2017;31(1):199-205.
6
7
8
9
10
11 74. Hughes M.J., E., Harrison, S., Paterson-Brown. Post-operative antibiotics after
12 appendectomy and post-operative abscess development: A retrospective analysis. *Surg*
13 *Infect.* 2013;14(1):56-61.
14
15
16
17 75. Hussain M.I., M.K., Alam, H.H., Al-Qahatani, M.H., Al-Akeely. Role of postoperative
18 antibiotics after appendectomy in non-perforated appendicitis. *J Coll Phys Surg Pak.*
19 2012;22(12):756-9.
20
21
22
23
24
25
26 76. Iqbal M., M., Jawaid, A., Qureshi, S., Iqbal. Effect of povidone-iodine irrigation on post
27 appendectomy wound infection: Randomized control trial. *J Postgrad Med Inst.*
28 2015;29(3):160-4.
29
30
31
32
33 77. Iñigo J.J., B., Bermejo, B., Oronoz, J., Herrera, A., Tarifa, F., Pérez, et al. Surgical site
34 infection in general surgery: 5-year analysis and assessment of the National Nosocomial
35 Infection Surveillance (NNIS) index. *Infección de sitio quirúrgico en un servicio de cirugía*
36 *general Análisis de cinco años y valoración del índice National Nosocomial Infection*
37 *Surveillance (NNIS).* 2006;79(4):224-30.
38
39
40
41
42
43
44 78. Javadi S.M.R., S.Y., Zarghami, P., Ghaderzadeh, M., Ghorbanpoor, H.R., Makarchian, A.,
45 Derakhshanfar, et al. Comparison of small access and classic McBurney's incisions for open
46 appendectomy: A randomized controlled trial. *Shiraz E Med J.* 2017;18(10).
47
48
49
50
51 79. Jenkins P.C., M.K., Oerline, A.J., Mullard, M.J., Englesbe, D.A., Campbell, M.R.,
52 Hemmila. Hospital variation in outcomes following appendectomy in a regional quality
53 improvement program. *Am J Surg.* 2016;212(5):857-62.
54
55
56
57
58
59 80. Kang J., B.N., Bae, G., Gwak, I., Park, H., Cho, K., Yang, et al. Comparative study of a
60

1
2
3 single-incision laparoscopic and a conventional laparoscopic appendectomy for the
4 treatment of acute appendicitis. *J Korean Soc Coloproctology*. 2012;28(6):304-8.
5
6
7
8

9 81. Kapischke M., A., Pries, A., Caliebe. Short term and long term results after open vs.
10 Laparoscopic appendectomy in childhood and adolescence: A subgroup analysis. *BMC*
11 *Pediatr*. 2013;13(1).
12
13
14

15
16 82. Karam P.A., A., Hiuser, D., Magnuson, F.G.F., Seifarth. Intracorporeal hybrid single port
17 vs conventional laparoscopic appendectomy in children. *Pediatr Med Chir*. 2016;38(3):89-
18 92.
19
20
21
22

23
24 83. Karam P.A., A., Mohan, M.R., Buta, F.G., Seifarth. Comparison of Transumbilical
25 Laparoscopically Assisted Appendectomy to Conventional Laparoscopic Appendectomy in
26 Children. *Surg Laparoscopy Endosc Percutaneous Tech*. 2016;26(6):508-12.
27
28
29

30
31 84. Kasatpibal N., S., Jamulitrat, V., Chongsuvivatwong. Standardized incidence rates of
32 surgical site infection: A multicenter study in Thailand. *Am J Infect Control*.
33 2005;33(10):587-94.
34
35
36
37

38
39 85. Kasatpibal N., M., Nørgaard, H.T., Sørensen, H.C., Schönheyder, S., Jamulitrat, V.,
40 Chongsuvivatwong. Risk of surgical site infection and efficacy of antibiotic prophylaxis: A
41 cohort study of appendectomy patients in Thailand. *BMC Infect Dis*. 2006;6.
42
43
44

45
46 86. Kato Y., T., Marusasa, S., Ichikawa, G.J., Lane, T., Okazaki, A., Yamataka. Lapprotector
47 use decreases incisional wound infections in cases of perforated appendicitis: a prospective
48 study. *Asian J Surg*. 2008;31(3):101-3.
49
50
51
52

53
54 87. Kell M.R., K., Power, D.C., Winter, C., Power, C., Shields, W.O., Kirwan, et al. Predicting
55 outcome after appendectomy. *Ir J Med Sci*. 2003;172(2):63-5.
56
57
58

59
60 88. Khan M.N., T., Fayyad, T.D., Cecil, B.J., Moran. Laparoscopic versus open appendectomy:

- 1
2
3 the risk of postoperative infectious complications. *JLS*. 2007;11(3):363-7.
4
5
6
7
8 89. Khan K.I., S., Mahmood, M., Akmal, A., Waqas. Comparison of rate of surgical wound
9 infection, length of hospital stay and patient convenience in complicated appendicitis
10 between primary closure and delayed primary closure. *J Pak Med Assoc*. 2012;62(8):596-
11 8.
12
13
14
15
16 90. Khan I., M.I., Khan, M., Jawed, U., Shaikh, S., Ahmed, A., Arif. To compare the frequency
17 of superficial surgical site infection after laparoscopic versus open appendectomy. *Med*
18 *Forum Monthly*. 2014;25(11):52-5.
19
20
21
22
23
24 91. Khiria L.S., R., Ardhari, N., Mohan, P., Kumar, R., Nambiar. Laparoscopic
25 appendectomy for complicated appendicitis: Is it safe and justified? A retrospective
26 analysis. *Surg Laparoscopy Endosc Percutaneous Tech*. 2011;21(3):142-5.
27
28
29
30
31 92. Kılıç Ş.S., S., Ekinçi, İ., Karnak, A.Ö., Çiftçi, F.C., Tanyel, M.E., Şenocak. Drainage
32 systems' effect on surgical site infection in children with perforated appendicitis. *Drenaj*
33 *Sistemlerinin perforé apandisitli çocuklarda cerrahi alan enfeksiyonuna etkisi*.
34 2016;7(5):591-4.
35
36
37
38
39
40 93. Kim M.J., F.J., Fleming, D.D., Gunzler, S., Messing, R.M., Salloum, J.R.T., Monson.
41 Laparoscopic appendectomy is safe and efficacious for the elderly: An analysis using the
42 National Surgical Quality Improvement Project database. *Surg Endosc Interv Tech*.
43 2011;25(6):1802-7.
44
45
46
47
48
49 94. Kim J.H., H.Y., Kim, S.K., Park, J.S., Lee, D.S., Heo, S.W., Park, et al. Single-incision
50 laparoscopic appendectomy versus conventional laparoscopic appendectomy: Experiences
51 from 1208 cases of single-incision laparoscopic appendectomy. *Ann Surg*.
52 2015;262(6):1054-8.
53
54
55
56
57
58 95. Kim J.K., J., Kang, W.R., Kim, E.J., Park, S.H., Baik, K.Y., Lee. Does Conversion
59
60

- 1
2
3 Adversely Impact the Clinical Outcomes for Patients with Complicated Appendicitis? J
4 Laparoendosc Adv Surg Techn. 2016;26(8):635-40.
5
6
7
8
9 96. Kiriakopoulos A., D., Tsakayannis, D., Linos. Laparoscopic management of complicated
10 appendicitis. JSLS. 2006;10(4):453-6.
11
12
13
14
15 97. Kirshtein B., Z.H., Perry, S., Mizrahi, L., Lantsberg. Value of laparoscopic appendectomy
16 in the elderly patient. World J Surg. 2009;33(5):918-22.
17
18
19
20
21 98. Kiudelis M., P., Ignatavicius, K., Zviniene, S., Grizas. Analysis of intracorporeal knotting
22 with invaginating suture versus endoloops in appendiceal stump closure. Wideochir Inne
23 Tech Ma?oinwazyjne. 2013;8(1):69-73.
24
25
26
27
28 99. Kleif J., L., Rasmussen, S., Fonnes, P., Tibæk, A., Daoud, H., Lund, et al. Enteral
29 Antibiotics are Non-inferior to Intravenous Antibiotics After Complicated Appendicitis in
30 Adults: A Retrospective Multicentre Non-inferiority Study. World J Surg.
31 2017;41(11):2706-14.
32
33
34
35
36
37
38 100. Koizumi N., H., Kobayashi, Y., Nakase, T., Takagi, K., Fukumoto. Efficacy of
39 transumbilical laparoscopic-assisted appendectomy for appendicitis: a four-year experience
40 at a single center. Surg Today. 2015;45(10):1245-9.
41
42
43
44
45 101. Shimizu T., M., Ishizuka, K., Kubota. The preoperative serum C-reactive protein level
46 is a useful predictor of surgical site infections in patients undergoing appendectomy. Surg
47 Today. 2015;45(11):1404-10.
48
49
50
51
52
53 102. Kumamoto K., H., Imaizumi, N., Hokama, T., Ishiguro, K., Ishibashi, K., Baba, et al.
54 Recent trend of acute appendicitis during pregnancy. Surg Today. 2015;45(12):1521-6.
55
56
57
58 103. Kumar B., A., Samad, T.W., Khanzada, M.H., Laghari, A.R., Shaikh. Superiority of
59 laparoscopic appendectomy over open appendectomy: The Hyderabad experience. Rawal
60

- 1
2
3 Med J. 2008;33(2):165-8.
4
5
6
7
8 104. Kumar S., A., Jalan, B.N., Patowary, S., Shrestha. Laparoscopic appendectomy versus
9 open appendectomy for acute appendicitis: A prospective comparative study. Kathmandu
10 Univ Med J. 2016;14(55):244-8.
11
12
13
14
15 105. Lacher M., O.J., Muensterer, G.R., Yannam, C.J., Aprahamian, L., Perger, M., Megison,
16 et al. Feasibility of single-incision pediatric endosurgery for treatment of appendicitis in
17 415 children. J Laparoendosc Adv Surg Techn. 2012;22(6):604-8.
18
19
20
21
22 106. Lasses-Martínez B., E., Ortiz-Oshiro, J.L., Cabañas-Ojeda, P., Benito-Expósito, C.,
23 Fernández-Pérez, J., Alvarez Fernández-Represa. Cost is not a drawback to perform
24 laparoscopic appendectomy in an academic hospital. Surg Laparoscopy Endosc
25 Percutaneous Tech. 2014;24(4):e123-e7.
26
27
28
29
30
31
32 107. Le D., W., Rusin, B., Hill, J., Langell. Post-operative antibiotic use in nonperforated
33 appendicitis. Am J Surg.2009;198(6):748-52.
34
35
36
37
38
39 108. Lee P., K., Waxman, B., Taylor, S., Yim. Use of wound-protection system and
40 postoperative wound-infection rates in open appendectomy: A randomized prospective
41 trial. Arch Surg. 2009;144(9):872-5.
42
43
44
45
46 109. Lee S.L., S., Shekherdimian, V.Y., Chiu. Comparison of pediatric appendicitis
47 outcomes between teaching and nonteaching hospitals. J Pediatr Surg. 2010;45(5):894-7.
48
49
50
51 110. Lee J.A., K.Y., Sung, J.H., Lee, D.S., Lee. Laparoscopic appendectomy with a single
52 incision in a single institute. J Korean Soc Coloproctology. 2010;26(4):260-4.
53
54
55
56
57 111. Lee S.-Y., H.-M., Lee, C.-S., Hsieh, J.-H., Chuang. Transumbilical laparoscopic
58 appendectomy for acute appendicitis: A reliable one-port procedure. Surg Endosc Interv
59
60

- 1
2
3 Tech. 2011;25(4):1115-20.
4
5
6
7 112. Lee S.L., A., Yaghoubian, A., Kaji. Laparoscopic vs open appendectomy in children:
8 Outcomes comparison based on age, sex, and perforation status. Arch Surg.
9 2011;146(10):1118-21.
10
11
12
13
14
15 113. Lee S.M., G.S., Hwang, D.S., Lee. Single-incision laparoscopic appendectomy using
16 homemade glove port at low cost. J Minimal Access Surg. 2016;12(2):124-8.
17
18
19
20
21 114. Levy S.M., G., Holzmann-Pazgal, K.P., Lally, K., Davis, L.S., Kao, K., Tsao. Quality
22 check of a quality measure: Surgical wound classification discrepancies impact risk-
23 stratified surgical site infection rates in pediatric appendicitis. J Am Coll Surg.
24 2013;217(6):969-73.
25
26
27
28
29
30 115. Li P., Q., Xu, Z., Ji, Y., Gao, X., Zhang, Y., Duan, et al. Comparison of surgical stress
31 between laparoscopic and open appendectomy in children. J Pediatr Surg. 2005;40(8):1279-
32 83.
33
34
35
36
37
38
39 116. Lim S.G., E.J., Ahn, S.Y., Kim, I.Y., Chung, J.-M., Park, S.H., Park, et al. A clinical
40 comparison of laparoscopic versus open appendectomy for complicated appendicitis. J
41 Korean Soc Coloproctology. 2011;27(6):293-7.
42
43
44
45
46 117. Geraldo José de Souza Lima, Silva, Alcino Lázaro da, Castro, Eduardo Godoy, Abras,
47 Gustavo Munayer, Pires, Lívio José Suretti, Leite, Rodrigo Fabiano Guedes. Efetividade e
48 segurança da apendicectomia videoassistida em porta única transumbilical em adolescentes
49 e adultos X1 Effectiveness and safeness of single-port trans-umbilical laparoscopic
50 appendectomy done in adolescents and adults. Revista do Colégio Brasileiro de Cirurgiões.
51 2008;35(4):244-51.
52
53
54
55
56
57
58
59 118. Lin H.-F., J.-M., Wu, L.-M., Tseng, K.-H., Chen, S.-H., Huang, I.-R., Lai. Laparoscopic
60

- Versus Open Appendectomy for Perforated Appendicitis. *J Gastrointest Surg.* 2006;10(6):906-10.
119. Litz C.N., S.M., Farach, P.D., Danielson, N.M., Chandler. Obesity and single-incision laparoscopic appendectomy in children. *J Surg Res.* 2016;203(2):283-6.
120. Liu C., W., Wang, Y., Sun, M., Xu, H., Zhuang, H., Chen, et al. Efficacy and complications of laparoscopic appendectomy for pediatric appendicitis. *Int J Clin Exp Med.* 2017;10(9):13784-9.
121. Mahmood M.M., A., Shahab, M.A., Razzaq. Surgical site infection in open versus laparoscopic appendectomy. *Pak J Med Health Sci.* 2016;10(3):1076-8.
122. Mbah N., W.Ek., Opara, N.P., Agwu. Waiting time among acute abdominal emergencies in a Nigerian teaching hospital: Causes of delay and consequences. *Niger J Surg Res.* 2006;8(1):69-73.
123. Memon G.A., A.I., Memon, S.K.A., Shah, R.A., Sahito, Habib-Ur-Rehman, S., Leghari, et al. An experience of treatment outcome in acute appendicitis with antibiotics and appendectomy at a tertiary care hospital. *Med Forum Monthly.* 2017;28(3):136-40.
124. Menezes M., L., Das, M., Alagtal, J., Haroun, P., Puri. Laparoscopic appendectomy is recommended for the treatment of complicated appendicitis in children. *Pediatr Surg Int.* 2008;24(3):303-5.
125. Merenda M., A., Litarski, P., Kabziński, D., Janczak. Laparoscopic appendectomy as an alternative to conventional procedure - results in our own material. *Pol Przegl Chir.* 2013;85(6):323-8.
126. Michailidou M., M.G., Sacco Casamassima, S.D., Goldstein, C., Gause, O., Karim, J.H., Salazar, et al. The impact of obesity on laparoscopic appendectomy: Results from the ACS

- 1
2
3 National Surgical Quality Improvement Program pediatric database. *J Pediatr Surg.*
4 2015;50(11):1880-4.
5
6
7
8
9 127. Michailidou M., S.D., Goldstein, M.G., Sacco Casamassima, J.H., Salazar, R., Elliott,
10 J., Hundt, et al. Laparoscopic versus open appendectomy in children: The effect of surgical
11 technique on healthcare costs. *Am J Surg.* 2015;210(2):270-5.
12
13
14
15
16 128. Mickovic I.N., Z., Golubovic, S., Mickovic, D., Vukovic, S., Trajkovic, S.S.,
17 Antunovic, et al. A comparative analysis of laparoscopic appendectomy in relation to the
18 open appendectomy in children. *Uporedna analiza laparoscopske apendektomije u odnosu*
19 *na otvorenu apendektomiju kod dece.* 2016;17(1):49-53.
20
21
22
23
24
25
26 129. Ming P.C., T.Y., Yee Yan, L.H., Tat. Risk factors of postoperative infections in adults
27 with complicated appendicitis. *Surg Laparoscopy Endosc Percutaneous Tech.*
28 2009;19(3):244-8.
29
30
31
32
33 130. Mingmalairak C., P., Ungbhakorn, V., Paocharoen. Efficacy of antimicrobial coating
34 suture coated polyglactin 910 with tricosan (Vicryl Plus) compared with polyglactin 910
35 (Vicryl) in reduced surgical site infection of appendicitis, double blind randomized control
36 trial, preliminary safety report. *J Med Assoc Thailand.* 2009;92(6):770-5.
37
38
39
40
41
42
43 131. Miyano G., T., Okazaki, Y., Kato, T., Marusasa, T., Takahashi, G.J., Lane, et al. Open
44 versus laparoscopic treatment for pan-peritonitis secondary to perforated appendicitis in
45 children: A prospective analysis. *J Laparoendosc Adv Surg Techn.* 2010;20(7):655-7.
46
47
48
49
50 132. Moazzez A., R.J., Mason, N., Katkhouda. Thirty-day outcomes of laparoscopic versus
51 open appendectomy in elderly using ACS/NSQIP database. *Surg Endosc Interv Tech.*
52 2013;27(4):1061-71.
53
54
55
56
57 133. Mohammad Taghi Rajabi-Mashhadi, Mousavi, Seyed Hadi, Khosravi-Mashizi, M. H.,
58 Ghayour-Mobarhan, Majid, Sahebkar, Amirhossein. Optimum duration of perioperative
59
60

- 1
2
3 antibiotic therapy in patients with acute non-perforated appendicitis: a prospective
4 randomized trial. *Asian Biomedicine*. 2012;6(6):891-4.
5
6
7
- 8
9 134. Monge Jodra V., A., Robustillo Rodela, F., Martin Martinez, N., López Fresneña, S.,
10 Oña Compán, F., Calbo Torrecillas, et al. Standardized infection ratios for three general
11 surgery procedures: A comparison between Spanish hospitals and U.S. centers participating
12 in the national nosocomial infections surveillance system. *Infect Control Hosp Epidemiol*.
13 2003;24(10):744-8.
14
15
16
17
18
- 19 135. Mueck K.M., L.R., Putnam, K.T., Anderson, K.P., Lally, K., Tsao, L.S., Kao. Does
20 compliance with antibiotic prophylaxis in pediatric simple appendicitis matter? *J Surg Res*.
21 2017;216:1-8.
22
23
24
25
26
27
- 28 136. Muensterer O.J., C., Puga Nougues, O.O., Adibe, S.R., Amin, K.E., Georgeson, C.M.,
29 Harmon. Appendectomy using single-incision pediatric endosurgery for acute and
30 perforated appendicitis. *Surg Endosc Interv Tech*. 2010;24(12):3201-4.
31
32
33
34
35
- 36 137. Muensterer O.J., R., Keijzer. A simple vacuum dressing reduces the wound infection
37 rate of single-incision pediatric endosurgical appendectomy. *J Soc Laparoendoscopic Surg*.
38 2011;15(2):147-50.
39
40
41
42
- 43 138. Mustafa M.I.T., S.M., Chaudhry, R.I.T., Mustafa. Comparison of early outcome
44 between patients of open appendectomy with and without drain for perforated appendicitis.
45 *Pak J Med Health Sci*. 2016;10(3):890-3.
46
47
48
49
- 50 139. Nadler E.P., K.K., Reblock, H.R., Ford, B.A., Gaines. Monotherapy versus multi-drug
51 therapy for the treatment of perforated appendicitis in children. *Surg Infect*. 2003;4(4):327-
52 33.
53
54
55
56
57
- 58 140. Nataraja R.M., A., Bandi, S.A., Clarke, M.J., Haddad. Comparison of intra-abdominal
59 abscess formation following laparoscopic and open appendectomy in children. *J*
60

- Laparoendosc Adv Surg Techn. 2010;20(4):391-4.
141. Nataraja R.M., W.J., Teague, J., Galea, L., Moore, M.J., Haddad, T., Tsang, et al. Comparison of intraabdominal abscess formation after laparoscopic and open appendectomies in children. *J Pediatr Surg*. 2012;47(2):317-21.
142. Norton Pérez, Romero, Marcela, Castelblanco, María Isabel, Rodríguez, Emma Isabel. Infección del sitio operatorio de apendicectomías en un hospital de la orinoquia colombiana X1 Surgical site infection following appendectomy at a hospital in the Colombian Orinoco river basin (Colombian Orinoquia). *Revista Colombiana de Cirugía*. 2009;24(1):23-30.
143. Obayashi J., K., Ohyama, S., Manabe, K., Tanaka, H., Nagae, H., Shima, et al. Are there reliable indicators predicting post-operative complications in acute appendicitis? *Pediatr Surg Int*. 2015;31(12):1189-93.
144. Obinwa O., C., Peirce, M., Cassidy, T., Fahey, J., Flynn. A model predicting perforation and complications in paediatric appendectomy. *Int J Colorectal Dis*. 2015;30(4):559-65.
145. Ohene-Yeboah M., B., Togbe. An audit of appendicitis and appendectomy in Kumasi, Ghana. *West Afr J Med*. 2006;25(2):138-43.
146. Okkyung Suh, 신완균, 강성희, 양대현. Appropriate Duration of Prophylactic Antibiotics in Acute Nonperforated Appendicitis Z1 급성 비천공성 충수염 수술시 예방적 항균제의 사용기간. *Korean Journal of Clinical Pharmacy S1 한국임상약학회지*. 2002;12(2):65-70.
147. Francisco Gabriel Onieva, Roldán, Sara, Domínguez, José Ramón, Montero, Juan Pedro, Galnares, Alfonso, Peralta, Jordi. Abordaje laparoscópico frente a enfoque clásico

- 1
2
3 en el tratamiento de la apendicitis aguda X1 Laparoscopic approach versus classic open
4 procedure in the treatment of acute appendicitis. *Revista Colombiana de Cirugía*.
5 2017;32(1):26-31.
6
7
8
9
- 10 148. Page A.J., J.D., Pollock, S., Perez, S.S., Davis, E., Lin, J.F., Sweeney. Laparoscopic
11 Versus Open Appendectomy: An Analysis of Outcomes in 17,199 Patients Using
12 ACS/NSQIP. *J Gastrointest Surg*. 2010;14(12):1955-62.
13
14
15
16
- 17 149. Palesty J.A., X.J., Wang, R.C., Rutland, J., Leighton, S.J., Dudrick, A., Benbrahim.
18 Fifty-five consecutive laparoscopic appendectomy procedures without conversion. *JSLs*.
19 2004;8(2):141-5.
20
21
22
23
- 24 150. Pandit R.K. Safe and feasible time limit for early appendectomy in appendiceal mass.
25 *Kathmandu Univ Med J*. 2016;14(55):210-4.
26
27
28
29
- 30 151. Parcels J.P., J.P., Mileski, F.T., Gnagy, A.F., Haragan, W.J., Mileski. Using
31 antimicrobial solution for irrigation in appendicitis to lower surgical site infection rates. *Am*
32 *J Surg*. 2009;198(6):875-80.
33
34
35
36
37
38
39
40
- 41 152. Park H.-C., M.J., Kim, B.H., Lee. Effect of a Standardized Protocol of Antibiotic
42 Therapy on Surgical Site Infection after Laparoscopic Surgery for Complicated
43 Appendicitis. *Surg Infect*. 2017;18(6):684-8.
44
45
46
47
48
- 49 153. Seongmun Park, Park, Min-Su, Lee, Kil-Yeon. Relationship between the Hospital Visit-
50 to-Operation Time Interval and the Risk of Appendiceal Perforation and Clinical Outcomes.
51 *Journal of Minimally Invasive Surgery*. 2018;21(1):31-7.
52
53
54
- 55 154. Reoyo Pascual J.F., R., León Miranda, C., Cartón Hernández, E., Alonso Alonso, R.M.,
56 Martínez Castro, J., Sánchez Manuel. Laparoscopic appendectomy by 'glove port'
57 system: Our first 100 cases. *Apendicectomía laparoscópica por sistema «glove port»:*
58
59
60

- 1
2
3 nuestros primeros 100 casos. 2017;69(6):467-71.
4
5
6
7 155. Patrice Lemieux., Pascal Rheaume., Isabelle Levesque., Emmanuel Bujold., Gaetan
8 Brochu. Laparoscopic appendectomy in pregnant patients: a review of 45 cases. Surg
9 Endosc.2009; 23:1701.
10
11
12
13
14 156. Patel S.C., G.F., Jumba, S., Akmal. Laparoscopic appendicectomy at the Aga Khan
15 Hospital, Nairobi. East Afr Med J. 2003;80(9):447-51.
16
17
18
19
20 157. Percy C., K., Almahmoud, T., Jackson, C., Hartline, A., Cahill, L., Spence, et al. Risky
21 business? Investigating outcomes of patients undergoing urgent laparoscopic
22 appendectomy on antithrombotic therapy. Am J Surg. 2017;214(6):1012-5.
23
24
25
26
27 158. Pishori T., A.R., Siddiqui, M., Ahmed. Surgical wound infection surveillance in general
28 surgery procedures at a teaching hospital in Pakistan. Am J Infect Control. 2003;31(5):296-
29 301.
30
31
32
33
34
35 159. Putnam L.R., T.G., Ostovar-Kermani, A., Le Blanc, K.T., Anderson, G., Holzmann-
36 Pazgal, K.P., Lally, et al. Surgical site infection reporting: more than meets the agar. J
37 Pediatr Surg. 2017;52(1):156-60.
38
39
40
41
42
43 160. Al-Qahtani S.M., H.M., Al-Amoudi, S., Al-Jehani, A.S., Ashour, M.R., Abd-Hammad,
44 O.R., Tawfik, et al. Post-appendectomy surgical site infection rate after using an
45 antimicrobial film incise drape: A prospective study. Surg Infect. 2015;16(2):155-8.
46
47
48
49
50 161. Quezada F., N., Quezada, R., Mejia, A., Brañes, O., Padilla, N., Jarufe, et al.
51 Laparoscopic versus open approach in the management of appendicitis complicated
52 exclusively with peritonitis: A single center experience. Int J Surg. 2015;13:80-3.
53
54
55
56
57
58 162. Raakow J., H.-G., Liesaus, P., Neuhaus, R., Raakow. Single-incision versus multiport
59 laparoscopic appendectomy: a case-matched comparative analysis. Surg Endosc Interv
60

- 1
2
3 Tech. 2015;29(6):1530-6.
4
5
6
7 163. Ríos J., C., Murillo, G., Carrasco, C., Humet. Increase in costs attributable to surgical
8 infection after appendicectomy and colectomy. Incremento de costes atribuible a la
9 infección quirúrgica de la apendicectomía y colectomía. 2003;17(3):218-25.
10
11
12
13
14 164. Rafiq M.S., M.M., Khan, A., Khan, H., Jan. Evaluation of postoperative antibiotics after
15 non-perforated appendectomy. J Pak Med Assoc. 2015;65(8):815-7.
16
17
18
19
20
21
22 165. Reinisch A., J., Heil, G., Woeste, W., Bechstein, J., Liese. The meteorological influence
23 on seasonal alterations in the course of acute appendicitis. J Surg Res. 2017;217:137-43.
24
25
26
27
28 166. Romano A., P., Parikh, P., Byers, N., Namias. Simple acute appendicitis versus non-
29 perforated gangrenous appendicitis: Is there a difference in the rate of post-operative
30 infectious complications? Surg Infect. 2014;15(5):517-20.
31
32
33
34
35
36 167. Romel Hilaire, Fernández, Zenén Rodríguez, García, Lázaro Ibrahim Romero, Sánchez,
37 Luis Pablo Rodríguez. Apendicectomía videolaparoscópica frente a apendicectomía
38 convencional X1 Laparoscopic versus conventional appendectomy. Revista Cubana de
39 Cirugía. 2014;53(1):30-40.
40
41
42
43
44
45 168. Romy S., M.-C., Eisenring, V., Bettschart, C., Petignat, P., Francioli, N., Troillet.
46 Laparoscope use and surgical site infections in digestive surgery. Ann Surg.
47 2008;247(4):627-32.
48
49
50
51
52
53 169. Rooh-ul-Muqim, M., Khan, M., Zarin. Experience of laparoscopic appendectomies
54 versus open appendectomies. Pak J Med Sci. 2010;26(2):324-8.
55
56
57
58
59 170. Van Rossem C.C., M.H.F., Schreinemacher, K., Treskes, R.M., Van Hogeand,
60 A.A.W., Van Geloven. Duration of antibiotic treatment after appendicectomy for acute

- 1
2
3 complicated appendicitis. *Br J Surg*. 2014;101(6):715-9.
4
5
6
7
8 171. Van Rossem C.C., M.D., Bolmers, M.H., Schreinemacher, A.A., van Geloven, W.A.,
9 Bemelman. Prospective nationwide outcome audit of surgery for suspected acute
10 appendicitis. *Br J Surg*. 2016;103(1):144-51.
11
12
13
14 172. Viet Hung N., T., Anh Thu, V.D., Rosenthal, D., Tat Thanh, N., Quoc Anh, N., Le Bao
15 Tien, et al. Surgical site infection rates in seven cities in Vietnam: Findings of the
16 international nosocomial infection control consortium. *Surg Infect*. 2016;17(2):243-9.
17
18
19
20
21 173. Rotermann M. Infection after cholecystectomy, hysterectomy or appendectomy. *Health*
22 *Rep*. 2004;15(4):11-23.
23
24
25
26 174. Saar S., P., Talving, J., Laos, T., Põdramägi, M., Sokirjanski, T., Lustenberger, et al.
27 Delay Between Onset of Symptoms and Surgery in Acute Appendicitis Increases
28 Perioperative Morbidity: A Prospective Study. *World J Surg*. 2016;40(6):1308-14.
29
30
31
32
33 175. Saber A.A., M.H., Elgamal, T.H., El-Ghazaly, A.V., Dewoolkar, A., Akl. Simple
34 technique for single incision transumbilical laparoscopic appendectomy. *Int J Surg*.
35 2010;8(2):128-30.
36
37
38
39
40 176. Sadraei-Moosavi S.-M., N., Nikhbakhsh, A.-A., Darzi. Postoperative antibiotic therapy
41 after appendectomy in patients with non-perforated appendicitis. *Caspian J Int Med*.
42 2017;8(2):104-7.
43
44
45
46
47 177. Saha N., D.K., Saha, M.A., Rahman, M.K., Islam, M.A., Aziz. Comparison of post
48 operative morbidity between laparoscopic and open appendectomy in children.
49 *Mymensingh Med J*. 2010;19(3):348-52.
50
51
52
53
54 178. Sahn M., R., Kube, S., Schmidt, C., Ritter, M., Pross, H., Lippert. Current analysis of
55 endoloops in appendiceal stump closure. *Surg Endosc Interv Tech*. 2011;25(1):124-9.
56
57
58
59 179. Sahn M., M., Pross, R., Otto, A., Koch, I., Gastinger, H., Lippert. Clinical health service
60

- 1
2
3 research on the surgical therapy of acute appendicitis: Comparison of outcomes based on 3
4 German multicenter quality assurance studies over 21 years. *Ann Surg.* 2015;262(2):338-
5 46.
6
7
8
9
- 10 180. Salö M., E., Järbur, M., Hambræus, B., Ohlsson, P., Stenström, E., Arnbjörnsson. Two-
11 trocar appendectomy in children - description of technique and comparison with
12 conventional laparoscopic appendectomy. *BMC Surg.* 2016;16(1):52.
13
14
15
16
- 17 181. Sánchez-Santana T., J.A., del-Moral-Luque, P., Gil-Yonte, L., Bañuelos-Andrío, M.,
18 Durán-Poveda, G., Rodríguez-Caravaca. Effect of compliance with an antibiotic
19 prophylaxis protocol in surgical site infections in appendectomies. Prospective cohort
20 study. Efecto de la adecuación a protocolo de la profilaxis antibiótica en la incidencia de
21 infección quirúrgica en apendicectomías Estudio de cohortes prospectivo. 2017;85(3):208-
22 13.
23
24
25
26
27
28
- 29 182. Sauvain M.-O., K., Slankamenac, M.K., Muller, S., Wildi, U., Metzger, W., Schmid, et
30 al. Delaying surgery to perform CT scans for suspected appendicitis decreases the rate of
31 negative appendectomies without increasing the rate of perforation nor postoperative
32 complications. *Langenbeck's Arch Surg.* 2016;401(5):643-9.
33
34
35
36
37
38
- 39 183. Scarborough J.E., K.M., Bennett, T.N., Pappas. Racial disparities in outcomes after
40 appendectomy for acute appendicitis. *Am J Surg.* 2012;204(1):11-7.
41
42
43
44
- 45 184. Seifarth F.G., N., Kundu, A.D., Guerron, M.M., Garland, M.W., Gaffley, S., Worley, et
46 al. Umbilical Negative Pressure Dressing for Transumbilical Appendectomy in Children.
47 *JLS.* 2016;20(4).
48
49
50
- 51 185. Federico G. Seifarth, Kundu, Neilendu, Guerron, Alfredo D., Garland, Mary M.,
52 Gaffley, Michaela W. G., Worley, Sarah, et al. Umbilical Negative Pressure Dressing for
53 Transumbilical Appendectomy in Children. *JLS-JOURNAL OF THE SOCIETY OF*
54 *LAPAROENDOSCOPIC SURGEONS.* 2016;20(4).
55
56
57
58
59
- 60 186. Senekjian L., R., Nirula. Tailoring the operative approach for appendicitis to the patient:

- 1
2
3 A prediction model from national surgical quality improvement program data. *J Am Coll*
4 *Surg.* 2013;216(1):34-40.
5
6
7
8
9 187. Sesia S.B., M., Frech, F.-M., Häcker, J., Mayr. Laparoscopic "single-port"
10 appendectomy in children. *Laparoskopische "single port"-appendektomie im Kindesalter.*
11 2011;136(1):50-5.
12
13
14
15 188. Shaikh A.R., S., Khatoon, M., Arif. Evaluation of re-admission after open
16 appendectomy. *Rawal Med J.* 2011;36(2):100-3.
17
18
19
20
21
22 189. Shang Q., Q., Geng, X., Zhang, C., Guo. The efficacy of combined therapy with
23 metronidazole and broad-spectrum antibiotics on postoperative outcomes for pediatric
24 patients with perforated appendicitis. *Medicine.* 2017;96(47).
25
26
27
28
29 190. Shindholimath V., K., Thinakaran, T., Rao, Y., Veerappa. Laparoscopic management
30 of appendicular mass. *J Minimal Access Surg.* 2011;7(2):136-40.
31
32
33
34 191. Shimizu T., M., Ishizuka, K., Kubota. The preoperative serum C-reactive protein level
35 is a useful predictor of surgical site infections in patients undergoing appendectomy. *Surg*
36 *Today.* 2015;45(11):1404-10.
37
38
39
40
41 192. Siam B., A., Al-Kurd, N., Simanovsky, H., Awesat, Y., Cohn, B., Helou, et al.
42 Comparison of appendectomy outcomes between senior general surgeons and general
43 surgery residents. *JAMA Surg.* 2017;152(7):679-85.
44
45
46
47
48 193. 서승원, 김신곤. Acute Appendicitis in Pregnant Patients and Non-Pregnant Patients:
49
50
51 Recent Clinical Experience of the Tertiary Hospital Z1 임신부와 가임기 여성의 급성
52
53
54
55
56
57
58
59
60
충수염의 비교 고찰: 최근 3차 병원의 임상적 경험. *Annals of Surgical Treatment and*
Research S1 대한외과학회지. 2002;62(6):486-90.

- 1
2
3
4
5
6
7 194. Alfredo Silva, M, Guido Vargas, A, Amparo Moreno, H, Pablo Becerra. Utilidad del
8 retractor elástico abdominal para disminuir el riesgo de infección de herida operatoria en
9 apendicitis aguda X1 Use of an elasticwall retractor during appendectomy to reduce wound
10 infection. *Revista chilena de cirugía*. 2008;60(6):527-33.
11
12
13
14
15 195. Singh V.K., K., Nishant, B., Kharga, A.K., Kalita, P., Bhutia, J., Jain. Randomized
16 controlled trial comparing open, conventional, and single port laparoscopic appendectomy.
17 *J Clin Diagn Res*. 2017;11(10):PC05-PC10.
18
19
20
21
22
23
24 196. Siribumrungwong B., K., Srikuea, A., Thakkinstian. Comparison of superficial surgical
25 site infection between delayed primary and primary wound closures in ruptured
26 appendicitis. *Asian J Surg*. 2014;37(3):120-4.
27
28
29
30
31 197. Sivrikoz E., E., Karamanos, E., Beale, P., Teixeira, K., Inaba, D., Demetriades. The
32 effect of diabetes on outcomes following emergency appendectomy in patients without
33 comorbidities: A propensity score-matched analysis of National Surgical Quality
34 Improvement Program database. *Am J Surg*. 2015;209(1):206-11.
35
36
37
38
39 198. Soll C., P., Wyss, H., Gelpke, D.A., Raptis, S., Breitenstein. Appendiceal stump closure
40 using polymeric clips reduces intra-abdominal abscesses. *Langenbeck's Arch Surg*.
41 2016;401(5):661-6.
42
43
44
45
46 199. Sozutek A., T., Colak, M., Dirlik, K., Ocal, O., Turkmenoglu, A., Dag. A prospective
47 randomized comparison of single-port laparoscopic procedure with open and standard 3-
48 port laparoscopic procedures in the treatment of acute appendicitis. *Surg Laparoscopy*
49 *Endosc Percutaneous Tech*. 2013;23(1):74-8.
50
51
52
53
54 200. Srishewachart P., S., Narksut. Incidence of abnormal preoperative blood testing and
55 postoperative complication in appendectomy patients in Siriraj Hospital. *J Med Assoc*
56 *Thailand*. 2016;99(5):517-24.
57
58
59
60

- 1
2
3 201. Staszewicz W., M.-C., Eisenring, V., Bettschart, S., Harbarth, N., Troillet. Thirteen
4 years of surgical site infection surveillance in Swiss hospitals. *J Hosp Infect.* 2014;88(1):40-
5 7.
6
7
8
9
10 202. Suttie S.A., S., Seth, C.P., Driver, A.A., Mahomed. Outcome after intra- and extra-
11 corporeal laparoscopic appendectomy techniques. *Surg Endosc.* 2004;18(7):1123-5.
12
13
14
15 203. Svensson J.F., B., Patkova, M., Almström, S., Eaton, T., Wester. Outcome after
16 introduction of laparoscopic appendectomy in children: A cohort study. *J Pediatr Surg.*
17 2016;51(3):449-53.
18
19
20
21
22 204. Taguchi Y., S., Komatsu, E., Sakamoto, S., Norimizu, Y., Shingu, H., Hasegawa.
23 Laparoscopic versus open surgery for complicated appendicitis in adults: a randomized
24 controlled trial. *Surg Endosc Interv Tech.* 2016;30(5):1705-12.
25
26
27
28
29 205. Tanaka S., D., Kubota, S.H., Lee, K., Oba, M., Matsuyama. Effectiveness of
30 laparoscopic approach for acute appendicitis. *Osaka City Med J.* 2007;53(1):1-8.
31
32
33
34 206. Tijerina J., R., Velasco-Rodríguez, C., Vásquez, V., Melnikov, S., Rodriguez.
35 Effectiveness of a systemic antibiotic followed by topical ionized solution as surgical site
36 infection prophylaxis. *J Int Med Res.* 2010;38(4):1287-93.
37
38
39
40
41 207. The SCARLESS Study Group. Single port/incision laparoscopic surgery compared with
42 standard three-port laparoscopic surgery for appendectomy: A randomized controlled
43 trial. *Surg Endosc Interv Tech.* 2015;29(1):77-85.
44
45
46
47
48 208. Juan Pablo Toro, Barrera, Óscar Javier, Morales, Carlos Hernando. Superioridad clínica
49 de la apendicectomía laparoscópica sobre la técnica abierta: ¿adopción lenta de un nuevo
50 estándar de tratamiento? X1 Clinical superiority of laparoscopic appendectomy over the
51 open technique: sluggish adoption of a new standard of treatment? *Revista Colombiana de*
52 *Cirugía.* 2017;32(1):32-9.
53
54
55
56
57
58 209. Towfigh S., T., Clarke, W., Yacoub, A.H., Pooli, R.J., Mason, N., Katkhouda, et al.
59 Significant reduction of wound infections with daily probing of contaminated wounds : A
60

- 1
2
3 prospective randomized clinical trial. *Arch Surg.* 2011;146(4):448-52.
4
5
6
7 210. Troillet N., E., Aghayev, M.-C., Eisenring, A.F., Widmer. First Results of the Swiss
8 National Surgical Site Infection Surveillance Program: Who Seeks Shall Find. *Infect*
9 *Control Hosp Epidemiol.* 2017;38(6):697-704.
10
11
12
13 211. Tsioplis C., C., Brockschmidt, S., Sander, D., Henne-Bruns, M., Kornmann. Factors
14 influencing the course of acute appendicitis in adults and children. *Langenbeck's Arch Surg.*
15 2013;398(6):857-67.
16
17
18
19
20 212. Vahdad M.R., M., Nissen, A., Semaan, T., Klein, E., Palade, T., Boemers, et al.
21 Experiences with LESS-appendectomy in Children. *Arch Iran Med.* 2016;19(1):57-63.
22
23
24
25 213. Van Rossem C.C., M.H.F., Schreinemacher, A.A.W., Van Geloven, W.A., Bemelman,
26 G.J.D., Van Acker, B., Akkermans, et al. Antibiotic duration after laparoscopic
27 appendectomy for acute complicated appendicitis. *JAMA Surg.* 2016;151(4):323-9.
28
29
30
31
32 214. Van Rossem C.C., M.H.F., Schreinemacher, K., Treskes, R.M., Van Hogeand,
33 A.A.W., Van Geloven. Duration of antibiotic treatment after appendectomy for acute
34 complicated appendicitis. *Br J Surg.* 2014;101(6):715-9.
35
36
37
38
39 215. Wang-Chan A., F.H., Hetzer, C., Gingert, C., Gingert, E., Angst, E., Angst, et al.
40 Clinical relevance and effect of surgical wound classification in appendicitis: Retrospective
41 evaluation of wound classification discrepancies between surgeons, Swissnosot-trained
42 infection control nurse, and histology as well as surgical site infection rates by wound class.
43 *J Surg Res.* 2017;215:132-9.
44
45
46
47
48
49 216. Watanabe A., S., Kohnoe, H., Sonoda, K., Shirabe, K., Fukuzawa, S., Maekawa, et al.
50 Effect of intra-abdominal absorbable sutures on surgical site infection. *Surg Today.*
51 2012;42(1):52-9.
52
53
54
55
56 217. Willis Z.I., E.M., Duggan, B.T., Bucher, J.B., Pietsch, M., Milovancev, W., Wharton,
57 et al. Effect of a clinical practice guideline for pediatric complicated appendicitis. *JAMA*
58 *Surg.* 2016;151(5).
59
60

- 1
2
3
4
5 218. Ramírez-Wong F.M., T., Atencio-Espinoza, V.D., Rosenthal, E., Ramirez, S.L., Torres-
6 Zegarra, Z.R., Díaz Tavera, et al. Surgical Site Infections Rates in More Than 13,000
7 Surgical Procedures in Three Cities in Peru: Findings of the International Nosocomial
8 Infection Control Consortium. *Surg Infect.* 2015;16(5):572-6.
9
10
11
12
13 219. Wu J.-M., K.-H., Chen, H.-F., Lin, L.-M., Tseng, S.-H., Tseng, S.-H., Huang.
14 Laparoscopic appendectomy in pregnancy. *J Laparoendosc Adv Surg Techn Part A.*
15 2005;15(5):447-50.
16
17
18
19
20 220. Wu H.-S., H.-W., Lai, S.-J., Kuo, Y.-T., Lee, D.-R., Chen, C.-W., Chi, et al. Competitive
21 edge of laparoscopic appendectomy versus open appendectomy: A subgroup comparison
22 analysis. *J Laparoendosc Adv Surg Techn.* 2011;21(3):197-202.
23
24
25
26
27 221. Wu K., L., Yang, A., Wu, J., Wang, S., Xu, H., Zhao, et al. Single-site laparoscopic
28 appendectomy in children using conventional instruments: a prospective, randomized,
29 control trial. *Pediatr Surg Int.* 2014;31(2):167-71.
30
31
32
33
34 222. Wu T.-C., Q., Lu, Z.-Y., Huang, X.-H., Liang. Efficacy of emergency laparoscopic
35 appendectomy in treating complicated appendicitis for elderly patients. *Saudi Med J.*
36 2017;38(11):1108-12.
37
38
39
40
41 223. Yaghoubian A., C., de Virgilio, V., Chiu, S.L., Lee. "July effect" and appendicitis. *J*
42 *Surg Educ.* 2010;67(3):157-60.
43
44
45
46 224. Yagnik V., J., Rathod, A., Phatak. A retrospective study of two-port appendectomy and
47 its comparison with open appendectomy and three-port appendectomy. *Saudi J*
48 *Gastroenterol.* 2010;16(4):268-71.
49
50
51
52
53 225. Yousef Y., F., Yousef, M., Homsy, T., Dinh, K., Pandya, H., Stagg, et al.
54 Standardization of care for pediatric perforated appendicitis improves outcomes. *J Pediatr*
55 *Surg.* 2017;52(12):1916-20.
56
57
58
59
60 226. Zhang Z., Y., Wang, R., Liu, L., Zhao, H., Liu, J., Zhang, et al. Suprapubic single-

1
2
3 incision versus conventional laparoscopic appendectomy. J Surg Res. 2016;200(1):131-8.
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

BMJ Open

Global incidence of surgical-site infection after appendectomy: a systematic review and meta-analysis

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2019-034266.R2
Article Type:	Original research
Date Submitted by the Author:	06-Jan-2020
Complete List of Authors:	Danwang, Celestin; Universite Libre de Bruxelles, School of Public Health; University of Yaoundé I, Department of Surgery and Specialties, Faculty of Medicine and Biomedical Sciences Bigna, Jean Joel; Centre Pasteur of Cameroon, Department of Epidemiology and Public Health Tochie, Joel Noutakdie; University of Yaoundé I, Department of Anesthesiology and Critical Care Medicine, Faculty of Medicine and Biomedical Sciences Mbonda, Aimé; Djohong District Hospital, Department of General Medicine Mbanga, Clarence; Mankon Sub-divisional Hospital Nzalie, Rolf ; Ngong District Hospital, North Region, Cameroon Guifo, Marc Leroy ; University of Yaoundé I, Department of Surgery and Specialties, Faculty of Medicine and Biomedical Sciences Essomba, Arthur ; University of Yaoundé I, Department of Surgery and Specialties, Faculty of Medicine and Biomedical Sciences
Primary Subject Heading:	Surgery
Secondary Subject Heading:	Global health, Infectious diseases, Public health
Keywords:	EPIDEMIOLOGY, Health policy < HEALTH SERVICES ADMINISTRATION & MANAGEMENT, Quality in health care < HEALTH SERVICES ADMINISTRATION & MANAGEMENT, Epidemiology < INFECTIOUS DISEASES, Public health < INFECTIOUS DISEASES

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

1 Global incidence of surgical-site infection after appendectomy: a systematic 2 review and meta-analysis

3 Celestin Danwang^{1,2}, Jean Joel Bigna^{3*}, Joel Noutakdie Tochie⁴, Aime Mbonda⁵, Clarence
4 Mvalo Mbanga⁶, Rolf Nyah Tuku Nzalieu⁷, Marc Leroy Guifo¹, Arthur Essomba¹

- 5
- 6 1. Department of Surgery and Specialties, Faculty of Medicine and Biomedical Sciences,
7 University of Yaoundé I, Yaoundé, Cameroon.
- 8 2. School of Public Health, Université Libre de Bruxelles, Brussels, Belgium.
- 9 3. Department of Epidemiology and Public Health, Centre Pasteur of Cameroon, Yaoundé,
10 Cameroon.
- 11 4. Department of Anesthesiology and Critical Care Medicine, Faculty of Medicine and Biomedical
12 Sciences, University of Yaoundé I, Yaoundé, Cameroon.
- 13 5. Department of General Medicine, Djohong District Hospital, Djohong, Cameroon.
- 14 6. Department of General Medicine, Mankon Sub-Divisional Hospital, Bamenda,
15 Cameroon.
- 16 7. Department of General Medicine, Ngong District Hospital, Ngong, Cameroon.

17
18 *Corresponding author: Dr. Jean Joel Bigna, MD, MPH

19 Department of Epidemiology and Public Health, Centre Pasteur of Cameroon, Yaoundé,
20 Cameroon. E-Mail: bignarimjj@yahoo.fr

21
22 E-mail addresses:

23 CD: danram07@yahoo.fr; JJB: bignarimjj@yahoo.fr; JNT: joeltochie@gmail.com; AM:
24 aimembonda@yahoo.fr; CMM: mbangaclarence@gmail.com; RNN: nzalierolf@yahoo.com;
25 GML: marcleroyguifo@gmail.com; AE: artgessomba@gmail.com

26
27 **Word count:** 2,774.

28 **Abstract**

29 **Background:** Although surgical-site infection (SSI) is one of the most studied healthcare-
30 associated infections, the global burden of SSI after appendectomy remains unknown.

31 **Objectives:** We estimated the incidence of SSI after appendectomy at global and regional
32 levels.

33 **Design:** Systematic review and meta-analysis.

34 **Participants:** Patients with appendectomy.

35 **Data sources:** EMBASE, PubMed, and Web of Science were searched to identify observational
36 studies and clinical trials, published between January 1, 2000 and December 30, 2018 and
37 reporting on the incidence of the SSI after an appendectomy; with no language restriction. A
38 random-effect models meta-analysis served to obtain the pooled incidence of SSI after 100
39 surgical procedures in patients with appendectomy.

40 **Results:** In total, 226 studies (729,434 participants from 49 countries) were included in the
41 meta-analysis. Concerning the methodological quality, 59 (26.1%) studies had a low risk, 147
42 (65.0%) a moderate risk, and 20 (8.8%) a high risk of bias. We found an overall incidence of
43 SSIs of 7.0 per 100 surgical procedures (95% prediction interval: 1.0-17.6) for appendectomy
44 varying from 0 to 37.4 per 100 surgical procedures. Subgroup analysis for identifying sources
45 of heterogeneity showed that the incidence varied from 5.8 in Europe to 12.6 per 100 surgical
46 procedures in Africa, $p < 0.0001$. The incidence of SSI after appendectomy increased when the
47 level of income decreased; from 6.2 in high-income countries to 11.1 per 100 surgical
48 procedures in low-income countries ($p = 0.015$). Open appendectomy (11.0 per 100 surgical
49 procedures) was found to have a higher incidence of SSI compared to laparoscopy (4.6 per 100
50 surgical procedures), $p = 0.0002$.

1
2
3 51 **Conclusion:** This study suggests a high burden of SSIs after appendectomy in some regions
4
5 52 (especially Africa) and in low-income countries. Strategies are needed to implement and
6
7 53 vulgarize WHO guidelines to decrease the burden of SSI after appendectomy in these regions.
8
9

10 54 **Registration:** PROSPERO, CRD42017075257.
11
12
13 55
14
15 56
16

17 57 **Keywords:**

18
19 58 Surgical wound infection; Global Health; Hospital infections; Cross infection; Healthcare
20
21
22 59 associated infection
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

60 **Strengths and limitations of this study**

- 61 • This meta-analysis is the first to summarize the global incidence of SSIs after
62 appendectomy.
- 63 • We investigated WHO regions, level of income, and surgical procedure as sources of
64 heterogeneity.
- 65 • We were not able to investigate all sources of heterogeneity because of missing information
66 in the original studies.
- 67 • There were few studies from low income countries and from Africa.

Introduction

Defined as an acute inflammation of the vermiform appendix,¹ evidence abounds that acute appendicitis is the most common abdominal surgical emergency,² with an incidence of almost 100 per 100,000 person-years reported in Australia, Europe and North America.^{3 4} Evidence suggests appendectomy, a surgical remove of the vermiform appendix as first-line treatment for acute appendicitis, although antibiotic therapy may be efficacious for a selected group of patients with uncomplicated acute appendicitis.⁵⁻⁷ Appendectomy is a relatively safe surgical intervention with a case fatality rate of 2.1 - 2.4 per 1000 patients as reported in studies conducted in Europe.^{8 9}

Innovations in appendectomy, especially with the advent of minimally invasive or laparoscopic surgery in 1983,¹⁰ which has replaced the traditional open appendectomy in most of high-income countries, has led to a drastic reduction in the morbidity and mortality related to appendectomy.¹¹⁻¹³ Laparoscopic appendectomy is now recognized as the gold standard surgical approach for uncomplicated acute appendicitis owing to its merits over open surgery; due to less postoperative pain, reduced postoperative ileus, shorter hospital stay, rapid postoperative recovery, and better aesthetic scars.¹⁴⁻¹⁹

However, regardless of the surgical technique (laparoscopic or open surgery), appendectomy remains a sceptical surgical intervention associated with a substantial risk of surgical-site infections (SSIs). SSIs after appendectomy are postoperative nosocomial infections affecting the incision site, deep tissues, organs at the operative site within 30 days after the surgical procedure.²⁰⁻²² SSI following appendectomy is a serious post-operative medical concern that increases the financial burden for both healthcare systems and patient, and also have a negative impact on the patients' health related quality of life.²³⁻²⁸

SSI is both the most frequently studied and the leading healthcare-associated infections reported hospital-wide in low- and middle-income countries.²⁹ A recently published prospective

1
2
3 international multicentre cohort study suggested a high burden of SSIs after any gastrointestinal
4 surgery in low-income countries compared to high-income countries.³⁰ Actually, there is no
5 global systematic review with meta-analysis reporting the burden of SSI after appendectomy or
6
7 comparing the burden between regions and between country level of income. It would be
8 interesting to have such accurately estimated data to construct efficient strategies to curb
9
10 globally the burden of SSIs after appendectomy. To fill this gap, the current systematic review
11 and meta-analysis aimed at summarizing contemporary data on the occurrence of SSIs after
12
13 appendectomy.
14
15
16
17
18
19
20
21
22
23

24 **Methods**

25 **Design**

26
27
28
29 This systematic review and meta-analysis was registered in the International Prospective
30 Register of Systematic Reviews (PROSPERO) under the registration number
31 CRD42017075257. The protocol has been published in a peer-review journal.³¹ This review is
32 reported according to MOOSE and PRISMA guidelines.^{32 33}
33
34
35
36
37
38

39 **Eligibility criteria**

40
41 We considered observational studies (cross-sectional, case-control, and cohort) and clinical
42 trials of patients with appendectomy. Outcome of interest was incidence of SSI of enough data
43 (number of cases of SSI and sample size) to compute this estimate. We excluded letters,
44 reviews, commentaries and editorials, and studies lacking key data and/or explicit method
45 description as well as studies in which relevant data on SSIs after appendectomy was impossible
46 to extract even after contacting the corresponding author.
47
48
49
50
51
52
53
54

55 **Search strategy**

1
2
3 We searched EMBASE, PubMed, and Web of Science (Web of Science Core Collection,
4 Current Contents Connect, KCI-Korean Journal Database, SciELO Citation Index, Russian
5 Science Citation Index) to identify observational studies, published between January 1, 2000
6 and December 30, 2018. No language restriction was applied. The initial search strategy was
7 designed for EMBASE and was adapted for the use in other databases. The search strategy as
8 illustrated in the Supplementary Table 1 and in the study protocol,³¹ was based on the
9 combination of relevant text words and medical subject headings related to SSIs. Moreover, the
10 references of all relevant articles found were scrutinized for potential additional data sources.
11 When a full text was not available, it was requested via the corresponding author by email. For
12 duplicates or studies published in more than one report, the one reporting the largest sample
13 size was considered.

24 **Study selection**

25 Two reviewers (CD and AM) independently screened the titles and abstract of articles for
26 eligibility. Full texts of potentially eligible articles were retrieved and screened for final
27 inclusion. Disagreements between the two reviewers were solved by discussion and when a
28 consensus was not reached, a third reviewer (JNT) resolved discrepancies. Studies in other
29 languages than French, English, and Spanish were translated using Google Translate.

30 **Data extraction and management**

31 A standardized and pretested data extraction form was used by five reviewers (CD, JNT, AM,
32 RNZ, CMM) to independently extract data from individual studies. A sixth reviewer (JJB)
33 independently extracted data for accuracy. The last name of the first author, year of publication,
34 country, study design, age groups, sample size, mean or median age, proportion of males,
35 specific conditions of the study population, the surgical method (open surgery or laparoscopy),
36 and incidence of SSIs after appendectomy in the study population (or enough data to compute
37 this estimate) were extracted.

1
2
3 To assess the methodological quality of each study, two reviewers (CD and CMM) used an
4 adapted version of the tool of bias assessment for prevalence studies developed by Hoy and
5
6 colleagues.³⁴
7
8
9

10 **Data synthesis and analysis**

11
12
13 A meta-analysis was used to summarize data concerning incidence of SSIs, by pooling together
14 data of studies reporting the incidence of SSIs. Study-specific estimates were then pooled
15 through a Dersimonian and Laird random-effects meta-analysis model to obtain an overall
16 summary estimate of the incidence across studies, after stabilizing the variance of individual
17 studies using the Freeman-Tukey double arc-sine transformation.³⁵ Incidence was expressed by
18 100 surgical procedures with their 95% confidence interval and 95% prediction interval.
19
20 Heterogeneity was evaluated by the χ^2 test on Q statistic which is quantified by I^2 values,³⁶
21 assuming that I^2 values of 25%, 50% and 75% represent low, medium and high heterogeneity
22 respectively.³⁷ Where substantial heterogeneity ($I^2 > 50\%$) was detected, a subgroup analysis
23 was performed to detect its possible sources using the following grouping variables: type of
24 surgery (laparoscopy or open), World Health Organization regions, and country level of
25 income. A p value < 0.05 was indicative of significant difference. The meta-regression analysis
26 was performed to estimate the explained heterogeneity of each covariate included in the
27 subgroup analysis. Inter-rater agreement for study inclusion was assessed using Cohen's κ
28 coefficient.³⁸ Funnel plots analysis and Egger's test ($p < 0.10$) were performed to detect the
29 presence of publication bias.³⁹ Since we believe that the incidence estimates of interest would likely
30 be published even if substantially different from previously reported estimates, we have not reported
31 adjusted incidence estimate in the case of publication bias. Data were analysed using the '*meta*' package
32 in R, version 3.6.1.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55

56 **Patient and public involvement**

1
2
3 Patients or the public were not involved in the design, or conduct, or reporting, or dissemination
4
5 of our research.
6
7
8
9

10 **Results**

11 **Study selection and characteristics**

12
13 Overall, 619 records were initially identified. After removal of duplicates, screening of study
14
15 titles, abstracts, and full texts; 226 studies including 729,434 patients were finally retained for
16
17 meta-analysis (Supplementary Figure 1). The full list of included studies is in the Appendix.
18
19 Concerning the methodological quality, 59 (26.1%) studies had a low risk, 147 (65.0%) a
20
21 moderate risk and 20 (8.8%) a high risk of bias. Supplementary Table 2 presents characteristics
22
23 of included studies. Among the included studies, 154 were done in high-income, 36 upper-
24
25 middle, 27 lower-middle, and nine in low-income countries. Overall, most of studies were from
26
27 Europe (n = 68) and Americas (n = 67). SSIs were defined according to Center of Disease
28
29 Control and Prevention criteria in 50 studies while 25 studies used other criteria. The definition
30
31 of SSIs was not clearly given in 151 studies. Individuals characteristics of included studies are
32
33 in the Supplementary Table 3.
34
35
36
37
38
39

40 **Overall prevalence**

41
42 The overall incidence of SSI after appendectomy was 7.0 per 100 surgical procedures (95%
43
44 prediction interval: 1.0-17.6) varying from 0% to 37.4% with substantial heterogeneity and
45
46 publication bias (Supplementary Figure 2). The sensitive analysis including only studies with
47
48 low risk of bias yielded a very close incidence to crude analysis (Table 1).
49
50
51

52 **Sources of heterogeneity**

1
2
3 According to country level of income (Figure 1), the incidence of SSI after appendectomy
4 increased when the level of income decreased; from 6.2 in high income countries to 11.1 per
5 100 surgical procedures in low income countries ($p = 0.015$) (Table 1).
6
7

8
9
10 The incidence varied widely across WHO regions (Figure 2). The incidence varied from 5.8 in
11 Europe to 12.6 per 100 surgical procedures in Africa, $p < 0.0001$ (Table 1). Two regions
12 (Europe and Americas) had an incidence < 6 per 100 surgical procedures, three an incidence
13 between 6-10 per 100 procedures (South-East Asia, Eastern Mediterranean, and Western
14 Pacific), and one an incidence > 10 per 100 procedures (Africa) (Table 1). The incidence also
15 varied widely in different regions. The incidence varied from 0.2 to 32.0 in Africa, from 1.9 to
16 37.4 in Western Pacific, from 1.3 to 33.8 in Eastern Mediterranean, from 1.2 to 25.8 in South-
17 East Asia, from 0.1 to 37.4 in Americas, and from 0 to 20.0 per 100 surgical procedures in
18 Europe (Figure 2).
19
20
21
22
23
24
25
26
27
28
29

30 Open appendectomy with an incidence of 11.0 (95% prediction interval: 0.0-39.3) per 100
31 surgical procedures was found to have a higher incidence of SSI compared to laparoscopic
32 appendectomy with an incidence of 4.6 (95% prediction interval: 0.0-14.3) per 100 surgical
33 procedures, $p = 0.0002$ (Figure 3).
34
35
36
37
38
39

40 Heterogeneity of the overall incidence of SSI after appendectomy was explained by WHO
41 regions (17.1%), country level of income (11.1%), and type of surgical procedure (4.9%). We
42 conducted a post hoc analysis; then in a meta-regression analysis of 119 studies reporting the
43 information of the use of antibiotics, there was no association between the variation of SSI
44 incidence and proportion of patients with the use of antibiotics (coefficient: 0.0010 [95%CI: -
45 0.0004; 0.0023]; $p = 0.170$). however, most (79.5%) of these studies reported using antibiotics
46 for all patients.
47
48
49
50
51
52
53
54
55
56
57

58 Discussion

59
60

1
2
3 This first systematic review and meta-analysis of data of 729,434 surgical procedures in 226
4 studies from 49 countries found an overall incidence of SSIs of 7.0 per 100 surgical procedures
5 for appendectomy varying from 0 to 37.4 per 100 surgical procedures with substantial
6 heterogeneity according to WHO regions, country level of income, and type of surgical
7 procedure. The incidence increased with decreasing country level of income and was higher
8 when using open surgery compared to laparoscopy. The incidence significantly varied by WHO
9 regions with Africa having the highest burden followed by Western Pacific, Eastern-
10 Mediterranean, and South-East Asia. We found no association between SSI incidence and
11 proportion of using antibiotics.

12
13
14
15
16
17
18
19
20
21
22
23
24 Health care-associated infections are acquired by patients when receiving care and are the most
25 frequent adverse event affecting patient safety worldwide. This includes SSIs after
26 appendectomy.⁴⁰ As reported in a previous systematic review and meta-analysis, SSIs were the
27 leading infection in hospitals in developed countries.²⁹ The high incidence we found in this
28 study suggests that SSIs after appendectomy remains a global public concern. WHO reported
29 that of every 100 hospitalized patients at any given time, seven in developed and 15 in
30 developing countries will acquire at least one health care-associated infection.⁴⁰ SSIs are mainly
31 caused by micro-organisms resistant to commonly-used antimicrobials, which can be
32 multidrug-resistant. Indeed, more than 50% of SSIs can be antibiotic-resistant.⁴¹ The leading
33 micro-organisms identified in SSIs are *Staphylococcus aureus*, coagulase-negative
34 staphylococci, and *Escherichia coli* as reported by National Healthcare Safety Network.⁴¹ It is
35 important to worry since *Staphylococcus aureus* and *Escherichia coli* are the micro-organisms
36 with highest proportion of antibiotic resistance, respectively resistant to oxacillin/methicillin in
37 43% of cases and to fluoroquinolones in 25% of cases.⁴¹ A recent international prospective
38 cohort study shown that 21.6% of patients with SSI after any gastrointestinal surgery had an
39 infection that was resistant to the prophylactic antibiotic used.³⁰ There are many factors that can
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 favour SSI including patient-related and procedural-related variable.⁴² These factors can be
4
5 classified in two categories; non-modifiable like age and sex and modifiable including
6
7 nutritional status, tobacco use, correct use of antibiotics, obesity, diabetes, prolonged surgery
8
9 duration, pre-surgery hospital stay of at least two days, lower volume of hospital and surgeons,
10
11 and the intraoperative techniques.⁴⁰ Strategies to curb the burden of SSIs should therefore focus
12
13 on addressing these identified factors. However, we were not able to find an association
14
15 between SSI with the use antibiotics, may be due to the low variability in the proportion of
16
17 antibiotics in the original studies.
18
19

20
21 In our present study looking at specifically SSI after appendectomy, we also found that SSI was
22
23 higher in low income countries. Interestingly, there was a trend with increasing incidence when
24
25 the country income decreased. The WHO Africa region essentially constituted with sub-
26
27 Saharan Africa was the region with highest incidence in this study. The WHO estimates that
28
29 the endemic burden of health care-associated infections is two to three time significantly higher
30
31 in low- and middle-income countries than in high-income nations.⁴⁰ The highest burden found
32
33 in Africa may be associated with the fact most of countries in this continent are low income
34
35 countries compared to other regions. Indeed, factors associated with increased risk of SSI after
36
37 appendectomy may be higher in low-income settings. The burden of diabetes, obesity, and
38
39 undernutrition are increasing in low-income countries.^{43 44} There is also inadequate use of
40
41 antimicrobial in low- and middle-income countries and micro-organisms are more resistant to
42
43 prophylactic antibiotics used to prevent SSI in low-income countries compared to high-income
44
45 countries.^{30 45 46} Lower level income is also associated with lower volume of surgeon and
46
47 hospital, factors recognised as associated increased risk of SSIs.⁴⁰ The higher incidence found
48
49 in low income countries may also be explained by the fact open surgery is the most used surgical
50
51 procedure in this setting. Indeed, we found as in other studies that open surgery is associated
52
53 with higher incidence of SSIs compared to laparoscopy.^{47 48} Laparoscopy is generally indicated
54
55
56
57
58
59
60

1
2
3 for uncomplicated appendicitis where the dissemination of micro-organism is lower compared
4
5 open surgery indicated for perforated appendicitis with peritonitis for example. Moreover, only
6
7 few low-income countries have the necessary infrastructure to carry out laparoscopy procedures
8
9 compared to high-income countries.⁴⁹⁻⁵¹

10
11
12 Our findings have important implications for healthcare providers and health policy makers.
13
14 SSIs are among the most preventable healthcare-associated infections.⁵²⁻⁵³ They still represent
15
16 a significant burden in terms of patient morbidity and mortality and additional costs for
17
18 healthcare systems.⁴⁰ The prevention of SSI has received considerable attention from surgeons,
19
20 infection control professionals, health policy makers, the media and the public since there is a
21
22 perception among the public that SSIs may reflect a poor quality of care.⁵⁴ However, special
23
24 attention is needed for low-income countries and Africa. Strategy to curb the burden of SSIs
25
26 after appendectomy as for other surgery procedures should be focused on strategies that can
27
28 help to address factors associated with increased risk of SSIs. Therefore, strategies should be a
29
30 package including how to address the factors cited above. The 26 WHO recommendations to
31
32 avoid SSIs should be vulgarized and implemented,⁴⁰ especially in low-income countries.
33
34 Strengthening the healthcare systems of low-income countries and of countries in WHO Afro
35
36 region is also a paramount by education of healthcare providers and skilling them on the use of
37
38 very less invasive surgical procedures.

39
40 This study should however be interpreted in the context of some drawbacks. Firstly, the same
41
42 definition of SSIs was not used by all the included studies. In addition, there were some
43
44 heterogeneity according to the surgical procedure and the profile of patients. This may lead to
45
46 an overestimation or underestimation of the SSIs incidence by individual studies (depending on
47
48 the study characteristics). Secondly, few studies reported on the participants' characteristics
49
50 and details on the surgical procedure since this can modify the risk for developing SSIs. We
51
52 were not therefore able to measure the impact on our outcome of interest. Thirdly, only a quarter
53
54
55
56
57
58
59
60

1
2
3 of studies had low risk of bias, however our analysis including only studies with low risk of
4 bias yielded an estimate close to the crude incidence. Fourth, the various geographic regions
5 and countries were variably represented, with some countries with only one study or even no
6 study, which could affect the generalizability of our findings.
7
8
9
10

11
12 Despite these limitations, this is the first systematic review and meta-analysis providing a global
13 estimate of the burden of SSIs after appendectomy. A protocol had been published before, and
14 we used rigorous methodological and statistical procedures to obtain and pool data.
15
16 Furthermore, subgroup analyses were conducted to investigate the various factors likely
17 affecting our estimate.
18
19
20
21
22
23
24
25

26 **Conclusion**

27
28 This systematic review and meta-analysis compiled data from more than 700,000 people with
29 appendicitis in 49 countries and pointed a high incidence of SSIs after appendectomy, at 7 per
30 100 surgical procedures. This estimate seemed higher in some WHO regions (especially Africa)
31 and in low-income countries. These data suggest that less invasive procedure is associated with
32 low incidence of SSIs after appendectomy. Strategies are needed to implement already known
33 guidelines to decrease the burden of SSI after appendectomy. However, in low-income
34 countries which have weak health systems, cost-effectiveness studies are needed to inform
35 policies regarding the best strategies for decreasing the burden of SSI after appendectomy.
36
37
38
39
40
41
42
43
44
45
46
47
48

49 **Contributors**

50
51
52 CD and JJB conceived the idea of the study and developed the protocol. JJB, CD, and JNT did
53 the literature search. CD, AM, and JNT selected the studies, CD, JNT, RNZ, AM, CMM, JJB
54 extracted the relevant information. CD, JJB, and CMM synthesized the data. CD, JNT, CMM,
55 and JJB wrote the first draft of the paper. CD, JJB, JNT, AB, RNZ, CMM, GML, and AE
56
57
58
59
60

1
2
3 critically revised successive drafts of the paper and approved the final version. GML and AE
4
5 supervised the overall work, CD and JJB are the guarantors of the review.
6

7 **Acknowledgements**

8
9
10 None.

11 **Funding**

12
13
14 This research received no specific grant from any funding agency in the public, commercial or
15
16 not-for-profit sectors.
17

18 **Competing interests**

19
20
21 We declare no competing interests.
22

23 **Patient consent**

24
25
26 Not applicable.
27

28 **Data sharing statement**

29
30
31 All data generated for this study are in the manuscript and its supporting files.
32
33

34 **Figures Legend**

35
36
37 Figure 1. Global incidence of SSI (surgical site infection) after appendectomy by level of
38
39 country income
40

41
42 Figure 2. Global incidence of SSI (surgical site infection) after appendectomy by WHO regions
43

44
45 Figure 3. Global incidence of SSI (surgical site infection) after appendectomy by type of
46
47 surgical procedures
48

49 **References**

- 50
51
52
53
54 1. Giesen LJX, van den Boom AL, van Rossem CC, et al. Retrospective Multicenter Study on
55 Risk Factors for Surgical Site Infections after Appendectomy for Acute Appendicitis.
56 *Dig Surg* 2017;34(2):103-07. doi: 10.1159/000447647 [published Online First: 09/16]
57
58 2. Navarro Fernández JA, Tárraga López PJ, Rodríguez Montes JA, et al. Validity of tests
59 performed to diagnose acute abdominal pain in patients admitted at an emergency
60

- department. *Rev Esp Enferm Dig* 2009;101(9):610-18. doi: 10.4321/s1130-01082009000900003
3. Ohmann C, Franke C, Kraemer M, et al. Status report on epidemiology of acute appendicitis. *Chirurg* 2002;73(8):769-76. doi: 10.1007/s00104-002-0512-7
4. Körner H, Söreide JA, Pedersen EJ, et al. Stability in incidence of acute appendicitis. A population-based longitudinal study. *Dig Surg* 2001;18(1):61-66. doi: 10.1159/000050099
5. Martínez Carrilero J. Safety and efficacy of antibiotics compared with appendectomy for treatment of uncomplicated acute appendicitis: meta-analysis of randomised controlled trials. *Rev Clin Esp* 2012;212(9):460-60. doi: 10.1016/j.rce.2012.05.005
6. Varadhan KK, Neal KR, Lobo DN. Safety and efficacy of antibiotics compared with appendectomy for treatment of uncomplicated acute appendicitis: meta-analysis of randomised controlled trials. *BMJ (Clinical research ed)* 2012;344:e2156-e56. doi: 10.1136/bmj.e2156
7. Masoomi H, Nguyen NT, Dolich MO, et al. Laparoscopic appendectomy trends and outcomes in the United States: data from the Nationwide Inpatient Sample (NIS), 2004-2011. *Am Surg* 2014;80(10):1074-77.
8. Kotaluoto S, Ukkonen M, Pauniahio S-L, et al. Mortality Related to Appendectomy; a Population Based Analysis over Two Decades in Finland. *World J Surg* 2017;41(1):64-69. doi: 10.1007/s00268-016-3688-6
9. Blomqvist PG, Andersson RE, Granath F, et al. Mortality after appendectomy in Sweden, 1987-1996. *Ann Surg* 2001;233(4):455-60. doi: 10.1097/00000658-200104000-00001
10. Semm K. Endoscopic appendectomy. *Endoscopy* 1983;15(2):59-64. doi: 10.1055/s-2007-1021466
11. Xiao Y, Shi G, Zhang J, et al. Surgical site infection after laparoscopic and open appendectomy: a multicenter large consecutive cohort study. *Surg Endosc* 2015;29(6):1384-93. doi: 10.1007/s00464-014-3809-y [published Online First: 10/11]
12. Varela JE, Wilson SE, Nguyen NT. Laparoscopic surgery significantly reduces surgical-site infections compared with open surgery. *Surg Endosc* 2010;24(2):270-76. doi: 10.1007/s00464-009-0569-1 [published Online First: 06/17]
13. Bregendahl S, Nørgaard M, Laurberg S, et al. Risk of complications and 30-day mortality after laparoscopic and open appendectomy in a Danish region, 1998-2007; a population-based study of 18,426 patients. *Pol Przegl Chir* 2013;85(7):395-400. doi: 10.2478/pjs-2013-0060
14. Sauerland S, Jaschinski T, Neugebauer EA. Laparoscopic versus open surgery for suspected appendicitis. *The Cochrane database of systematic reviews* 2010(10):CD001546-CD46. doi: 10.1002/14651858.CD001546.pub3

15. Aziz O, Athanasiou T, Tekkis PP, et al. Laparoscopic versus open appendectomy in children: a meta-analysis. *Ann Surg* 2006;243(1):17-27. doi: 10.1097/01.sla.0000193602.74417.14
16. Dai L, Shuai J. Laparoscopic versus open appendectomy in adults and children: A meta-analysis of randomized controlled trials. *United European Gastroenterol J* 2017;5(4):542-53. doi: 10.1177/2050640616661931 [published Online First: 08/16]
17. Markides G, Subar D, Riyad K. Laparoscopic versus open appendectomy in adults with complicated appendicitis: systematic review and meta-analysis. *World J Surg* 2010;34(9):2026-40. doi: 10.1007/s00268-010-0669-z
18. Wei B, Qi C-L, Chen T-F, et al. Laparoscopic versus open appendectomy for acute appendicitis: a metaanalysis. *Surg Endosc* 2011;25(4):1199-208. doi: 10.1007/s00464-010-1344-z [published Online First: 09/17]
19. Li X, Zhang J, Sang L, et al. Laparoscopic versus conventional appendectomy--a meta-analysis of randomized controlled trials. *BMC Gastroenterol* 2010;10:129-29. doi: 10.1186/1471-230X-10-129
20. Horan TC, Gaynes RP, Martone WJ, et al. CDC definitions of nosocomial surgical site infections, 1992: a modification of CDC definitions of surgical wound infections. *Am J Infect Control* 1992;20(5):271-74. doi: 10.1016/s0196-6553(05)80201-9
21. Culver DH, Horan TC, Gaynes RP, et al. Surgical wound infection rates by wound class, operative procedure, and patient risk index. National Nosocomial Infections Surveillance System. *Am J Med* 1991;91(3B):152S-57S. doi: 10.1016/0002-9343(91)90361-z
22. Horan TC, Andrus M, Dudeck MA. CDC/NHSN surveillance definition of health care-associated infection and criteria for specific types of infections in the acute care setting. *Am J Infect Control* 2008;36(5):309-32. doi: 10.1016/j.ajic.2008.03.002
23. Badia JM, Casey AL, Petrosillo N, et al. Impact of surgical site infection on healthcare costs and patient outcomes: a systematic review in six European countries. *J Hosp Infect* 2017;96(1):1-15. doi: 10.1016/j.jhin.2017.03.004 [published Online First: 03/08]
24. Thompson KM, Oldenburg WA, Deschamps C, et al. Chasing zero: the drive to eliminate surgical site infections. *Ann Surg* 2011;254(3):430-37. doi: 10.1097/SLA.0b013e31822cc0ad
25. Hawn MT, Vick CC, Richman J, et al. Surgical site infection prevention: time to move beyond the surgical care improvement program. *Ann Surg* 2011;254(3):494-501. doi: 10.1097/SLA.0b013e31822c6929
26. Mehta JA, Sable SA, Nagral S. Updated recommendations for control of surgical site infections. *Ann Surg* 2015;261(3):e65-e65. doi: 10.1097/SLA.0b013e318289c5fd
27. Andersson RE. Short-term complications and long-term morbidity of laparoscopic and open appendectomy in a national cohort. *Br J Surg* 2014;101(9):1135-42. doi: 10.1002/bjs.9552 [published Online First: 06/30]

- 1
2
3 28. Pinkney TD, Calvert M, Bartlett DC, et al. Impact of wound edge protection devices on
4 surgical site infection after laparotomy: multicentre randomised controlled trial
5 (ROSSINI Trial). *BMJ (Clinical research ed)* 2013;347:f4305-f05. doi:
6 10.1136/bmj.f4305
7
- 8
9 29. Allegranzi B, Bagheri Nejad S, Combescure C, et al. Burden of endemic health-care-
10 associated infection in developing countries: systematic review and meta-analysis.
11 *Lancet (London, England)* 2011;377(9761):228-41. doi: 10.1016/S0140-
12 6736(10)61458-4 [published Online First: 12/09]
13
- 14
15 30. GlobalSurg Collaborative. Surgical site infection after gastrointestinal surgery in high-
16 income, middle-income, and low-income countries: a prospective, international,
17 multicentre cohort study. *Lancet Infect Dis* 2018;18(5):516-25. doi: 10.1016/S1473-
18 3099(18)30101-4 [published Online First: 02/13]
19
- 20
21 31. Danwang C, Mazou TN, Tochie JN, et al. Global prevalence and incidence of surgical site
22 infections after appendectomy: a systematic review and meta-analysis protocol. *BMJ*
23 *open* 2018;8(8):e020101-e01. doi: 10.1136/bmjopen-2017-020101
24
- 25
26 32. Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of observational studies in
27 epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in
28 Epidemiology (MOOSE) group. *JAMA* 2000;283(15):2008-12. doi:
29 10.1001/jama.283.15.2008
30
- 31
32 33. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic
33 reviews and meta-analyses of studies that evaluate healthcare interventions: explanation
34 and elaboration. *BMJ* 2009;339:b2700-b00. doi: 10.1136/bmj.b2700
35
- 36
37 34. Hoy D, Brooks P, Woolf A, et al. Assessing risk of bias in prevalence studies: modification
38 of an existing tool and evidence of interrater agreement. *J Clin Epidemiol*
39 2012;65(9):934-9. doi: 10.1016/j.jclinepi.2011.11.014 [published Online First:
40 2012/06/30]
41
- 42
43 35. Barendregt JJ, Doi SA, Lee YY, et al. Meta-analysis of prevalence. *Journal of epidemiology*
44 *and community health* 2013;67(11):974-78. doi: 10.1136/jech-2013-203104 [published
45 Online First: 08/20]
46
- 47
48 36. Cochran WG. The Combination of Estimates from Different Experiments. *Biometrics*
49 1954;10(1):101-29.
50
- 51
52 37. Higgins JPT, Thompson SG, Deeks JJ, et al. Measuring inconsistency in meta-analyses.
53 *BMJ (Clinical research ed)* 2003;327(7414):557-60. doi: 10.1136/bmj.327.7414.557
54
- 55
56 38. Viera AJ, Garrett JM. Understanding interobserver agreement: the kappa statistic. *Fam Med*
57 2005;37(5):360-63.
58
- 59
60 39. Egger M, Davey Smith G, Schneider M, et al. Bias in meta-analysis detected by a simple,
graphical test. *BMJ (Clinical research ed)* 1997;315(7109):629-34. doi:
10.1136/bmj.315.7109.629 [published Online First: 1997/10/06]

- 1
2
3 40. WHO. Global guidelines on the prevention of surgical site infection: WHO; 2016 [Available
4 from: [https://apps.who.int/iris/bitstream/handle/10665/250680/9789241549882-](https://apps.who.int/iris/bitstream/handle/10665/250680/9789241549882-eng.pdf?sequence=8)
5 [eng.pdf?sequence=8](https://apps.who.int/iris/bitstream/handle/10665/250680/9789241549882-eng.pdf?sequence=8) accessed Nov 23 2019.
6
7
8 41. Sievert DM, Ricks P, Edwards JR, et al. Antimicrobial-resistant pathogens associated with
9 healthcare-associated infections: summary of data reported to the National Healthcare
10 Safety Network at the Centers for Disease Control and Prevention, 2009-2010. *Infect*
11 *Control Hosp Epidemiol* 2013;34(1):1-14. doi: 10.1086/668770 [published Online First:
12 11/27]
13
14 42. Buggy D. Can anaesthetic management influence surgical-wound healing? *Lancet (London,*
15 *England)* 2000;356(9227):355-57. doi: 10.1016/S0140-6736(00)02523-X
16
17 43. Seidell JC, Halberstadt J. The global burden of obesity and the challenges of prevention.
18 *Ann Nutr Metab* 2015;66 Suppl 2:7-12. doi: 10.1159/000375143 [published Online
19 First: 06/02]
20
21 44. Checkley W, Ghannem H, Irazola V, et al. Management of NCD in low- and middle-income
22 countries. *Glob Heart* 2014;9(4):431-43. doi: 10.1016/j.ghheart.2014.11.003
23
24 45. Versporten A, Zarb P, Caniaux I, et al. Antimicrobial consumption and resistance in adult
25 hospital inpatients in 53 countries: results of an internet-based global point prevalence
26 survey. *Lancet Glob Health* 2018;6(6):e619-e29. doi: 10.1016/S2214-109X(18)30186-
27 4 [published Online First: 04/23]
28
29 46. Klein EY, Van Boeckel TP, Martinez EM, et al. Global increase and geographic
30 convergence in antibiotic consumption between 2000 and 2015. *Proceedings of the*
31 *National Academy of Sciences of the United States of America* 2018;115(15):E3463-
32 E70. doi: 10.1073/pnas.1717295115 [published Online First: 03/26]
33
34 47. Foster D, Kethman W, Cai LZ, et al. Surgical Site Infections after Appendectomy
35 Performed in Low and Middle Human Development-Index Countries: A Systematic
36 Review. *Surg Infect (Larchmt)* 2018;19(3):237-44. doi: 10.1089/sur.2017.188
37 [published Online First: 10/23]
38
39 48. Marchi M, Pan A, Gagliotti C, et al. The Italian national surgical site infection surveillance
40 programme and its positive impact, 2009 to 2011. *Euro Surveill* 2014;19(21):20815.
41 doi: 10.2807/1560-7917.es2014.19.21.20815
42
43 49. Udwardia TE. Diagnostic laparoscopy. *Surg Endosc* 2004;18(1):6-10. doi: 10.1007/s00464-
44 002-8872-0 [published Online First: 09/10]
45
46 50. Adisa AO, Lawal OO, Arowolo OA, et al. Local adaptations aid establishment of
47 laparoscopic surgery in a semiurban Nigerian hospital. *Surg Endosc* 2013;27(2):390-
48 93. doi: 10.1007/s00464-012-2463-5 [published Online First: 07/18]
49
50 51. Alfa-Wali M, Osaghae S. Practice, training and safety of laparoscopic surgery in low and
51 middle-income countries. *World J Gastrointest Surg* 2017;9(1):13-18. doi:
52 10.4240/wjgs.v9.i1.13
53
54
55
56
57
58
59
60


- 1
2
3 52. Haley RW, Culver DH, White JW, et al. The efficacy of infection surveillance and control
4 programs in preventing nosocomial infections in US hospitals. *American journal of*
5 *epidemiology* 1985;121(2):182-205. doi: 10.1093/oxfordjournals.aje.a113990
6
7
8 53. Harbarth S, Sax H, Gastmeier P. The preventable proportion of nosocomial infections: an
9 overview of published reports. *J Hosp Infect* 2003;54(4):258-321. doi: 10.1016/s0195-
10 6701(03)00150-6
11
12 54. Birgand G, Lepelletier D, Baron G, et al. Agreement among healthcare professionals in ten
13 European countries in diagnosing case-vignettes of surgical-site infections. *PloS one*
14 2013;8(7):e68618-e18. doi: 10.1371/journal.pone.0068618
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60


For peer review only

Table 1. Summary statistics of meta-analysis incidence of surgery site infections after appendectomy


	Incidence per 100 surgical procedures (95%CI)	95% Prediction interval	N studies	N participants	H (95%CI)	I ² (95%CI)	P heterogeneity	P Egger test	P difference
Global	7.0 (6.4-7.7)	1.0-17.7	226	729,434	8.9 (8.7-9.1)	98.7 (98.7-98.8)	< 0.0001	< 0.0001	-
- Low risk of bias	6.9 (6.0-7.9)	1.6-15.2	59	204,450	6.7 (6.3-7.1)	97.7 (97.4-98.0)	< 0.0001	< 0.0001	-
By Level of income									
- Low	11.1 (5.5-18.2)	0.0-42.2	9	1,496	3.8 (3.0-4.8)	93.1 (89.0-95.6)	< 0.0001	0.735	0.015
- Lower-middle	9.2 (6.3-12.6)	0.0-31.6	27	10,379	5.1 (4.6-5.7)	96.2 (95.3-96.9)	< 0.0001	0.960	
- Upper-middle	8.5 (6.5-10.8)	0.3-25.3	36	26,557	5.4 (2.9-5.9)	96.6 (95.9-97.1)	< 0.0001	0.392	
- High	6.2 (5.6-6.9)	0.9-15.3	154	691,002	9.5 (9.2-9.8)	98.9 (98.8-99.0)	< 0.0001	< 0.0001	
By WHO regions									
- Africa	12.6 (3.3-26.4)	0.0-72.5	8	3,001	9.1 (7.9-10.5)	98.8 (98.4-99.1)	< 0.0001	0.628	< 0.0001
- Western Pacific	9.6 (8.1-11.2)	2.3-20.8	43	30,822	3.8 (3.5-4.2)	93.2 (91.7-94.4)	< 0.0001	0.150	
- Eastern Mediterranean	8.2 (6.4-10.2)	1.7-18.6	23	7,779	2.6 (2.2-3.1)	85.3 (79.1-89.6)	< 0.0001	0.515	
- South-East Asia	7.6 (4.7-11.1)	0.0-24.6	16	5,782	3.8 (3.2-4.5)	93.0 (90.1-95.0)	< 0.0001	0.0001	
- Americas	5.9 (5.2-6.6)	1.9-11.7	67	401,931	7.5 (7.1-7.9)	98.2 (98.0-98.4)	< 0.0001	0.0004	
- Europe	5.8 (4.6-7.0)	0.0-19.1	68	276,793	10.4 (10.0-10.8)	99.1 (99.0-99.1)	< 0.0001	< 0.0001	
By type of surgical procedure									
- Laparoscopy with open surgery	4.6 (2.5-7.2)	0.0-15.6	10	4,892	3.2 (2.6-4.2)	90.7 (85.0-94.2)	< 0.0001	0.942	0.0002
- Laparoscopy	4.6 (3.4-5.9)	0.0-14.3	40	33,873	4.4 (4.0-4.8)	94.7 (93.6-95.7)	< 0.0001	0.0002	
- Open surgery	11.0 (7.9-14.4)	0.0-39.3	44	13,120	5.7 (5.2-6.1)	96.9 (96.4-97.3)	< 0.0001	0.077	

WHO: World Health Organization; CI: confidence interval; H: H statistics





1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Global incidence of surgical-site infection after appendectomy: a systematic review and meta-analysis

APPENDIX


Celestin **Danwang**, Jean Joel **Bigna**, Joel Noutakdie **Tochie**,
 Aime **Mbonda**, Clarence Mvalo **Mbanga**, Rolf Nyah Tuku **Nzalie**,
 Marc Leroy **Guifo**, Arthur **Essomba**

Supplementary Figure 1. Study flow	2
Supplementary Figure 2. Funnel plot for publication bias	3
Supplementary Table 1. Search strategy in EMBASE	4
Supplementary Table 2 : Characteristics of included studies	5
Supplementary Table 3. Individual characteristics of included studies	7
Reference list of included studies.....	23



Supplementary Figure 1. Study flow

review only



Supplementary Figure 2. Funnel plot for publication bias

Supplementary Table 1. Search strategy in EMBASE

	Search terms
#1	'appendectomy'/exp OR appendectomy OR 'appendicectomy'/exp OR appendicectomy OR appendices OR 'appendix epiploica' OR 'omental appendix' OR 'appendicitis'/exp OR appendicitis
#2	'surgical site infection'/exp OR 'surgical site infection' OR 'surgical wound infection'/exp OR 'surgical wound infection' OR 'surgical wound infections'/exp OR 'surgical wound infections' OR 'surgical site infections' OR 'operative site infections' OR 'postoperative wound infections'/exp OR 'postoperative wound infections' OR 'postoperative wound infection'/exp OR 'postoperative wound infection'
#3	[2000-2018]/py
#4	#1 AND #2 AND #3

Supplementary Table 2 : Characteristics of included studies

Characteristics	N = 226
Year of publication, range	2000-2018
%Male, range	0-100 (n = 195)
Mean/median age, range	7-74 (n = 186)
%HIV	0-13.1 (n = 2)
%Diabetes	0-95.7 (n = 34)
%Obesity	0-7.4 (n = 18)
Design, n	
- Cross sectional	120
- Cohort study	99
- Case control	7
WHO regions, n	
- Africa	8
- Americas	67
- Eastern Mediterranean	23
- Europe	68
- Multiregional	1
- South-East Asia	16
- Western Pacific	43
Level of income, n	
- Low	9
- Lower-middle	27
- Upper-middle	36
- High	154
Timing of data collection	
- Retrospective	123
- Prospective	101
- Unclear	2
Sampling	

1		
2		
3	- Consecutive	131
4		
5	- Systematic	37
6		
7	- Random	32
8		
9	- Exhaustive	11
10		
11	- Unclear	15
12		
13	Number of sites	
14		
15	- Multisite	51
16		
17	- One site	170
18		
19	- Unclear	5
20		
21	Pattern of appendicitis, range	
22		
23	- %Catarrhal	0-100 (n = 84)
24		
25	- %Perforated	0-100 (n = 110)
26		
27	- %Suppurated	0-100 (n = 70)
28		
29	- %Gangrenous	0-46.7 (n = 89)
30	%With administered antibiotics	24.1-100 (n = 109)
31		
32	%With administered analgesics	64.5-100 (n = 20)
33		
34	%With diet > 6 or 8 hours	50-100 (n = 3)
35		
36	Type of surgery	
37		
38	- %Open surgery	0-100 (n = 134)
39		
40	- %Laparoscopy	0-100 (n = 187)
41		
42	Mean/median time to complete the intervention (in hours), range	0.1-2.2 (n = 106)
43		
44	Type of anesthesia, n	
45		
46	- General	118
47		
48	- Spinal and general	2
49		
50	- Unclear	106
51		
52	SSI definition, n	
53		
54	- CDC-NNIS criteria	50
55		
56	- Other criteria	25
57		
58	- Not reported/Unclear	151
59		
60		

Supplementary Table 3. Individual characteristics of included studies

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Adejumo (1)	2015	Moderate	Cohort	Nigeria	Retrospective	Consecutive	One site	2007-2014	Adults	39	26	NR	Catarrhal, Perforated, Suppurated, Gangrenous	48.9	NR	NR	NR	100	NR	NR	NR	NR	141
Aguillo (2)	2005	Moderate	Cohort	Spain	Prospective	Consecutive	Unclear	NR	Children, Adolescents, Adults, Elderly	63.1	NR	NR	Unclear	NR	NR	NR	NR	100	NR	NR	NR	NR	792
Adhikar (3)	2008	Moderate	Cohort	Nepal	Prospective	Consecutive	One site	2005-2006	Children, Adolescents, Adults, Elderly	NR	NR	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy	0.5	General	NR	50
Ahmad(4)	2014	Moderate	Clinical trial	Pakistan	Prospective	Consecutive	One site	2012	Adults	35.4	27.4	NR	Perforated	0	100	0	0	100	Open Surgery	NR	General	NR	158
Ahmed (5)	2015	Moderate	Cross sectional	Pakistan	Retrospective	Consecutive	One site	2009-2010	Children, Adolescents	51.89	10.1	NR	Unclear	NR	NR	NR	NR	100	Open Surgery	NR	NR	NR	79
Akkoyun (6)	2012	Moderate	Case control	Turkey	Retrospective	Consecutive	One site	1998-2011	Children	64.5	8.9	NR	Perforated	0	100	0	0	100	Open Surgery	0.6	General	NR	234
Al-Saadi (7)	2006	Moderate	Cohort	Yemen	Retrospective	Consecutive	One site	2003-2005	Children, Adolescents, Adults	75	NR	NR	Catarrhal, Perforated, Suppurated, Gangrenous	NR	60	NR	13	100	Laparoscopy	NR	NR	NR	160
Al-Temimi (8)	2017	Low	Cohort	USA	Prospective	Systematic	One site	2016	Children, Adolescents, Adults, Elderly	40.2	30	NR	Catarrhal, Perforated, Suppurated, Gangrenous, Normal	73	17.4	3.3	3.3		Laparoscopy	0.6	NR	NR	92
Ali (9)	2012	Moderate	Cohort	Nigeria	Prospective	Consecutive	One site	2002-2009	Children, Adolescents, Adults, Elderly	33.9	32	NR	Catarrhal, Perforated, Suppurated, Gangrenous	NR	23.47	NR	NR		Open Surgery	NR	NR	NR	1257
Ali (10)	2015	Moderate	Cross sectional	Pakistan	Prospective	Consecutive	One site	2014	Adults	46.3	27.4	NR	Unclear	NR	NR	NR	NR	100	Open Surgery	NR	General	NR	121
Almström (11)	2016	Moderate	Cohort	Sweden	Retrospective	Systematic	One site	2006-2013	Children, Adolescents	59.5	NR	NR	Perforated, Non Perforated 76%	NR	24	NR	NR		Laparoscopy or Open Surgery	0.8	NR	NR	2756
Álvarez-Moreno (12)	2014	Low	Cohort	Colombia	Prospective	Systematic	Multisite	2008-2010	Children, Adolescents, Adults, Elderly	NR	NR	NR	Unclear	NR	NR	NR	NR		NR	NR	NR	According to CDC-NNIS diagnostic criteria	947
Andert (13)	2017	Moderate	Cohort	Germany	Retrospective	Consecutive	One site	2003-2014	Adults	48.6	30.5	NR	Catarrhal, Perforated, Suppurated, Gangrenous	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	NR	NR	Local signs of inflammation	2136
Anderson (14)	2014	Moderate	Cohort	Sweden	Retrospective	Exhaustive	Multisite	1992-2008	Adults	54	NR	NR	Perforated, Not perforated	NR	19.4	0	0		Laparoscopy or Open Surgery	NR	NR	NR	169896
Aranda-Narváez (15)	2010	Low	Cohort	Spain	Retrospective	Random	One site	1997-2009	Children, Adolescents, Adults, Elderly	63.3	35	NR	Suppurated, Gangrenous	0	0	53.3	46.7	100	Laparoscopy or Open Surgery	NR	General	According to CDC-NNIS diagnostic Criteria	30
Aranda-Narvaez (16)	2014	Moderate	Cohort	Spain	Not reported/Unclear	Not clear	One site	2007-2010	Adults	57	29	NR	Catarrhal, Perforated, Suppurated, Gangrenous	65.8	NR	NR	NR	62.00	Laparoscopy or Open Surgery	0.92	NR	According to CDC-NNIS diagnostic Criteria	868

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Arthur (17)	2017	Low	Cross sectional	Australia	Prospective	Systematic	Multisite	2016	Children, Adolescents, Adults, Elderly	49.5	31.4	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	1	NR	NR	1189
Asefa (18)	2002	High	Cross sectional	Ethiopia	Retrospective	Consecutive	One site	1997-1999	Adults	79.5	25.6	NR	Catarrhal, Perforated	45.4	44	0	0		NR	NR	NR	NR	200
Asefa (19)	2014	Moderate	Cross sectional	Ethiopia	Retrospective	Consecutive	One site	2006-2010	Children	62.1	10	NR	Catarrhal, Perforated, Suppurated, Gangrenous	32.3	59.6	6.2	1.9		NR	NR	NR	NR	322
Bae (20)	2016	Moderate	Cross sectional	USA	Retrospective	Systematic	One site	2010-2013	Children, Adolescents, Adults	NR	32	NR	Unclear	NR	NR	NR	NR	36.4	Laparoscopy	NR	NR	According to CDC-NNIS diagnostic criteria	143
Bae (21)	2016	Low	Cross sectional	Korea	Prospective	Systematic	One site	2014-2016	Adults	52	62	NR	Perforated, Suppurated, Gangrenous	NR	4	72	24	100	Laparoscopy	1.2	General	According to CDC-NNIS diagnostic criteria	25
Baek (22)	2011	Moderate	Cross sectional	Korea	Retrospective	Exhaustive	One site	2007-2009	Elderly	45.5	68.2	NR	Catarrhal, Perforated, Suppurated, Gangrenous	22.1	29.9	32.5	15.6	100	Laparoscopy or Open Surgery	1.05	General	NR	77
Bali (23)	2015	Moderate	Cohort	Turkey	Prospective	Consecutive	One site	2009-2013	Adults	35.7	32.33	NR	Unclear	NR	NR	NR	NR		Laparoscopy	1	NR	NR	126
Banghu (24)	2013	Low	Cohort	UK, Spain, Japan, Hong Kong, Australia, New Zealand	Prospective	Consecutive	Multisite	2012	Children, Adolescents, Adults, Elderly	51.1	NR	NR	Unclear	NR	NR	NR	NR	96.9	Laparoscopy or Open Surgery	NR	NR	According to CDC-NNIS diagnostic criteria	3326
Bansal (25)	2012	Low	Cohort	Switzerland	Prospective	Consecutive	One site	NR	Children	62	9.8	NR	Catarrhal, Perforated	74.3	25.7	NR	NR	49.2	Laparoscopy or Open Surgery	1.0	NR	According to CDC-NNIS diagnostic Criteria	187
Batajoo(26)	2012	Moderate	Cross sectional	Nepal	Retrospective	Consecutive	One site	2009-2012	Children, Adolescents, Adults, Elderly	45.6	29.6	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	0.8	NR	NR	226
Bharathi (27)	2011	Moderate	Cohort	Nepal	Prospective	Consecutive	One site	2008-2009	Children, Adolescents, Adults, Elderly	50	22.9	NR	Catarrhal, Perforated, Suppurated, Gangrenous	80	NR	NR	NR	100	Laparoscopy with Open Surgery	0.5	NR	NR	26
Bıçakcı (28)	2011	High	Cross sectional	Turkey	Retrospective	Systematic	Unclear	2006-2009	Children, Adolescents	64.5	10	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	NR	NR	279
Bickel (28)	2011	Moderate	Clinical trial	Israel	Prospective	Random	One site	2006-2009	Adults	73	28	NR	Catarrhal, Gangrenous, Phlegmonous 58.6 , Normal 4.3%	17	NR	NR	20.5	100	Open Surgery	0.5	General	NR	210
Blackwood (29)	2017	Moderate	Cross sectional	USA	Retrospective	Random	One site	2010-2015	Children	55.6	10.4	29.6	Unclear	NR	NR	NR	NR		Open Surgery	2	General	According to CDC-NNIS diagnostic Criteria	115
Blakely (31)	2011	Low	Clinical trial	USA	Prospective	Random	One site	2006-2009	Children, Adolescents	55.7	10.2	NR	Perforated	0	100	0	0	100	Laparoscopy or Open Surgery	1.9	NR	NR	131

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Bonadio (32)	2017	Moderate	Cross sectional	USA	Retrospective	Consecutive	One site	2008-2015	Children, Adolescents	56.2	8.4	NR	Perforated	0	100	0	0	100	Laparoscopy	NR	General	NR	203
Boomer (33)	2014	Low	Cohort	USA	Retrospective	Consecutive	One site	2010-2012	Children, Adolescents	61.1	10.9	NR	Catarrhal, Perforated, Gangrenous	66.2	NR	NR	NR	97.8	Laparoscopy or Open Surgery	NR	General	Wound infection or abdominal/pelvic abscess	1388
Boomer (34)	2016	Low	Cross sectional	USA	Retrospective	Systematic	Multisite	2010-2012	Children, Adolescents	60.3	11.0	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	NR	According to CDC-NNIS diagnostic criteria	1338
Bozkurt (35)	2014	Moderate	Case control	Turkey	Retrospective	Consecutive	One site	2008-2012	Children, Adolescents, Adults, Elderly	54	30.4	NR	Catarrhal	100	0	0	0		Laparoscopy	0.8	General	NR	1849
Brandt (36)	2008	Moderate	Cross sectional	Germany	Retrospective	Systematic	Multisite	2000-2004	Children, Adolescents, Adults, Elderly	NR	NR	NR	NR	NR	NR	NR	NR		NR	NR	NR	According to CDC-NNIS diagnostic criteria	10969
Brümmer (37)	2009	Moderate	Cohort	Germany	Retrospective	Consecutive	Multisite	2004-2007	Children, Adolescents, Adults, Elderly	NR	NR	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	NR	According to CDC-NNIS diagnostic Criteria	14209
Cairo (38)	2017	Moderate	Cohort	USA	Retrospective	Consecutive	Multisite	2012-2015	Children, Adolescents	61.3	11.0	29.9	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	General	NR	20981
Cameron (39)	2017	Low	Cohort	USA	Retrospective	Systematic	Multisite	2012-2015	Children, Adolescents	60.4	11	11.7	Unclear	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	NR	NR	According to CDC-NNIS diagnostic criteria	1389
Cao (40)	2015	Moderate	Cohort	China	Retrospective	Consecutive	Multisite	2011-2013	Adults	54.2	37.3	12.4	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	0.8	General	NR	12586
Cervantes-sánchez (41)	2000	Low	Clinical trial	Mexico	Prospective	Random	One site	1994-1995	Children, Adults	53.4	28	NR	Unclear	NR	NR	NR	NR		Open Surgery	NR	General	Pus or a positive bacteriologic culture from a wound discharge	350
Chamisa (42)	2009	High	Cross sectional	South Africa	Retrospective	Exhaustive	One site	2002-2004	Children, Adolescents, Adults, Elderly	78.4	NR	NR	Catarrhal, Perforated, Gangrenous, Normal	53	30.5	NR	10.2		Laparoscopy or Open Surgery	NR	NR	NR	324
Chaudhary (43)	2005	Moderate	Clinical trial	Pakistan	Prospective	Random	One site	1999-2003	Children, Adolescents, Adults, Elderly	45.4	NR	NR	Catarrhal	NR	0	0	0		Open Surgery	NR	NR	NR	677
Chen (44)	2011	High	Cross sectional	China	Prospective	Systematic	One site	2008-2009	Adults	NR	NR	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	0.8	Spinal and General	NR	69
Chen (45)	2012	High	Cross sectional	Taiwan	Prospective	Consecutive	One site	2010	Adults	60	38	NR	Unclear	NR	NR	NR	NR	73	Laparoscopy or Open Surgery	NR	General	NR	117
Chen (46)	2015	Moderate	Cohort	Taiwan	Retrospective	Consecutive	One site	2010-2012	Adults	43.6	42.5	NR	Catarrhal, Perforated	87.3	12.7	0	0		Laparoscopy or Open Surgery	NR	NR	NR	236
Chiang (47)	2006	Moderate	Cross sectional	Taiwan	Retrospective	Exhaustive	One site	2002-2004	Adults	59.7	35	NR	Catarrhal, Perforated	68	17	0	0	100	Open Surgery	1.1	General	NR	390

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Chiang (48)	2012	Moderate	Cohort	Taiwan	Prospective	Consecutive	One site	2008-2009	Adults	58.6	37.8	10	Unclear	NR	NR	NR	NR	100	Open Surgery	NR	General	Presence of gross or purulent discharge at the incision site with or without a positive bacterial culture	70
Cho (49)	2014	Low	Cross sectional	Korea	Prospective	Consecutive	One site	2011-2012	Adults	53	38.7	18.8	Unclear	NR	NR	NR	NR		Laparoscopy	NR	General	According to CDC-NNIS diagnostic Criteria	101
Choudhary (50)	2014	Moderate	Cross sectional	India	Prospective	Random	One site	2010-2013	Adults	67	NR	NR	Appendicular mass	0	0	0	0		NR	NR	NR	NR	92
Clyde (51)	2008	High	Cross sectional	USA	Retrospective	Systematic	One site	2002-2007	Children, Adolescents, Adults, Elderly	52	35	NR	Catarrhal, Perforated, Unclear	77	14	0	0		Laparoscopy or Open Surgery	NR	NR	NR	1198
Coakley (52)	2011	Low	Cohort	USA	Retrospective	Exhaustive	One site	2005-2010	Adults	47.3	28.7	NR	Catarrhal, Perforated, Suppurated, Gangrenous	38.3	1.2	47.1	9.8	100	Laparoscopy or Open Surgery	1.0	NR	According to CDC-NNIS diagnostic Criteria	728
Crandall (53)	2009	High	Cross sectional	USA	Retrospective	Not clear	One site	2004-2005	Adults	54	32.5	NR	NR	NR	74	NR	NR		Laparoscopy or Open Surgery	0.9	General	NR	176
Dede (54)	2008	Moderate	Cohort	Hungary	Prospective	Consecutive	One site	2005-2007	Children, Adolescents, Adults, Elderly	NR	NR	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	NR	NR	273
Dhiman (55)	2013	High	Cross sectional	USA	Retrospective	Not clear	Multisite	2003-2009	Adults	58	30.1	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	General	NR	1250
Dimitriou (56)	2013	Moderate	Cohort	Germany	Retrospective	Consecutive	One site	2007-2010	Children, Adolescents, Adults, Elderly	53.5	34.9	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	1	NR	NR	404
Durkin (57)	2015	Moderate	Cohort	USA	Retrospective	Consecutive	Multisite	2007-2012	Adults	NR	NR	NR	Unclear	NR	NR	NR	NR		Laparoscopy	NR	NR	NR	14763
Ein (58)	2013	Moderate	Cross sectional	Canada	Retrospective	Consecutive	One site	1969-2003	Children	70	7	NR	Perforated	0	100	0	0	78.8	Open Surgery	NR	General	1. Wound infection=pus draining from between the stitches or staples 2. Intra-abdominal abscess=presence of fever, abdominal pain and or gastrointestinal dysfunction and confirmed by radiologic evidence of intra-abdominal fluid collection	496

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Fukuda (59)	2016	Moderate	Cohort	Japan	Retrospective	Consecutive	Multisite	2007-2011	Children, Adolescents, Adults	54.4	64.5	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	1.3	General	NR	2074
Gandaglia (60)	2014	Low	Cohort	USA	Prospective	Consecutive	Multisite	2005-2011	Adolescents, Adults, Elderly	NR	NR	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	NR	General	According to CDC-NNIS diagnostic Criteria	97780
Garcell (61)	2016	Low	Cohort	Cuba	Prospective	Consecutive	One site	2013-2015	Children, Adolescents, Adults, Elderly	95.3	30.7	2.1	Unclear	NR	NR	NR	NR		Open Surgery	NR	NR	According to CDC-NNIS diagnostic Criteria	603
Ghnnam(62)	2011	Moderate	Cross sectional	Saudi Arabia	Retrospective	Not clear	One site	2007-2010	Adults	63.4	49.0	NR	Perforated, Unclear	NR	38.1	NR	NR		NR	NR	NR	NR	63
Giesen(63)	2017	Moderate	Cohort	Netherlands	Retrospective	Consecutive	Multisite	2014-2015	Children, Adults	54.3	31	NR	Catarrhal, Perforated, Suppurated, Gangrenous	48.2	17.3	23.2	11.3	100	Laparoscopy or Open Surgery	0.52	NR	According to CDC-NNIS diagnostic criteria	637
Giiti (64)	2010	Moderate	Cross sectional	Tanzania	Prospective	Systematic	One site	2008-2009	Children, Adolescents, Adults, Elderly	44.7	27	NR	Catarrhal, Perforated, Suppurated, Mass	87.4	7.0	1.5	0		NR	NR	NR	NR	199
Golub (65)	2016	Moderate	Cohort	Russia	Retrospective	Consecutive	Multisite	2012	Adolescents, Adults	NR	34.8	NR	Unclear	NR	NR	NR	NR	100	Open Surgery	NR	General	NR	332
Gross (66)	2016	Moderate	Cross sectional	USA	Retrospective	Consecutive	Multisite	2012-2013	Children, Adolescents	60.1	NR	17.8	Perforated	0	100	0	0		Laparoscopy or Open Surgery	NT	General	NR	2585
Gurien (67)	2016	Moderate	Cohort	USA	Retrospective	Consecutive	One site	2009-2012	Children, Adolescents	62	10.5	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	NR	General	Wound infections or intra abdominal abscesses	484
Hamzaoglu (68)	2004	Low	Cross sectional	Turkey	Prospective	Consecutive	One site	1999-2001	Adults	57	46.7	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	General	NR	100
Harmon (69)	2016	Low	Cohort	USA	Retrospective	Systematic	One site	2007-2012	Children, Adolescents, Adults, Elderly	47.4	39.7	NR	Non perforated	0	0	0	0		Laparoscopy or Open Surgery	NR	NR	According to CDC-NNIS diagnostic criteria	411
Helling (70)	2017	Low	Cross sectional	USA	Retrospective	Systematic	One site	2009-2014	Adults	64.3	34.4	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	NR	NR	According to CDC-NNIS diagnostic criteria	611
Helmer (71)	2002	Low	Cross sectional	USA	Retrospective	Systematic	One site	1998-1999	Children, Adolescents, Adults, Elderly	NR	NR	NR	Perforated, Non perforated	NR	19.4	NR	NR	100	NR	NR	NR	A surgical wound infection was defined as purulent drainage from the wound, cellulitis requiring antibiotics, or the opening of a closed wound. An intra-abdominal abscess was defined as an intraabdominal fluid collection that contained purulent material.	438
Hesami (72)	2014	Low	Clinical trial	Iran	Prospective	Random	Unclear	2010-2011	Children, Adolescents, Adults	58.9	27	NR	Unclear	NR	NR	NR	NR	100	Open Surgery	NR	NR	1..wound unfection=Purulent discharge, redness, inflammation, and the need to reoopen	90

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Hussain (73)	2012	Moderate	Clinical trial	Saudi Arabia	Prospective	Consecutive	One site	2010-2011	Adults	NR	32.2	NR	Catarrhal	100	0	0	0	100	Open Surgery	NR	NR	the wound 2...intra-abdominal abscess=abdominal pain, fullness, fever and confirmed by ecography 1..SSI=Pus discharge from wound needing its opening and drainage 2..Intra-abdominal collection=fluid collection inside the peritoneal cavity confirmed by ultrasound or CT scan that required drainage	377
Horvath (74)	2016	Moderate	Cross sectional	Germany	Retrospective	Consecutive	One site	2005-2013	Adults	47	28.6	NR	Perforated, phelgmonous	NR	52	NR	NR		Laparoscopy or Open Surgery	1.0	General	According to CDC-NNIS diagnostic Criteria	1516
Hughes (75)	2013	Moderate	Cross sectional	United Kingdom	Retrospective	Systematic	One site	2009-2010	Adults	55.6	30	NR	Unclear, simple and complicated	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	NR	General	NR	266
Hung (76)	2016	Moderate	Cross sectional	Vietnam	Prospective	Systematic	Multisite	2008-2010	Adults	45	41.6	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	1	General	NR	752
Iñigo (77)	2006	Low	Cohort	Spain	Prospective	Consecutive	One site	1998-2002	Adults	NR	NR	NR	Unclear	NR	NR	NR	NR		NR	0.7	NR	According to CDC-NNIS diagnostic Criteria	721
Iqbal (78)	2015	Low	Clinical trial	Pakistan	Prospective	Random	One site	2011	Adolescents, Adults, Elderly	66.3	26	NR	Catarrhal	100	0	0	0	100	Open Surgery	NR	General	According to Southampton criteria. Southampton grade 2 and above was considered as surgical site infection.	166
Javadi (79)	2017	Moderate	Clinical trial	Iran	Prospective	Random	One site	2016	Children, Adolescents, Adults	65	19.3	NR	Catarrhal, Suppurated, Gangrenous	NR	0	NR	NR		Open Surgery	0.5	General	NR	69
Jenkins (80)	2016	Low	Cohort	USA	Prospective	Systematic	Multisite	2006-2011	Children, Adolescents, Adults, Elderly	51.3	40.1	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	NR	According to CDC-NNIS diagnostic criteria	12410
Kang (81)	2012	Moderate	Case control	Korea	Retrospective	Random	One site	2010-2012	Adults	54.4	31.7	NR	Catarrhal, Perforated, Suppurated, Gangrenous	NR	5.1	NR	NR	100	Laparoscopy	1.1	General	NR	217
Kapischke (82)	2013	Low	Case control	Germany	Retrospective	Consecutive	One site	1999-2001	Children, Adolescents	47.8	11.5	NR	Catarrhal, Perforated	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	0.6	General	According to CDC-NNIS diagnostic criteria	159
Karam (83)	2016	Moderate	Cross sectional	USA	Retrospective	Not clear	One site	2010-2015	Children	62	12	NR	Perforated, Gangrenous	NR	20.6	NR	6.2		Laparoscopy	NR	General	NR	625
Karam (84)	2016	Moderate	Cross sectional	USA	Retrospective	Consecutive	One site	2010-2015	Children, Adolescents	63	NR	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy	1	General	NR	101
Kasatpibal (85)	2005	Low	Cross sectional	Thailand	Prospective	Systematic	Multisite	2003-2004	Children, Adolescents, Adults, Elderly	26.6	37.2	NR	Unclear	NR	NR	NR	NR	24.1	NR	0.8	NR	According to CDC-NNIS diagnostic criteria	1487
Kasatpibal (85)	2006	Moderate	Cohort	Thailand	Prospective	Not clear	Multisite	2003-2004	Children, Adolescents, Adults, Elderly	46.9	26	NR	Catarrhal	100	NR	NR	NR	92.2	Open Surgery	0.97	NR	According to CDC-NNIS diagnostic criteria	2139

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Kato (87)	2008	Low	Cohort	Japan	Prospective	Systematic	One site	2004-2006	Children	NR	9.4	NR	Perforated, Non perforated 75%	NR	25	NR	NR	100	Open Surgery	NR	NR	NR	64
Kell (88)	2003	Moderate	Cohort	Ireland	Prospective	Consecutive	Unclear	NR	Children, Adolescents, Adults, Elderly	75.2	20.7	NR	Unclear	NR	NR	NR	NR	100	Open Surgery	NR	General	Defined based on clinical and microbiological criteria	149
Khan (89)	2007	Low	Cohort	United Kingdom	Prospective	Consecutive	One site	2006	Children, Adolescents, Adults, Elderly	47.0	24	NR	Catarrhal, Perforated	63.4	20.1	0	0	100	Laparoscopy or Open Surgery	0.9	General	According to CDC-NNIS diagnostic Criteria	134
Khan (90)	2012	Moderate	Clinical trial	Pakistan	Prospective	Random	Multisite	2006-2009	Adults	69	33.3	NR	Unclear	NR	NR	NR	NR	100	Open Surgery	0.6	General	NR	100
Khan (91)	2014	Moderate	Clinical trial	Pakistan	Prospective	Random	Multisite	2013-2014	Children, Adolescents, Adults, Elderly	56.7	24	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	NR	NR	Observation of pain, redness, tenderness and purulent discharge	270
Khiria (92)	2011	Moderate	Cross sectional	India	Retrospective	Consecutive	One site	1999-2009	Adults	66	33.4	NR	Perforated, Gangrenous	NR	14.3	NR	5.2	100	Laparoscopy or Open Surgery	1.2	General	Any evidence of infection(erythema, purulent discharge, induration...) and requiring suture removal, antibiotic treatment, or evidence of dehiscence	497
Kilic (93)	2016	Moderate	Cross sectional	Turkey	Retrospective	Consecutive	One site	2004-2010	Children	62.1	9.5	NR	Perforated	0	100	0	0	100	Open Surgery	NR	NR	According to CDC-NNIS diagnostic Criteria	110
Kim (94)	2015	Low	Cross sectional	Korea	Retrospective	Systematic	One site	2008-2013	Children, Adolescents, Adults, Elderly	47.8	32.6	NR	Perforated, Suppurated, Gangrenous, Normal	6	13.8	64.5	7.1	100	Laparoscopy	0.7	NR	According to CDC-NNIS diagnostic criteria	2587
Kim (95)	2011	Moderate	Cross sectional	USA	Prospective	Consecutive	One site	2005-2008	Elderly	48.1	73.4	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	NR	General	According to CDC-NNIS diagnostic Criteria	3335
Kim(96)	2016	Moderate	Cohort	Korea	Retrospective	Consecutive	One site	2005-2012	Adults	59	NR	NR	Perforated, Gangrenous	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	1.9	General	NR	207
Kiriakopoulos (97)	2006	Moderate	Cross sectional	Greece	Retrospective	Consecutive	One site	2000-2004	Adults	73.8	42.3	NR	Perforated, Suppurated, Generalized peritonitis	0	61.9	9.5	0	100	Laparoscopy with Open Surgery	1.1	General	NRR	42
Kirshtein (98)	2009	Moderate	Cross sectional	Israel	Retrospective	Consecutive	One site	2000-2007	Adults	31.9	70.1	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	0.7	General	NR	1435
Kiudelis (99)	2013	Moderate	Cross sectional	Lithuania	Prospective	Consecutive	One site	2004-2009	Adults	46.3	32.3	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy	1.1	General	NR	152
Kleif (100)	2017	Moderate	Cross sectional	Denmark	Retrospective	Not clear	Multisite	2012-2014	Adults	53	47	NR	Suppurated, Gangrenous	NR	NR	NR	NR	98	Laparoscopy or Open Surgery	NR	General	NR	1151
Koizumi (101)	2014	Moderate	Cross sectional	Japan	Prospective	Consecutive	One site	2010	Adults	57.9	39.8	NR	Catarrhal, Perforated, Gangrenous, phelmong	6.4	6.4	NR	25 4	100	Laparoscopy or Open Surgery	0.9	General	NR	185
Kubota (102)	2014	Low	Clinical trial	Japan	Prospective	Random	One site	2008-2012	Children	63.6	NR	NR	Unclear	NR	NR	NR	NR	100	Open Surgery	NR	General	According to CDC-NNIS diagnostic Criteria	10

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Kumamoto (103)	2014	Low	Cohort	Japan	Prospective	Consecutive	One site	1997-2011	Adults	0	28	NR	Catarrhal, Gangrenous, Phlegmonous	21.8	NR	NR	33.4	100	Open Surgery	0.7	General	NR	124
Kumar (104)	2008	Moderate	Cohort	Pakistan	Prospective	Consecutive	One site	1997-2000	Children, Adolescents, Adults, Elderly	NR	NR	NR	Unclear	NR	NR	NR	NR	50	Laparoscopy or Open Surgery	0.7	NR	NR	100
Kumar (105)	2016	Moderate	Cohort	Nepal	Prospective	Consecutive	One site	2015-2016	Adolescents, Adults	49	33.9	NR	Catarrhal, Perforated, Suppurated, Gangrenous, Normal	88.6	4.7	1.9	2.4	Laparoscopy or Open Surgery	0.7	General	According to CDC-NNIS diagnostic Criteria	212	
Lacher (106)	2012	Moderate	Cohort	USA	Prospective	Consecutive	One site	2009-2011	Children, Adolescents	64.1	10.9	22.4	Catarrhal, Perforated	71.8	19	NR	NR	100	Laparoscopy	0.7	General	NR	415
Le (107)	2009	Moderate	Cross sectional	USA	Retrospective	Systematic	One site	1997-2007	Children, Adolescents, Adults, Elderly	52.1	31.8	NR	Catarrhal, Perforated, Gangrenous, Normal Appendix	92.9	2	0	2.2	86	Laparoscopy or Open Surgery	0.9	NR	According to CDC-NNIS diagnostic criteria	507
Lee (108)	2009	Low	Clinical trial	USA	Prospective	Random	One site	2006-2008	Children, Adolescents, Adults, Elderly	64.2	34.2	NR	Catarrhal, Perforated, Suppurated, Gangrenous	46.8	26.6	16.5	10.1	100	Open Surgery	NR	NR	Any significant subcutaneous SSI necessitating wound opening or treatment with antibiotics. This also included any subject who was prescribed a separate course of antibiotics after discharge from the hospital.	109
Lee (109)	2010	Moderate	Cross sectional	Taiwan	Prospective	Consecutive	One site	2006-2008	Children	58	11.1	NR	Unclear	NR	NR	NR	NR	Laparoscopy or Open Surgery	2	General	NR	264	
Lee (110)	2010	Moderate	Cross sectional	Korea	Retrospective	Consecutive	One site	2008-2009	Adults	49.3	26.7	NR	Perforated, Suppurated	0	26.7	73.3	0	Laparoscopy	1.0	General	NR	75	
Lee (111)	2011	Moderate	Cohort	USA	Retrospective	Consecutive	Multisite	1998-2007	Children, Adolescents	61.5	11	NR	Perforated, Non perforated	NR	25.7	NR	NR	Laparoscopy or Open Surgery	NR	General	NR	2462	
Lee (112)	2011	Moderate	Cohort	USA	Retrospective	Systematic	Multisite	1998-2007	Children, Adolescents	61	11.6	NR	Perforated, Non perforated 70.8%	NR	29.2	NR	NR	Laparoscopy or Open Surgery	NR	NR	NR	7650	
Lemieux (113)	2008	Moderate	Cohort	Canada	Retrospective	Consecutive	One site	1997-2007	Adults	0	28.8	NR	Perforated	NR	NR	NR	NR	Laparoscopy with Open Surgery	0.8	NR	NR	45	
Levy(114)	2013	Moderate	Cohort	USA	Retrospective	Consecutive	One site	2010-2011	Children	NR	NR	NR	Catarrhal, Perforated, Suppurated, Gangrenous	56.4	32.7	4.2	6.7	Laparoscopy or Open Surgery	NR	NR	NSQIP criteria	312	

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Li (115)	2005	Moderate	Cohort	China	Prospective	Consecutive	One site	2002-2004	Children, Adolescents	71.3	7.9	NR	Catarrhal, Suppurated, Gangrenous	11.0	0	69.4	19.7	100	Laparoscopy or Open Surgery	0.65	General	NR	160
Li (115)	2017	Moderate	Cohort	China	Retrospective	Consecutive	One site	2005-2016	Children	58.8	5.2	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	1	NR	Erythema, swelling and pus at the site of operation	398
Lim (116)	2011	Low	Cohort	Korea	Retrospective	Consecutive	One site	2009-2011	Adults	47.8	50.8	NR	Perforated, Gangrenous	0	61.6	NR	18.3	100	Laparoscopy or Open Surgery	1.3	General	Any evidence of infection (e.g., erythema, purulent discharge, induration, etc) requiring suture removal, antibiotics or dehiscence.	60
Lima (117)	2007	Moderate	Cross sectional	Spain	Retrospective	Consecutive	One site	2001-2006	Children, Adolescents, Adults, Elderly	NR	NR	NR	Catarrhal, Perforated, Suppurated, Gangrenous	53	2	26.3	9.7		NR	NR	NR	NR	300
Lin (118)	2006	Moderate	Cross sectional	Taiwan	Retrospective	Consecutive	One site	2001-2003	Adults	57.6	37.5	NR	Perforated	NR	100	NR	NR	100	Laparoscopy or Open Surgery	1.4	Not described	NR	229
Litz (119)	2016	Moderate	Cohort	USA	Retrospective	Consecutive	One site	2012-2015	Children, Adolescents	NR	11.4	17.7	Catarrhal, Perforated, Suppurated, Gangrenous, Interval, Normal	54	11.4	15.0	11.9		Laparoscopy	0.5	General	NR	413
Liu (120)	2017	High	Cross sectional	China	Retrospective	Consecutive	Unclear	2015-2016	Children	53.6	6.6	NR	Catarrhal, Suppurated, Gangrenous	34.8	0	38.4	26.8		Laparoscopy or Open Surgery	1.0	General	NR	112
Mahmood (121)	2016	Moderate	Clinical trial	Pakistan	Prospective	Random	One site	2012	Children, Adolescents, Adults	55.5	22.3	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	NR	General	Based on Wound Asepsis Score	200
Martinez (122)	2012	Moderate	Cross sectional	Spain	Retrospective	Random	One site	2011	Adults	60	35.8	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	General	NR	75
Mbah (123)	2006	Moderate	Cohort	Nigeria	Prospective	Consecutive	One site	2005	Children, Adolescents, Adults, Elderly	70	25	NR	Unclear	NR	NR	NR	NR		Open Surgery	NR	General	NR	136
Memon (124)	2017	Moderate	Clinical trial	Pakistan	Prospective	Random	One site	2014-2016	Adults	53.3	26	NR	Catarrhal	100	0	0	0	100	Laparoscopy or Open Surgery	NR	NR	NR	227
Menezes (125)	2008	Moderate	Cross sectional	Ireland	Retrospective	Consecutive	One site	2000-2006	Children, Adolescents	62.7	10.5	NR	Perforated, Gangrenous	0	81.4	0	17.8		Laparoscopy or Open Surgery	NR	NR	NR	118
Merenda (126)	2013	Moderate	Cross sectional	Poland	Retrospective	Consecutive	One site	2006-2012	Children, Adolescents, Adults, Elderly	NR	NR	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	NR	NR	317
Michailidou (127)	2015	Low	Cross sectional	USA	Retrospective	Systematic	One site	2007-2013	Children, Adolescents	56.1	9.6	NR	Perforated, Negative appendectomy	NR	26.5	NR	NR		Laparoscopy or Open Surgery	1.3	NR	According to CDC-NNIS diagnostic criteria	264
Michailidou (128)	2015	Moderate	Cross sectional	USA	Retrospective	Consecutive	Multisite	2012	Children, Adolescents	60.1	11.2	22.5	Catarrhal, Perforated, Suppurated, Gangrenous	NR	NR	NR	NR	100	Laparoscopy	0	General	NR	2812

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Mickovic (129)	2015	Moderate	Cross sectional	Serbia	Retrospective	Not clear	One site	2010	Children	46.4	11.7	NR	Catarrhal, Perforated, Gangrenous	45.9	2.2	NR	19.5	100	Laparoscopy or Open Surgery	0.7	General	NR	218
Ming (130)	2009	Moderate	Cross sectional	China	Retrospective	Consecutive	One site	2003-2005	Adults	57.2	48.8	NR	Perforated, Gangrenous, Appendicular abscess	NR	72.3	NR	38.2	100	Laparoscopy or Open Surgery	NR	General	NR	173
Mingmalairak(131)	2009	Low	Clinical trial	Thailand	Prospective	Random	One site	2006-2007	Adults	61	29.5	0	Catarrhal, Perforated, Suppurated, Gangrenous	24	16	52	8.0	100	Open Surgery	43	General	NR	100
Miyano (132)	2010	Low	Cohort	Japan	Prospective	Consecutive	One site	2004-2008	Children, Adolescents	56.5	7.7	NR	Peritonitis complicating appendicitis	0	100	0	0	100	Laparoscopy or Open Surgery	1.9	General	NR	23
Moazzez (133)	2013	Low	Cohort	USA	Retrospective	Not clear	One site	2005-2009	Elderly	49.3	74	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	General	NR	2060
Monge Jodra (134)	2003	Moderate	Cohort	Spain	Prospective	Consecutive	Multisite	1997-2000	Children, Adolescents, Adults, Elderly	NR	NR	NR	Unclear	NR	NR	NR	NR		NR	NR	NR	According to CDC-NNIS diagnostic Criteria	5780
Mueck (135)	2017	Moderate	Cohort	USA	Prospective	Consecutive	One site	2012-2015	Children, Adolescents	62.5	11.0	NR	Catarrhal, Suppurated	NR	NR	NR	NR	95	Laparoscopy with Open Surgery	NR	General	NR	697
Muensterer (136)	2011	Low	Cohort	USA	Prospective	Consecutive	One site	2009-2010	Children, Adolescents	NR	11.2	NR	Catarrhal, Perforated	78.1	10.4	0	0	100	Laparoscopy	0.6	General	Infected umbilicus requiring antibiotics, or incision and drainage	183
Muensterer (137)	2009	Moderate	Cross sectional	USA	Prospective	Consecutive	One site	2009	Children	61.3	11	NR	Perforated	NR	21.4	NR	NR		Laparoscopy	0.73	General	NR	75
Mustafa (138)	2016	Low	Clinical trial	Pakistan	Prospective	Random	One site	2015-2016	Adults	52.9	26.6	NR	Perforated	0	100	0	0	100	Open Surgery	NR	NR	Redness around the wound, serosanguinous discharge, fever > 100°F	68
Nadler (139)	2003	High	Cross sectional	USA	Retrospective	Systematic	One site	1998-2001	Children	62.2	9.35	NR	Perforated	NR	100	NR	NR	100	Open Surgery	NR	General	NR	94
Nataraja (140)	2010	Moderate	Cohort	United Kingdom	Retrospective	Consecutive	One site	2008-2010	Children, Adolescents	59.1	11	NR	Catarrhal, Perforated, Suppurated	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	General	NR	191
Nataraja (141)	2012	Moderate	Case control	United Kingdom	Retrospective	Consecutive	Multisite	2003-2010	Children, Adolescents	58.2	11.3	NR	Catarrhal, Perforated, Suppurated, Gangrenous	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	General	Post op intra abdominal abscess	1205
Obayashi (142)	2015	Moderate	Cross sectional	Japan	Retrospective	Consecutive	One site	2006-2014	Children, Adolescents	60	11	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	NR	NR	485
Obinwa (143)	2015	Moderate	Cohort	Ireland	Retrospective	Consecutive	One site	1995-2008	Children	54.5	9.6	NR	Catarrhal, Perforated, Suppurated, Gangrenous	62.7	NR	NR	4.2	100	NR	NR	NR	NR	1037
Ohene (144)	2006	Moderate	Cross sectional	Ghana	Prospective	Consecutive	One site	1998-2004	Adults	63.9	32.4	NR	Unclear	NR	NR	NR	NR	100	Open Surgery	NR	General	NR	638
Okkyung (145)	2002	Moderate	Clinical trial	Korea	Prospective	Random	One site	2002	Children, Adolescents, Adults, Elderly	54.7	30.5	NR	Catarrhal, Suppurated, Gangrenous	NR	0	50	27.3	100	Open Surgery	NR	General	NR	84

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Onieva (146)	2017	Moderate	Cross sectional	Spain	Retrospective	Consecutive	One site	2012-2014	Children, Adolescents, Adults, Elderly	53.7	32	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	NR	NR	294
Page (147)	2010	Moderate	Cross sectional	USA	Retrospective	Exhaustive	Multisite	2008	Adults	51.4	39.2	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	0.94	General	NR	17199
Palesty (148)	2004	Moderate	Cross sectional	USA	Retrospective	Consecutive	One site	2000-2002	Adults	47	25.2	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy	1.2	General	NR	50
Pandit (149)	2016	High	Cohort	Nepal	Retrospective	Consecutive	Multisite	2009-2014	Children, Adolescents, Adults	51	24.3	NR	Perforated, Suppurated	NR	2.6	97.4	0		Open Surgery	0.6	Spinal and General	NR	101
Parcells (150)	2009	Low	Cohort	USA	Retrospective	Systematic	One site	1997-2007	Adults	NR	39.3	NR	Perforated, Not perforated	NR	33.1	NR	NR		Laparoscopy or Open Surgery	NR	NR	According to CDC-NNIS diagnostic criteria	1063
Park (151)	2017	Low	Cohort	Korea	Prospective	Systematic	One site	2012-2014	Adults	53.3	37.45	NR	Perforated, Gangrenous	NR	38.7	NR	45.9		Laparoscopy	1.1	General	According to CDC-NNIS diagnostic Criteria	1343
Park (152)	2018	Moderate	Cohort	Korea	Retrospective	Consecutive	One site	2009-2013	Adults	53.7	34.2	NR	Perforated	NR	13.2	NR	NR		Laparoscopy with Open Surgery	1.1	General	NR	986
Pascual (153)	2017	Moderate	Cohort	Spain	Prospective	Consecutive	One site	2013-2017	Adults	49	41	NR	Unclear	NR	NR	NR	NR		Laparoscopy	0.1	General	NR	100
Patel (154)	2003	High	Cohort	Kenya	Retrospective	Consecutive	One site	1996-2002	Children, Adolescents, Adults	30.2	30.6	NR	Catarrhal, Suppurated, Gangrenous, Carcinoid tumor	94.3	0	0.9	2.8	100	Laparoscopy with Open Surgery	1.5	General	NR	106
Pearcy (155)	2017	Moderate	Case control	USA	Retrospective	Random	Multisite	2010-2014	Adults	54	36	NR	Unclear	NR	NR	NR	NR		NR	1.1	NR	NR	2903
Pishori (156)	2003	Low	Cross sectional	Pakistan	Prospective	Systematic	One site	1997-1999	Children, Adolescents, Adults, Elderly	NR	NR	NR	Unclear	NR	NR	NR	NR		NR	NR	NR	According to CDC-NNIS diagnostic criteria	3304
Putnam (157)	2016	Moderate	Cross sectional	USA	Prospective	Consecutive	One site	2012-2015	Children, Adolescents	61	9.4	NR	Perforated, Suppurated, Gangrenous	0	100	0	0	100	Laparoscopy or Open Surgery	0.9	General	NR	410
Qahtani (158)	2014	Moderate	Cohort	Saudi Arabia	Prospective	Random	One site	2012	Adolescents, Adults	68	23.6	NR	Catarrhal, Perforated, Gangrenous	9.6	19.9	NR	22.4	100	Open Surgery	1.5	General	NR	91
Quezada(159)	2015	Moderate	Cohort	Chile	Retrospective	Consecutive	One site	2003-2013	Adults	43	39	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	2.2	NR	NR	227
Raakow (160)	2014	High	Cohort	Germany	Prospective	Not clear	One site	2009-2013	Adolescents, Adults	28.8	27.2	NR	Catarrhal, Suppurated, Gangrenous	12.8	0	16	4		Laparoscopy	0.8	General	NR	156
Rafiq (161)	2015	Low	Clinical trial	Pakistan	Prospective	Random	One site	2012-2014	Adolescents, Adults, Elderly	48.5	22.6	0	Unclear	NR	NR	NR	NR	100	Open Surgery	0.7	General	NR	390
Rajabi-Mashhadi (162)	2012	Moderate	Clinical trial	Iran	Prospective	Random	One site	2006-2007	Adults	62.5	26.2	NR	Unclear, Non perforated	NR	NA	NR	NR	100	Open Surgery	NR	NR	NR	291

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Reinisch (163)	2017	Moderate	Cross sectional	Germany	Retrospective	Consecutive	One site	2008-2015	Adults	56	32	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy with Open Surgery	NR	General	NR	680
Rios (164)	2004	High	Cross sectional	Peru	Not reported/Unclear	Consecutive	One site	2001-2002	Children, Adolescents, Adults, Elderly	NR	30.6	NR	Catarrhal, Perforated, Suppurated, Gangrenous	15.38	16.35	44.23	24.04	100	NR	0.98	Unclear	NR	104
Romano (165)	2014	Moderate	Cross sectional	USA	Retrospective	Systematic	One site	2010-2012	Adults	66	35.7	NR	Gangrenous	NR	NR	NR	9.7	86	Laparoscopy	NR	General	NR	372
Romel Hilaire (166)	2014	Moderate	Cross sectional	Cuba	Retrospective	Consecutive	One site	2007-2009	Adults	100	NR	NR	Suppurated	0	0	100	0		Laparoscopy or Open Surgery	NR	NR	NR	720
Romero (167)	2009	Moderate	Cross sectional	Colombia	Retrospective	Consecutive	One site	1997	Children, Adolescents, Adults, Elderly	NR	NR	NR	Unclear	NR	NR	NR	NR		Laparoscopy	NR	NR	NR	310
Romy (168)	2008	Low	Cross sectional	Switzerland	Prospective	Systematic	Multisite	1998-2004	Children, Adolescents, Adults, Elderly	53.9	32.7	NR	Unclear	NR	NR	NR	NR	59.5	Laparoscopy or Open Surgery	NR	NR	According to CDC-NNIS diagnostic criteria	2468
Rooh-ul-Muqim (169)	2010	Moderate	Cohort	Pakistan	Prospective	Consecutive	One site	2008-2009	Adolescents, Adults, Elderly	48.5	24	NR	Catarrhal, Perforated, Suppurated, Gangrenous	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	0.5	General	NR	165
Rossem (170)	2015	Moderate	Cohort	Netherlands	Prospective	Consecutive	Multisite	2014	Adults	47.5	44	NR	Perforated, Gangrenous	NR	68	10.4	21.7	100	Laparoscopy with Open Surgery	0.9	General	NR	415
Rossem (171)	2016	Low	Cohort	Netherlands	Prospective	Not clear	Multisite	2014	Children, Adolescents, Adults	46.2	28.0	NR	Unclear	NR	NR	NR	NR		Laparoscopy	0.8	General	Superficial surgical site infection: recorded when administration of antibiotics, opening of the incision or both was necessary. An intra-abdominal abscess was defined as a postoperative intra-abdominal fluid collection diagnosed by cross-sectional imaging for which administration of antibiotics or a radiological or surgical intervention was needed.	1995
Rotermann (172)	2004	Moderate	Cohort	Canada	Retrospective	Consecutive	Multisite	1997-2000	Children, Adolescents, Adults, Elderly	55.2	NR	NR	Unclear	NR	NR	NR	NR		NR	NR	NR	According to CDC-NNIS diagnostic criteria	80867

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Saar (173)	2016	Low	Cross sectional	Estonia	Prospective	Consecutive	One site	2013-2014	Adults	48.9	35.4	NR	Perforated, Gangrenous	NR	15.4	NR	59.4	95.1	Laparoscopy or Open Surgery	0.7	General	According to CDC-NNIS diagnostic Criteria	266
Saber (174)	2010	Moderate	Clinical trial	USA	Prospective	Consecutive	One site	2008-2009	Adults	42.3	33	NR	Catarrhal	100	0	0	0		Laparoscopy	0.8	NR	NR	26
Sadraei-Mosavi (175)	2017	Moderate	Clinical trial	Iran	Prospective	Random	One site	2013-2014	Adults	NR	28.4	NR	Catarrhal	100	0	0	0	100	Open Surgery	NR	NR	SSI=pus discharge from wound, redness, tenderness, edema	152
Saha (176)	2010	Moderate	Cohort	Bangladesh	Prospective	Consecutive	One site	2007-2008	Children	NR	NR	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	NR	NR	NR	60
Sahm (177)	2010	Moderate	Cross sectional	Germany	Prospective	Systematic	One site	1998-2006	Adults	54	39	NR	Catarrhal, Perforated, Gangrenous	50.7	17.0	NR	6.9	100	Laparoscopy or Open Surgery	1.0	General	NR	1710
Sahm (178)	2015	Moderate	Cross sectional	Germany	Prospective	Exhaustive	Multisite	1988-2009	Children, Adolescents, Adults, Elderly	43	31	NR	Perforated, Non Perforated	91.5	NR	8.5	NR	NR	Laparoscopy or Open Surgery	NR	NR	NR	12570
Saló (179)	2016	High	Cohort	Sweden	Retrospective	Consecutive	One site	2006-2014	Children	55.6	10.4	NR	Perforated, Gangrenous, Phlegmonous	NR	7.3	NR	11.6	100	Laparoscopy	0.94	NR	NR	259
Sanchez-Santana (180)	2017	Low	Cohort	Spain	Prospective	Consecutive	One site	2007-2015	Adults	55.2	32.9	2.6	Unclear	NR	NR	NR	NR	71.3	Laparoscopy or Open Surgery	NR	NR	According to CDC-NNIS diagnostic criteria	930
Sauvain (181)	2016	Moderate	Cohort	Switzerland	Retrospective	Consecutive	Multisite	2007-2011	Adults	53.2	34	NR	Catarrhal, Perforated, Suppurated, Gangrenous	NR	19	NR	NR		Laparoscopy or Open Surgery	NR	NR	NR	2559
Scarborough (182)	2012	Low	Cross sectional	USA	Retrospective	Systematic	Multisite	2005-2009	Children, Adolescents, Adults, Elderly	52	38.9	NR	Perforated, Non rupture	NR	11.2	NR	NR		Laparoscopy or Open Surgery	0.9	NR	According to CDC-NNIS diagnostic criteria	39122
Scarless (182)	2013	Moderate	Clinical trial	Scotland	Prospective	Random	One site	2011	Adults	53	32	NR	Unclear	NR	NR	NR	NR		Laparoscopy	1.4	General	NR	38
Seifarth (183)	2016	Moderate	Cohort	USA	Retrospective	Consecutive	Multisite	2007-2012	Children, Adults	60	12	NR	Catarrhal	100	0	0	0		Laparoscopy	NR	General	NR	1283
Seifarth (184)	2016	Low	Cohort	USA	Retrospective	Consecutive	Multisite	2007-2012	Children, Adolescents, Adults	60	12	NR	Perforated, Suppurated, Gangrenous	100	0	0	0		Laparoscopy	NR	NR	NR	1283
Senekjan (185)	2013	Moderate	Cohort	USA	Retrospective	Consecutive	Multisite	2005-2009	Adolescents	56.5	40.3	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	0.9	NR	1) SSI (superficial and deep incisional)...infection within 30 days of operation and involved skin, subcutaneous tissue or deep soft tissue 2) Organ space infection (OSI)...infection within 30 days of operation when the infection appeared to be related to the operation and involved any part of the anatomy other than the incision	61830

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Seo (186)	2002	Moderate	Cross sectional	Korea	Retrospective	Systematic	One site	2000	Adults	0	NR	NR	Catarrhal, Perforated, Suppurated, Gangrenous	14.7	15.6	49.5	20.2		NR	NR	NR	NR	129
Sesia (187)	2011	Moderate	Cohort	Germany	Prospective	Consecutive	One site	2006-2008	Children, Adolescents, Adults, Elderly	NR	NR	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	1	NR	NR	265
Shaikh (188)	2011	Moderate	Cross sectional	Pakistan	Prospective	Consecutive	One site	2007-2009	Adults	51.4	29	NR	Catarrhal, Perforated, Suppurated	82.86	8.67	1.51	0		Open Surgery	NR	General	NR	461
Shang (189)	2017	Moderate	Cohort	China	Retrospective	Consecutive	One site	2013-2016	Adults	54.3	2.2	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	NR	Erythema, swelling and purulent discharge	398
Shimizu (102)	2014	Low	Cross sectional	Japan	Retrospective	Not clear	One site	2000-2012	Adults	44	35	NR	Catarrhal, Gangrenous	19	NR	NR	37		Open Surgery	NR	General	According to CDC-NNIS diagnostic Criteria	300
Shindholimath (190)	2011	Moderate	Cross sectional	India	Retrospective	Consecutive	One site	2007-2009	Adults	68.4	NR	NR	Perforated, Suppurated, Gangrenous, Appendicular abscess	0	36.8	5.3	26.3	100	Laparoscopy	1.6	General	NR	19
Siam (191)	2017	Moderate	Cohort	Israel	Retrospective	Consecutive	One site	2008-2015	Adults	62.8	34.1	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy with Open Surgery	0.7	General	NR	1649
Silva (192)	2008	Moderate	Cohort	Chile	Prospective	Random	One site	2005-2006	Adults	58.9	NR	NR	Unclear	NR	NR	NR	NR		Open Surgery	NR	General	NR	433
Singh (193)	2017	Moderate	Clinical trial	India	Prospective	Consecutive	One site	2014-2015	Adults	43.2	28.7	11.4	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	General	NR	44
Siribumrungwong (194)	2013	Low	Cohort	Thailand	Retrospective	Systematic	One site	2006	Adults	65	37	NR	Perforated	NR	100	NR	NR	100	Open Surgery	1.2	NR	According to CDC-NNIS diagnostic criteria	128
Sivrikoz (195)	2015	Moderate	Cohort	USA	Retrospective	Exhaustive	Multisite	2004-2010	Children, Adolescents, Adults, Elderly	52.1	48	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	0.9	NR	NR	4844
Soll (196)	2016	Low	Cohort	Switzerland	Retrospective	Consecutive	One site	2009-2013	Children, Adolescents, Adults, Elderly	54.7	26.5	NR	Catarrhal, Perforated, Suppurated, Gangrenous	NR	46	NR	NR	100	Laparoscopy	1	NR	According to CDC-NNIS diagnostic Criteria	813
Sozutek (197)	2013	Low	Clinical trial	Turkey	Retrospective	Consecutive	One site	2010-2011	Adults	44	30.9	NR	Catarrhal, Perforated	57	20	NR	NR		Laparoscopy or Open Surgery	0.5	General	NR	75
Srishewachart (198)	2016	Moderate	Cross sectional	Thailand	Retrospective	Consecutive	One site	2012-2014	Children, Adolescents, Adults, Elderly	52	43.7	7.4	Unclear	NR	NR	NR	NR		Open Surgery	NR	General	NR	450
Staszewicz (199)	2014	Moderate	Cohort	Switzerland	Prospective	Systematic	Multisite	1998-2011	Children, Adolescents, Adults, Elderly	54	34.2	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	1	NR	According to CDC-NNIS diagnostic Criteria	6383
Suttie (200)	2004	High	Case control	Scotland	Retrospective	Not clear	One site	1997-2002	Children	50	10.8	NR	Perforated, Suppurated, Gangrenous	0	2	50	14		Laparoscopy	1	General	NR	60
Svensson (201)	2016	Moderate	Cohort	Sweden	Prospective	Consecutive	One site	2006-2010	Children, Adolescents	60.2	11.3	NR	Catarrhal, Perforated, Suppurated, Gangrenous, not described	6.6	21.8	44.6	29.8	100	Laparoscopy or Open Surgery	0.7	General	NR	1745

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Taguchi (202)	2015	Moderate	Clinical trial	Japan	Prospective	Random	One site	2009-2014	Adults	65.43	47.5	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	1.2	General	According to CDC-NNIS diagnostic Criteria	81
Tanaka (203)	2007	Moderate	Cohort	Japan	Retrospective	Consecutive	One site	2002-2005	Children	54.3	2.2	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	NR	NR	95
Tijerina (204)	2010	Low	Clinical trial	Mexico	Prospective	Exhaustive	One site	2005-2007	Children, Adolescents, Adults, Elderly	46	NR	NR	Unclear	NR	NR	NR	NR	100	Open Surgery	NR	General	NR	529
Toro Pablo (205)	2017	Moderate	Cohort	Spain	Retrospective	Consecutive	One site	2012-2016	Children, Adolescents, Adults	NR	26	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	NR	NR	600
Towfigh (206)	2011	Low	Clinical trial	USA	Prospective	Random	One site	2007-2009	Adults	77.3	33	NR	Perforated	0	100	0	0		Open Surgery	NR	NR	NR	75
Troillet (207)	2017	Low	Cohort	Switzerland	Prospective	Consecutive	Multisite	2011-2015	Children, Adolescents, Adults, Elderly	NR	NR	NR	Unclear	NR	NR	NR	NR	92.2	Laparoscopy or Open Surgery	NR	NR	According to CDC-NNIS diagnostic Criteria	15439
Tsioplis (208)	2013	Moderate	Cross sectional	Germany	Retrospective	Consecutive	One site	1999-2008	Children, Adolescents, Adults, Elderly	51	23	9	Catarhal, Perforated, Suppurated, Gangrenous	19	NR	50	25	75	Laparoscopy or Open Surgery	NR	Not reported	NR	1439
Vahdad (209)	2016	Moderate	Cross sectional	Germany	Retrospective	Systematic	One site	2008-2012	Children, Adolescents	52.4	NR	NR	Catarhal, Perforated, Phelgmonous in 43% of cases	48.2	8.7	NR	NR		Laparoscopy	1.1	NR	NR	309
Van Rossem (210)	2016	High	Cohort	Netherlands	Prospective	Consecutive	Multisite	2014	Adults	49.7	39.0	NR	Catarhal, Perforated, Gangrenous	73.7	11.0	NR	9.9	96.6	Laparoscopy or Open Surgery	0.72	NR	NR	1378
Van Rossem (170)	2014	High	Cohort	Netherlands	Retrospective	Consecutive	Multisite	2004-2010	Adults	53.2	49	NR	Perforated	0	100	0	0	100	Laparoscopy or Open Surgery	0.85	NR	NR	267
Wang-Chan (211)	2017	Low	Cross sectional	Switzerland	Retrospective	Consecutive	One site	2013-2014	Children, Adolescents, Adults, Elderly	55.3	47	13.8	Unclear	NR	NR	NR	NR		Laparoscopy with Open Surgery	NR	General	According to CDC-NNIS diagnostic Criteria	246
Watanabe (212)	2011	Low	Cross sectional	Japan	Prospective	Consecutive	Multisite	2005-2006	Adults	59.4	63.8	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	General	According to CDC-NNIS diagnostic Criteria	903
Willis (213)	2016	Moderate	Cohort	USA	Prospective	Consecutive	One site	2013-2014	Children, Adolescents	58.5	8.8	NR	Unclear	NR	NR	NR	NR	100	Laparoscopy or Open Surgery	NR	General	NR	313
Wong (214)	2015	High	Cohort	Peru	Prospective	Not clear	Multisite	2005-2010	Adults	NR	NR	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	General	NR	352
Wu (215)	2006	Low	Cross sectional	Taiwan	Retrospective	Not clear	One site	2001-2005	Adults	75	42	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	0.95	General	NR	1795

Author	Year	Risk of bias	Design	Country	Timing	Sampling method	Sites	Period	Population	%Male	Mean or median age	%Obesity	Pattern of appendicitis	%Catarrhal	%Perforated	%Suppurated	%Gangrenous	% with antibiotic therapy	Type of surgery	Time to complete the surgery intervention (in hours)	Type of anesthesia	SSI Definition	Sample
Wu (216)	2011	Moderate	Cohort	Taiwan	Retrospective	Exhaustive	Multisite	2004-2009	Children, Adolescents, Adults, Elderly	58.1	36.4	NR	Unclear	NR	NR	NR	NR		Laparoscopy or Open Surgery	NR	General	NR	1366
Wu (217)	2014	Moderate	Clinical trial	China	Prospective	Random	One site	2011-2013	Children, Adolescents	60	8.5	NR	Catarrhal	100	0	0	0		Laparoscopy	1	General	NR	60
Wu (218)	2017	Moderate	Cross sectional	China	Retrospective	Consecutive	One site	2014-2016	Elderly	59	71	NR	Perforated, Suppurated, Gangrenous	0	61.7	10	28.7	100	Laparoscopy or Open Surgery	1	General	NR	115
Yaghoubian (219)	2010	High	Cross sectional	USA	Retrospective	Exhaustive	Multisite	1998-2007	Children, Adolescents, Adults, Elderly	61.5	29.2	NR	Catarrhal, Perforated	73.4	26.6	0	0		Laparoscopy or Open Surgery	NR	NR	NR	4325
Yagnik (220)	2010	Moderate	Cross sectional	India	Retrospective	Consecutive	One site	2007-2009	Children, Adolescents, Adults	32.5	23.41	NR	Catarrhal	100	0	0	0	100	Laparoscopy or Open Surgery	1	General	NR	151
Yousef (221)	2017	Moderate	Cohort	Canada	Prospective	Consecutive	One site	2015-2016	Children, Adolescents	63.1	9.3	NR	Perforated	0	100	0	0	100	Laparoscopy	NR	General	NR	122
Zhang (222)	2015	Moderate	Clinical trial	China	Prospective	Random	One site	2012-2013	Adults	47.2	30.8	NR	Unclear	10.2	7.4	54.6	9.3		Laparoscopy	0.9	General	NR	108

NR: not reported

References

References

1. Adejumo A.A., N.M., Mshelia, Y.M., Saleh. Clinicopathological presentation and management outcome of appendicitis in gombe, north-east nigeria: a 7-year retrospective audit. Niger J Med. 2015;24(4):337-43.
2. Aguiló J., S., Peiró, C., Muñoz, J., García del Caño, M., Garay, V., Viciano, et al. Adverse outcomes in the surgical treatment of acute appendicitis. Efectos adversos en la cirugía de la apendicitis aguda. 2005;78(5):312-7.
3. Adhikary S., S., Tyagi, G., Sapkota, A., Afaq, B.K., Bhattarai, C.S., Agrawal. Port exteriorization appendectomy: is it the future? Nepal Med Coll J. 2008;10(1):30-4.
4. Ahmad M., K., Ali, H., Latif, S., Naz, K., Said. Comparison of primary wound closure with delayed primary closure in perforated appendicitis. J Ayub Med Coll Abbottabad. 2014;26(2):153-7.
5. Ahmed I., J., Burr, M., Castillo, D., Collins, J.A., Cook, M., Campbell, et al. Single port/incision laparoscopic surgery compared with standard three-port laparoscopic surgery for appendectomy: A randomized controlled trial. Surg Endosc Interv Tech.

- 1
2
3 2015;29(1):77-85.
4
5
6
7 6. Akkoyun I., A., Taş Tuna. Advantages of abandoning abdominal cavity irrigation and
8 drainage in operations performed on children with perforated appendicitis. *J Pediatr Surg.*
9 2012;47(10):1886-90.
10
11
12
13
14 7. Al-Saadi A.S., A.H., Al-Wadan, S.A., Hamarnah, H., Amin. Is abandoning routine
15 peritoneal cultures during appendectomy justified? *Saudi Med J.* 2007;28(12):1827-9.
16
17
18
19
20 8. Al-Temimi M.H., M.A., Berglin, E.G., Kim, D.J., Tessier, S.D., Johna. Endostapler versus
21 Hem-O-Lok clip to secure the appendiceal stump and mesoappendix during laparoscopic
22 appendectomy. *Am J Surg.* 2017;214(6):1143-8.
23
24
25
26
27 9. Ali N., S., Aliyu. Appendicitis and its surgical management experience at the University of
28 Maiduguri Teaching Hospital Nigeria. *Niger J Med.* 2012;21(2):223-6.
29
30
31
32
33
34 10. Ali K., H., Latif, S., Ahmad. Frequency of wound infection in non-perforated appendicitis
35 with use of single dose preoperative antibiotics. *J Ayub Med Coll Abbottabad.*
36 2015;27(2):378-80.
37
38
39
40
41 11. Almström M., J.F., Svensson, B., Patkova, A., Svenningsson, T., Wester. In-hospital
42 surgical delay does not increase the risk for perforated appendicitis in children. *Ann Surg.*
43 2017;265(3):616-21.
44
45
46
47
48 12. Álvarez-Moreno C., A.M., Pérez-Fernández, V.D., Rosenthal, J., Quintero, E., Chapeta-
49 Parada, C., Linares, et al. Surgical site infection rates in 4 cities in Colombia: Findings of
50 the International Nosocomial Infection Control Consortium (INICC). *Am J Infect Control.*
51 2014;42(10):1089-92.
52
53
54
55
56
57 13. Andert A., H.P., Alizai, C.D., Klink, N., Neitzke, C., Fitzner, C., Heidenhain, et al. Risk
58 factors for morbidity after appendectomy. *Langenbeck's Arch Surg.* 2017;402(6):987-93.
59
60

- 1
2
3
4
5 14. Andersson R.E. Short-term complications and long-term morbidity of laparoscopic and
6
7
8
9
10
11 15. Aranda-Narváez J.M., A.J., González-Sánchez, N., Marín-Camero, C., Montiel-Casado, P.,
12
13 López-Ruiz, B., Sánchez-Pérez, et al. Conservative approach versus urgent appendectomy
14
15 in surgical management of acute appendicitis with abscess or phlegmon. Resultados del
16
17 tratamiento conservador inicial y de la cirugía urgente en la apenaicitis aguda evolucionada.
18
19 2010;102(11):648-52.
- 20
21
22 16. Aranda-Narváez J.M., T., Prieto-Puga Arjona, B., García-Albiach, M.C., Montiel-Casado,
23
24 A.J., González-Sánchez, B., Sánchez-Pérez, et al. Postappendectomy surgical site infection:
25
26 Overall rate and type according to open/laparoscopic approach. Infección de sitio quirúrgico
27
28 tras apendicectomía urgente: tasa global y tipo según la vía de abordaje
29
30 (abierta/laparoscópica). 2014;32(2):76-81.
- 31
32
33 17. Arthur T., R., Gartrell, B., Manoharan, D., Parker. Emergency appendicectomy in Australia:
34
35 findings from a multicentre, prospective study. ANZ J Surg. 2017;87(9):656-60.
- 36
37
38 18. Asefa Z. Acute appendicitis in Yirgalem Hospital, southern Ethiopia. Ethiop Med J.
39
40 2002;40(2):155-62.
- 41
42
43 19. Assefa Z., A., G/yesuse. Acute appendicitis in children admitted to zewditu memorial
44
45 hospital. Ethiop Med J. 2014;52(4):189-95.
- 46
47
48 20. Atif M.L., F., Sadaoui, A., Bezzaoucha, C.A., BezzaouchaKaddache, R., Boukari, S.,
49
50 Djelato, et al. Intra-abdominal abscesses and laparoscopic versus open appendectomies.
51
52 Infect Control Hosp Epidemiol. 2009;30(7):713-5
- 53
54
55
56 21. Bae E., A., Dehal, V., Franz, M., Joannides, N., Sakis, J., Scurlock, et al. Postoperative
57
58 antibiotic use and the incidence of intra-abdominal abscess in the setting of suppurative
59
60 appendicitis: a retrospective analysis. Am J Surg.2016;212(6):1121-5.

- 1
2
3
4
5
6 22. Bae S.U., W.K., Jeong, S.K., Baek. Single-port laparoscopic interval appendectomy for
7 perforated appendicitis with a periappendiceal abscess. *Ann Coloproctol.* 2016;32(3):105-
8 10.
9
10
11
12
13 23. Baek H.N., Y.H., Jung, Y.H., Hwang. Laparoscopic versus open appendectomy for
14 appendicitis in elderly patients. *J Korean Soc Coloproctology.* 2011;27(5):241-5.
15
16
17
18 24. Bali İ., F., Karateke, S., Özyazıcı, A., Kuvvetli, C., Oruç, E., Menekşe, et al. Comparison
19 of intracorporeal knotting and endoloop for stump closure in laparoscopic appendectomy.
20 *Laparoskopik appendektomide intrakorporal düğüm ve endoloop ile güdük kapama*
21 *yöntemlerinin karşılaştırılması.* 2015;21(6):446-9.
22
23
24
25
26
27 25. Bhangu A., Richardson, C., Torrance, A., Pinkney, T., Collaborative, Natl Surg Res.
28 Multicentre observational study of performance variation in provision and outcome of
29 emergency appendicectomy. *British Journal Of Surgery.* 2013;100(9):1240-52.
30
31
32
33
34
35 26. Bansal V., S., Altermatt, D., Nadal, C., Berger. Lack of benefit of preoperative antimicrobial
36 prophylaxis in children with acute appendicitis: A prospective cohort study. *Infection.*
37 2012;40(6):635-41.
38
39
40
41
42 27. Batajoo H., N.K., Hazra. Laparoscopic versus open appendectomy in acute appendicitis. *J*
43 *Nepal Health Res Counc.* 2012;10(22):239-42.
44
45
46
47
48 28. Saranga Bharathi R., V., Sharma, A., Chakladar, P., Kumari. Port exteriorisation
49 appendectomy-our experience. *Med J Armed Forces India.* 2011;67(2):147-51.
50
51
52
53
54 29. Biçakci U., B., Tander, M., Günaydin, R., Rizalar, E., Aritürk, S.H., Ayyildiz, et al. The
55 comparison of open and laparoscopic appendectomy: Is there any outcome difference
56 between non-complicated and complicated appendicitis? *Balkan Med J.* 2011;28(3):304-6.
57
58
59
60

- 1
2
3 30. Bickel A., M., Gurevits, R., Vamos, S., Ivry, A., Eitan. Perioperative hyperoxygenation and
4 wound site infection following surgery for acute appendicitis : A randomized, prospective,
5 controlled trial. *Arch Surg.* 2011;146(4):464-70.
6
7
8
9
10
11 31. Blackwood B.P., C.D., Gause, J.C., Harris, C.M., Theodorou, I., Helenowski, T.B., Lautz,
12 et al. Overweight and obese pediatric patients have an increased risk of developing a
13 surgical site infection. *Surg Infect.* 2017;18(4):491-7.
14
15
16
17
18 32. Blakely M.L., R., Williams, M.S., Dassinger, J.W., Eubanks III, P., Fischer, E.Y., Huang,
19 et al. Early vs interval appendectomy for children with perforated appendicitis. *Arch Surg.*
20 2011;146(6):660-5.
21
22
23
24
25
26 33. Bonadio W., K., Rebillot, O., Ukwuoma, C., Saracino, A., Iskhakov. Management of
27 Pediatric Perforated Appendicitis: Comparing Outcomes Using Early Appendectomy
28 versus Solely Medical Management. *Pediatr Infect Dis J.* 2017;36(10):937-41.
29
30
31
32
33 34. Boomer L.A., J.N., Cooper, K.J., Deans, P.C., Minneci, K., Leonhart, K.A., Diefenbach, et
34 al. Does delay in appendectomy affect surgical site infection in children with appendicitis?
35 *J Pediatr Surg.* 2014;49(6):1026-9.
36
37
38
39
40
41 35. Boomer L.A., J.N., Cooper, S., Anandalwar, S.C., Fallon, D., Ostlie, C.M., Leys, et al.
42 Delaying appendectomy does not lead to higher rates of surgical site infections. *Ann Surg.*
43 2016;264(1):164-8.
44
45
46
47
48 36. Bozkurt M.A., M.G., Ünsal, S., Kapan, B., Kankaya, M.U., Kalaycii, H., Aliiş. Two
49 different methods for appendiceal stump closure: Metal clip and Hem-o-lok clip. *J*
50 *Laparoendosc Adv Surg Techn.* 2014;24(8):571-3.
51
52
53
54
55
56 37. Brandt C., U., Hott, D., Sohr, F., Daschner, P., Gastmeier, H., Rüden. Operating room
57 ventilation with laminar airflow shows no protective effect on the surgical site infection rate
58 in orthopedic and abdominal surgery. *Ann Surg.* 2008;248(5):695-700.
59
60

- 1
2
3
4
5
6 38. Cairo S.B., M.V., Raval, M., Browne, H., Meyers, D.H., Rothstein. Association of same-
7 day discharge with hospital readmission after appendectomy in pediatric patients. *JAMA*
8 *Surg.* 2017;152(12):1106-12.
9
10
11
12
13 39. Cameron D.B., P., Melvin, D.A., Graham, C.C., Glass, S.K., Serres, M.P., Kronman, et al.
14 Extended Versus Narrow-spectrum Antibiotics in the Management of Uncomplicated
15 Appendicitis in Children: A Propensity-matched Comparative Effectiveness Study. *Ann*
16 *Surg.* 2017.
17
18
19
20
21
22 40. Cao J.-G., F., Tao, X.-J., Zhou, X.-G., Wang, S.-S., Wang, H., Zhang, et al. Trends and
23 outcomes of laparoscopic appendectomy in China: A multicenter, retrospective cohort
24 study. *Surg Pract.* 2015;19(4):166-72.
25
26
27
28
29
30 41. Cervantes-Sánchez C.R., R., Gutiérrez-Vega, J.A., Vázquez-Carpizo, P., Clark, C., Athié-
31 Gutiérrez. Syringe pressure irrigation of subdermic tissue after appendectomy to decrease
32 the incidence of postoperative wound infection. *World J Surg.* 2000;24(1):38-42.
33
34
35
36
37 42. Chamisa I. A clinicopathological review of 324 appendices removed for acute appendicitis
38 in Durban, South Africa: a retrospective analysis. *Ann R Coll Surg Engl.* 2009;91(8):688-
39 92.
40
41
42
43
44
45 43. Chaudhary I.A., Samiullah, A.A., Mallhi, Z., Afridi, A., Bano. Is it necessary to invaginate
46 the stump after appendectomy? *Pak J Med Sci.* 2005;21(1):35-8.
47
48
49
50
51 44. Chen D., H., Shi, H., Dong, K., Liu, K., Ding. Gasless single-incision laparoscopic
52 appendectomy. *Surg Endosc Interv Tech.* 2011;25(5):1472-6.
53
54
55
56 45. Chen C.-Y., Y.-C., Chen, H.-N., Pu, C.-H., Tsai, W.-T., Chen, C.-H., Lin. Bacteriology of
57 acute appendicitis and its implication for the use of prophylactic antibiotics. *Surg Infect.*
58 2012;13(6):383-90.
59
60

- 1
2
3
4
5
6 46. Chen C.-C., C.-T., Ting, M.-J., Tsai, W.-C., Hsu, P.-C., Chen, M.-D., Lee, et al.
7 Appendectomy timing: Will delayed surgery increase the complications? *J Chin Med*
8 *Assoc.* 2015;78(7):395-9.
9
10
11
12 47. Chiang R.-A., S.-L., Chen, Y.-C., Tsai, M.-J., Bair. Comparison of primary wound closure
13 versus open wound management in perforated appendicitis. *J Formos Med Assoc.*
14 2006;105(10):791-5.
15
16
17
18
19 48. Chiang R.-A., S.-L., Chen, Y.-C., Tsai. Delayed primary closure versus primary closure for
20 wound management in perforated appendicitis: A prospective randomized controlled trial.
21 *J Chin Med Assoc.* 2012;75(4):156-9.
22
23
24
25
26
27 49. Cho M., J., Kang, I.-K., Kim, K.Y., Lee, S.-K., Sohn. Underweight body mass index as a
28 predictive factor for surgical site infections after laparoscopic appendectomy. *Yonsei Med*
29 *J.* 2014;55(6):1611-6.
30
31
32
33
34
35 50. Choudhary S.K., S.K., Dhakaita. Appendicular mass-early appendicectomy vs interval
36 appendicectomy. *Intl J Pharma Bio Sci.* 2014;5(1):B400-B4.
37
38
39
40
41 51. Clyde C., T., Bax, A., Merg, M., MacFarlane, P., Lin, S., Beyersdorf, et al. Timing of
42 intervention does not affect outcome in acute appendicitis in a large community practice.
43 *Am J Surg.* 2008;195(5):590-3.
44
45
46
47
48 52. Coakley B.A., E.S., Sussman, T.S., Wolfson, A.S., Bhagavath, J.J., Choi, N.E., Ranasinghe,
49 et al. Postoperative antibiotics correlate with worse outcomes after appendectomy for
50 nonperforated appendicitis. *J Am Coll Surg.* 2011;213(6):778-83.
51
52
53
54
55 53. Crandall M., M.B., Shapiro, M., Worley, M.A., West. Acute uncomplicated appendicitis:
56 case time of day influences hospital length of stay. *Surg Infect (Larchmt).* 2009;10(1):65-
57 9.
58
59
60

- 1
2
3
4
5
6 54. Dede K., T., Mersich, A., Zaránd, I., Besznyák, Z., Baranyai, B., Atkári, et al. Laparoscopic
7 or open appendectomy? *Laparoszkópos vagy nyílt appendectomia?* 2008;149(50):2357-61.
8
9
10
11 55. Dhiman N., A., Chi, T.M., Pawlik, D.T., Efron, E.R., Haut, E.B., Schneider, et al. Increased
12 complications after appendectomy in patients with cerebral palsy: Are special needs patients
13 at risk for disparities in outcomes? *Surgery*. 2013;154(3):479-85.
14
15
16
17 56. Dimitriou I., B., Reckmann, O., Nephuth, M., Betzler. Single institution's experience in
18 laparoscopic appendectomy as a suitable therapy for complicated appendicitis.
19 *Langenbeck's Arch Surg*. 2013;398(1):147-52.
20
21
22
23
24
25
26 57. Durkin M.J., K.V., Dicks, A.W., Baker, S.S., Lewis, R.W., Moehring, L.F., Chen, et al.
27 Seasonal variation of common surgical site infections: Does season matter? *Infect Control*
28 *Hosp Epidemiol*. 2015;36(9):1011-6.
29
30
31
32
33
34 58. Ein S.H., A., Nasr, A., Ein. Open appendectomy for pediatric ruptured appendicitis: a
35 historical clinical review of the prophylaxis of wound infection and postoperative intra-
36 abdominal abscess. *Can J Surg*. 2013;56(3):E7-E12.
37
38
39
40
41 59. Fukuda H. Patient-related risk factors for surgical site infection following eight types of
42 gastrointestinal surgery. *J Hosp Infect*. 2016;93(4):347-54.
43
44
45
46
47 60. Gandaglia G., K.R., Ghani, A., Sood, J.R., Meyers, J.D., Sammon, M., Schmid, et al. Effect
48 of minimally invasive surgery on the risk for surgical site infections results from the national
49 surgical quality improvement program (nsqip) database. *JAMA Surg*. 2014;149(10):1039-
50 44.
51
52
53
54
55
56 61. Garcell H.G., A.V., Arias, C.A., Pancorbo Sandoval, E.G., García, M.E., Valle Gamboa,
57 A.B., Sado, et al. Incidence and etiology of surgical site infections in appendectomies: A 3-
58 year prospective study. *Oman Med J*. 2017;32(1):31-5.
59
60

- 1
2
3
4
5
6 62. Ghnam W.M. Elderly versus young patients with appendicitis 3 years experience. *Alex J*
7 *Med.* 2012;48(1):9-12.
8
9
10
11
12 63. Giesen L.J., A.L., van den Boom, C.C., van Rossem, P.T., den Hoed, B.P., Wijnhoven.
13 Retrospective Multicenter Study on Risk Factors for Surgical Site Infections after
14 Appendectomy for Acute Appendicitis. *Dig Surg.* 2017;34(2):103-7.
15
16
17
18
19 64. Giiti G.C., H.D., Mazigo, J., Heukelbach, W., Mahalu. HIV, appendectomy and
20 postoperative complications at a reference hospital in Northwest Tanzania: Cross-sectional
21 study. *AIDS Res Ther.* 2010;7.
22
23
24
25
26
27 65. Golub A.V., R.S., Kozlov, V.G., Pleshkov, A.P., Moskalev, R.A., Alibegov, M.A.,
28 Chelombitko. Surgical Site Infections after Open Appendectomy and Effectiveness of
29 Complex Approach to Their Prevention. *Khirurgiia (Mosk).* 2016(6):68-76.
30
31
32
33
34
35 66. Gross T.S., C., McCracken, K.F., Heiss, M.L., Wulkan, M.V., Raval. The contribution of
36 practice variation to length of stay for children with perforated appendicitis. *J Pediatr Surg.*
37 2016;51(8):1292-7.
38
39
40
41
42 67. Gurien L.A., D.L., Wyrick, S.D., Smith, M.S., Dassinger. Optimal timing of appendectomy
43 in the pediatric population. *J Surg Res.* 2016;202(1):126-31.
44
45
46
47
48 68. Hamzaoglu I., B., Baca, D.E., Böler, E., Polat, Y., Özer. Is umbilical flora responsible for
49 wound infection after laparoscopic surgery? *Surg Laparoscopy Endosc Percutaneous Tech.*
50 2004;14(5):263-7.
51
52
53
54
55 69. Harmon L.A., M.L., Davis, D.C., Jupiter, R.C., Frazee, J.L., Regner. Computed tomography
56 to operating room in less than 3 hours minimizes complications from appendicitis. *Am J*
57 *Surg.* 2016;212(2):246-50.
58
59
60

- 1
2
3
4
5
6 70. Helling T.S., D.F., Soltys, S., Seals. Operative versus non-operative management in the care
7 of patients with complicated appendicitis. *Am J Surg.* 2017;214(6):1195-200.
8
9
10
11 71. Helmer K.S., E.K., Robinson, K.P., Lally, J.C., Vasquez, K.L., Kwong, T.H., Liu, et al.
12 Standardized patient care guidelines reduce infectious morbidity in appendectomy patients.
13 *Am J Surg.* 2002;183(6):608-13.
14
15
16
17
18 72. Hesami M.A., H., Alipour, H., Nikoupour Daylami, B., Alipour, S., Bazargan-Hejazi, A.,
19 Ahmadi. Irrigation of abdomen with imipenem solution decreases surgical site infections in
20 patients with perforated appendicitis: A randomized clinical trial. *Iran Red Crescent MedJ.*
21 2014;16(4).
22
23
24
25
26
27
28 73. Horvath P., J., Lange, R., Bachmann, F., Struller, A., Königsrainer, M., Zdichavsky.
29 Comparison of clinical outcome of laparoscopic versus open appendectomy for complicated
30 appendicitis. *Surg Endosc Interv Tech.* 2017;31(1):199-205.
31
32
33
34
35
36 74. Hughes M.J., E., Harrison, S., Paterson-Brown. Post-operative antibiotics after
37 appendectomy and post-operative abscess development: A retrospective analysis. *Surg*
38 *Infect.* 2013;14(1):56-61.
39
40
41
42
43 75. Hussain M.I., M.K., Alam, H.H., Al-Qahatani, M.H., Al-Akeely. Role of postoperative
44 antibiotics after appendectomy in non-perforated appendicitis. *J Coll Phys Surg Pak.*
45 2012;22(12):756-9.
46
47
48
49
50
51 76. Iqbal M., M., Jawaid, A., Qureshi, S., Iqbal. Effect of povidone-iodine irrigation on post
52 appendectomy wound infection: Randomized control trial. *J Postgrad Med Inst.*
53 2015;29(3):160-4.
54
55
56
57
58 77. Iñigo J.J., B., Bermejo, B., Oronoz, J., Herrera, A., Tarifa, F., Pérez, et al. Surgical site
59 infection in general surgery: 5-year analysis and assessment of the National Nosocomial
60

- 1
2
3 Infection Surveillance (NNIS) index. Infección de sitio quirúrgico en un servicio de cirugía
4 general Análisis de cinco años y valoración del índice National Nosocomial Infection
5 Surveillance (NNIS). 2006;79(4):224-30.
6
7
8
9
10
11 78. Javadi S.M.R., S.Y., Zarghami, P., Ghaderzadeh, M., Ghorbanpoor, H.R., Makarchian, A.,
12 Derakhshanfar, et al. Comparison of small access and classic McBurney's incisions for open
13 appendectomy: A randomized controlled trial. Shiraz E Med J. 2017;18(10).
14
15
16
17
18 79. Jenkins P.C., M.K., Oerline, A.J., Mullard, M.J., Englesbe, D.A., Campbell, M.R.,
19 Hemmila. Hospital variation in outcomes following appendectomy in a regional quality
20 improvement program. Am J Surg. 2016;212(5):857-62.
21
22
23
24
25
26 80. Kang J., B.N., Bae, G., Gwak, I., Park, H., Cho, K., Yang, et al. Comparative study of a
27 single-incision laparoscopic and a conventional laparoscopic appendectomy for the
28 treatment of acute appendicitis. J Korean Soc Coloproctology. 2012;28(6):304-8.
29
30
31
32
33
34 81. Kapischke M., A., Pries, A., Caliebe. Short term and long term results after open vs.
35 Laparoscopic appendectomy in childhood and adolescence: A subgroup analysis. BMC
36 Pediatr. 2013;13(1).
37
38
39
40
41 82. Karam P.A., A., Hiuser, D., Magnuson, F.G.F., Seifarth. Intracorporeal hybrid single port
42 vs conventional laparoscopic appendectomy in children. Pediatr Med Chir. 2016;38(3):89-
43 92.
44
45
46
47
48
49 83. Karam P.A., A., Mohan, M.R., Buta, F.G., Seifarth. Comparison of Transumbilical
50 Laparoscopically Assisted Appendectomy to Conventional Laparoscopic Appendectomy in
51 Children. Surg Laparoscopy Endosc Percutaneous Tech. 2016;26(6):508-12.
52
53
54
55
56 84. Kasatpibal N., S., Jamulitrat, V., Chongsuvivatwong. Standardized incidence rates of
57 surgical site infection: A multicenter study in Thailand. Am J Infect Control.
58 2005;33(10):587-94.
59
60

- 1
2
3
4
5
6 85. Kasatpibal N., M., Nørgaard, H.T., Sørensen, H.C., Schönheyder, S., Jamulitrat, V.,
7 Chongsuvivatwong. Risk of surgical site infection and efficacy of antibiotic prophylaxis: A
8 cohort study of appendectomy patients in Thailand. *BMC Infect Dis.* 2006;6.
9
10
11
12
13 86. Kato Y., T., Marusasa, S., Ichikawa, G.J., Lane, T., Okazaki, A., Yamataka. Lapprotector
14 use decreases incisional wound infections in cases of perforated appendicitis: a prospective
15 study. *Asian J Surg.* 2008;31(3):101-3.
16
17
18
19
20 87. Kell M.R., K., Power, D.C., Winter, C., Power, C., Shields, W.O., Kirwan, et al. Predicting
21 outcome after appendectomy. *Ir J Med Sci.* 2003;172(2):63-5.
22
23
24
25
26 88. Khan M.N., T., Fayyad, T.D., Cecil, B.J., Moran. Laparoscopic versus open appendectomy:
27 the risk of postoperative infectious complications. *JLS.* 2007;11(3):363-7.
28
29
30
31
32 89. Khan K.I., S., Mahmood, M., Akmal, A., Waqas. Comparison of rate of surgical wound
33 infection, length of hospital stay and patient convenience in complicated appendicitis
34 between primary closure and delayed primary closure. *J Pak Med Assoc.* 2012;62(8):596-
35 8.
36
37
38
39
40
41
42 90. Khan I., M.I., Khan, M., Jawed, U., Shaikh, S., Ahmed, A., Arif. To compare the frequency
43 of superficial surgical site infection after laparoscopic versus open appendectomy. *Med*
44 *Forum Monthly.* 2014;25(11):52-5.
45
46
47
48
49 91. Khiria L.S., R., Ardhari, N., Mohan, P., Kumar, R., Nambiar. Laparoscopic
50 appendectomy for complicated appendicitis: Is it safe and justified? A retrospective
51 analysis. *Surg Laparoscopy Endosc Percutaneous Tech.* 2011;21(3):142-5.
52
53
54
55
56 92. Kılıç Ş.S., S., Ekinçi, İ., Karnak, A.Ö., Çiftçi, F.C., Tanyel, M.E., Şenocak. Drainage
57 systems' effect on surgical site infection in children with perforated appendicitis. *Drenaj*
58 *Sistemlerinin perfore apandisitli çocuklarda cerrahi alan enfeksiyonuna etkisi.*
59
60

2016;7(5):591-4.

93. Kim M.J., F.J., Fleming, D.D., Gunzler, S., Messing, R.M., Salloum, J.R.T., Monson. Laparoscopic appendectomy is safe and efficacious for the elderly: An analysis using the National Surgical Quality Improvement Project database. *Surg Endosc Interv Tech.* 2011;25(6):1802-7.
94. Kim J.H., H.Y., Kim, S.K., Park, J.S., Lee, D.S., Heo, S.W., Park, et al. Single-incision laparoscopic appendectomy versus conventional laparoscopic appendectomy: Experiences from 1208 cases of single-incision laparoscopic appendectomy. *Ann Surg.* 2015;262(6):1054-8.
95. Kim J.K., J., Kang, W.R., Kim, E.J., Park, S.H., Baik, K.Y., Lee. Does Conversion Adversely Impact the Clinical Outcomes for Patients with Complicated Appendicitis? *J Laparoendosc Adv Surg Techn.* 2016;26(8):635-40.
96. Kiriakopoulos A., D., Tsakayannis, D., Linos. Laparoscopic management of complicated appendicitis. *JLS.* 2006;10(4):453-6.
97. Kirshtein B., Z.H., Perry, S., Mizrahi, L., Lantsberg. Value of laparoscopic appendectomy in the elderly patient. *World J Surg.* 2009;33(5):918-22.
98. Kiudelis M., P., Ignatavicius, K., Zviniene, S., Grizas. Analysis of intracorporeal knotting with invaginating suture versus endoloops in appendiceal stump closure. *Wideochir Inne Tech Ma?oinwazyjne.* 2013;8(1):69-73.
99. Kleif J., L., Rasmussen, S., Fonnes, P., Tibæk, A., Daoud, H., Lund, et al. Enteral Antibiotics are Non-inferior to Intravenous Antibiotics After Complicated Appendicitis in Adults: A Retrospective Multicentre Non-inferiority Study. *World J Surg.* 2017;41(11):2706-14.

- 1
2
3 100. Koizumi N., H., Kobayashi, Y., Nakase, T., Takagi, K., Fukumoto. Efficacy of
4 transumbilical laparoscopic-assisted appendectomy for appendicitis: a four-year experience
5 at a single center. *Surg Today*. 2015;45(10):1245-9.
6
7
8
9
10
11 101. Shimizu T., M., Ishizuka, K., Kubota. The preoperative serum C-reactive protein level
12 is a useful predictor of surgical site infections in patients undergoing appendectomy. *Surg*
13 *Today*. 2015;45(11):1404-10.
14
15
16
17
18 102. Kumamoto K., H., Imaizumi, N., Hokama, T., Ishiguro, K., Ishibashi, K., Baba, et al.
19 Recent trend of acute appendicitis during pregnancy. *Surg Today*. 2015;45(12):1521-6.
20
21
22
23
24 103. Kumar B., A., Samad, T.W., Khanzada, M.H., Laghari, A.R., Shaikh. Superiority of
25 laparoscopic appendectomy over open appendectomy: The Hyderabad experience. *Rawal*
26 *Med J*. 2008;33(2):165-8.
27
28
29
30
31
32 104. Kumar S., A., Jalan, B.N., Patowary, S., Shrestha. Laparoscopic appendectomy versus
33 open appendectomy for acute appendicitis: A prospective comparative study. *Kathmandu*
34 *Univ Med J*. 2016;14(55):244-8.
35
36
37
38
39 105. Lacher M., O.J., Muensterer, G.R., Yannam, C.J., Aprahamian, L., Perger, M., Megison,
40 et al. Feasibility of single-incision pediatric endosurgery for treatment of appendicitis in
41 415 children. *J Laparoendosc Adv Surg Techn*. 2012;22(6):604-8.
42
43
44
45
46 106. Lasses-Martínez B., E., Ortiz-Oshiro, J.L., Cabañas-Ojeda, P., Benito-Expósito, C.,
47 Fernández-Pérez, J., Alvarez Fernández-Represa. Cost is not a drawback to perform
48 laparoscopic appendectomy in an academic hospital. *Surg Laparoscopy Endosc*
49 *Percutaneous Tech*. 2014;24(4):e123-e7.
50
51
52
53
54
55
56 107. Le D., W., Rusin, B., Hill, J., Langell. Post-operative antibiotic use in nonperforated
57 appendicitis. *Am J Surg*. 2009;198(6):748-52.
58
59
60

- 1
2
3
4
5 108. Lee P., K., Waxman, B., Taylor, S., Yim. Use of wound-protection system and
6 postoperative wound-infection rates in open appendectomy: A randomized prospective
7 trial. *Arch Surg.* 2009;144(9):872-5.
8
9
10
11
12 109. Lee S.L., S., Shekherdimian, V.Y., Chiu. Comparison of pediatric appendicitis
13 outcomes between teaching and nonteaching hospitals. *J Pediatr Surg.* 2010;45(5):894-7.
14
15
16
17 110. Lee J.A., K.Y., Sung, J.H., Lee, D.S., Lee. Laparoscopic appendectomy with a single
18 incision in a single institute. *J Korean Soc Coloproctology.* 2010;26(4):260-4.
19
20
21
22
23 111. Lee S.-Y., H.-M., Lee, C.-S., Hsieh, J.-H., Chuang. Transumbilical laparoscopic
24 appendectomy for acute appendicitis: A reliable one-port procedure. *Surg Endosc Interv*
25 *Tech.* 2011;25(4):1115-20.
26
27
28
29
30 112. Lee S.L., A., Yaghoubian, A., Kaji. Laparoscopic vs open appendectomy in children:
31 Outcomes comparison based on age, sex, and perforation status. *Arch Surg.*
32 2011;146(10):1118-21.
33
34
35
36
37
38
39 113. Lee S.M., G.S., Hwang, D.S., Lee. Single-incision laparoscopic appendectomy using
40 homemade glove port at low cost. *J Minimal Access Surg.* 2016;12(2):124-8.
41
42
43
44
45 114. Levy S.M., G., Holzmann-Pazgal, K.P., Lally, K., Davis, L.S., Kao, K., Tsao. Quality
46 check of a quality measure: Surgical wound classification discrepancies impact risk-
47 stratified surgical site infection rates in pediatric appendicitis. *J Am Coll Surg.*
48 2013;217(6):969-73.
49
50
51
52
53 115. Li P., Q., Xu, Z., Ji, Y., Gao, X., Zhang, Y., Duan, et al. Comparison of surgical stress
54 between laparoscopic and open appendectomy in children. *J Pediatr Surg.* 2005;40(8):1279-
55 83.
56
57
58
59
60

- 1
2
3 116. Lim S.G., E.J., Ahn, S.Y., Kim, I.Y., Chung, J.-M., Park, S.H., Park, et al. A clinical
4 comparison of laparoscopic versus open appendectomy for complicated appendicitis. J
5 Korean Soc Coloproctology. 2011;27(6):293-7.
6
7
8
9
10
11 117. Geraldo José de Souza Lima, Silva, Alcino Lázaro da, Castro, Eduardo Godoy, Abras,
12 Gustavo Munayer, Pires, Lívio José Suretti, Leite, Rodrigo Fabiano Guedes. Efetividade e
13 segurança da apendicectomia videoassistida em porta única transumbilical em adolescentes
14 e adultos X1 Effectiveness and safeness of single-port trans-umbilical laparoscopic
15 appendectomy done in adolescents and adults. Revista do Colégio Brasileiro de Cirurgiões.
16 2008;35(4):244-51.
17
18
19
20
21
22
23 118. Lin H.-F., J.-M., Wu, L.-M., Tseng, K.-H., Chen, S.-H., Huang, I.-R., Lai. Laparoscopic
24 Versus Open Appendectomy for Perforated Appendicitis. J Gastrointest Surg.
25 2006;10(6):906-10.
26
27
28
29
30
31 119. Litz C.N., S.M., Farach, P.D., Danielson, N.M., Chandler. Obesity and single-incision
32 laparoscopic appendectomy in children. J Surg Res. 2016;203(2):283-6.
33
34
35
36
37 120. Liu C., W., Wang, Y., Sun, M., Xu, H., Zhuang, H., Chen, et al. Efficacy and
38 complications of laparoscopic appendectomy for pediatric appendicitis. Int J Clin Exp Med.
39 2017;10(9):13784-9.
40
41
42
43
44 121. Mahmood M.M., A., Shahab, M.A., Razzaq. Surgical site infection in open versus
45 laparoscopic appendectomy. Pak J Med Health Sci. 2016;10(3):1076-8.
46
47
48
49
50 122. Mbah N., W.Ek., Opara, N.P., Agwu. Waiting time among acute abdominal
51 emergencies in a Nigerian teaching hospital: Causes of delay and consequences. Niger J
52 Surg Res. 2006;8(1):69-73.
53
54
55
56
57 123. Memon G.A., A.I., Memon, S.K.A., Shah, R.A., Sahito, Habib-Ur-Rehman, S., Leghari,
58 et al. An experience of treatment outcome in acute appendicitis with antibiotics and
59
60

- 1
2
3 appendectomy at a tertiary care hospital. *Med Forum Monthly*. 2017;28(3):136-40.
4
5
6
7 124. Menezes M., L., Das, M., Alagtal, J., Haroun, P., Puri. Laparoscopic appendectomy is
8 recommended for the treatment of complicated appendicitis in children. *Pediatr Surg Int*.
9 2008;24(3):303-5.
10
11
12
13
14 125. Merenda M., A., Litarski, P., Kabziński, D., Janczak. Laparoscopic appendectomy as
15 an alternative to conventional procedure - results in our own material. *Pol Przegl Chir*.
16 2013;85(6):323-8.
17
18
19
20
21 126. Michailidou M., M.G., Sacco Casamassima, S.D., Goldstein, C., Gause, O., Karim, J.H.,
22 Salazar, et al. The impact of obesity on laparoscopic appendectomy: Results from the ACS
23 National Surgical Quality Improvement Program pediatric database. *J Pediatr Surg*.
24 2015;50(11):1880-4.
25
26
27
28
29
30 127. Michailidou M., S.D., Goldstein, M.G., Sacco Casamassima, J.H., Salazar, R., Elliott,
31 J., Hundt, et al. Laparoscopic versus open appendectomy in children: The effect of surgical
32 technique on healthcare costs. *Am J Surg*. 2015;210(2):270-5.
33
34
35
36
37
38 128. Mickovic I.N., Z., Golubovic, S., Mickovic, D., Vukovic, S., Trajkovic, S.S.,
39 Antunovic, et al. A comparative analysis of laparoscopic appendectomy in relation to the
40 open appendectomy in children. *Uporedna analiza laparoscopske apendektomije u odnosu*
41 *na otvorenu apendektomiju kod dece*. 2016;17(1):49-53.
42
43
44
45
46
47 129. Ming P.C., T.Y., Yee Yan, L.H., Tat. Risk factors of postoperative infections in adults
48 with complicated appendicitis. *Surg Laparoscopy Endosc Percutaneous Tech*.
49 2009;19(3):244-8.
50
51
52
53
54
55 130. Mingmalairak C., P., Ungbhakorn, V., Paocharoen. Efficacy of antimicrobial coating
56 suture coated polyglactin 910 with tricosan (Vicryl Plus) compared with polyglactin 910
57 (Vicryl) in reduced surgical site infection of appendicitis, double blind randomized control
58
59
60

- 1
2
3 trial, preliminary safety report. *J Med Assoc Thailand*. 2009;92(6):770-5.
4
5
6
7
8 131. Miyano G., T., Okazaki, Y., Kato, T., Marusasa, T., Takahashi, G.J., Lane, et al. Open
9 versus laparoscopic treatment for pan-peritonitis secondary to perforated appendicitis in
10 children: A prospective analysis. *J Laparoendosc Adv Surg Techn*. 2010;20(7):655-7.
11
12
13
14
15 132. Moazzez A., R.J., Mason, N., Katkhouda. Thirty-day outcomes of laparoscopic versus
16 open appendectomy in elderly using ACS/NSQIP database. *Surg Endosc Interv Tech*.
17 2013;27(4):1061-71.
18
19
20
21
22 133. Mohammad Taghi Rajabi-Mashhadi, Mousavi, Seyed Hadi, Khosravi-Mashizi, M. H.,
23 Ghayour-Mobarhan, Majid, Sahebkar, Amirhossein. Optimum duration of perioperative
24 antibiotic therapy in patients with acute non-perforated appendicitis: a prospective
25 randomized trial. *Asian Biomedicine*. 2012;6(6):891-4.
26
27
28
29
30
31 134. Monge Jodra V., A., Robustillo Rodela, F., Martin Martinez, N., López Fresneña, S.,
32 Oña Compán, F., Calbo Torrecillas, et al. Standardized infection ratios for three general
33 surgery procedures: A comparison between Spanish hospitals and U.S. centers participating
34 in the national nosocomial infections surveillance system. *Infect Control Hosp Epidemiol*.
35 2003;24(10):744-8.
36
37
38
39
40
41 135. Mueck K.M., L.R., Putnam, K.T., Anderson, K.P., Lally, K., Tsao, L.S., Kao. Does
42 compliance with antibiotic prophylaxis in pediatric simple appendicitis matter? *J Surg Res*.
43 2017;216:1-8.
44
45
46
47
48
49
50 136. Muensterer O.J., C., Puga Nougues, O.O., Adibe, S.R., Amin, K.E., Georgeson, C.M.,
51 Harmon. Appendectomy using single-incision pediatric endosurgery for acute and
52 perforated appendicitis. *Surg Endosc Interv Tech*. 2010;24(12):3201-4.
53
54
55
56
57 137. Muensterer O.J., R., Keijzer. A simple vacuum dressing reduces the wound infection
58 rate of single-incision pediatric endosurgical appendectomy. *J Soc Laparoendoscopic Surg*.
59
60

- 2011;15(2):147-50.
138. Mustafa M.I.T., S.M., Chaudhry, R.I.T., Mustafa. Comparison of early outcome between patients of open appendectomy with and without drain for perforated appendicitis. *Pak J Med Health Sci.* 2016;10(3):890-3.
139. Nadler E.P., K.K., Reblock, H.R., Ford, B.A., Gaines. Monotherapy versus multi-drug therapy for the treatment of perforated appendicitis in children. *Surg Infect.* 2003;4(4):327-33.
140. Nataraja R.M., A., Bandi, S.A., Clarke, M.J., Haddad. Comparison of intra-abdominal abscess formation following laparoscopic and open appendectomy in children. *J Laparoendosc Adv Surg Techn.* 2010;20(4):391-4.
141. Nataraja R.M., W.J., Teague, J., Galea, L., Moore, M.J., Haddad, T., Tsang, et al. Comparison of intraabdominal abscess formation after laparoscopic and open appendectomies in children. *J Pediatr Surg.* 2012;47(2):317-21.
142. Norton Pérez, Romero, Marcela, Castelblanco, María Isabel, Rodríguez, Emma Isabel. Infección del sitio operatorio de apendicectomías en un hospital de la orinoquia colombiana X1 Surgical site infection following appendectomy at a hospital in the Colombian Orinoco river basin (Colombian Orinoquia). *Revista Colombiana de Cirugía.* 2009;24(1):23-30.
143. Obayashi J., K., Ohyama, S., Manabe, K., Tanaka, H., Nagae, H., Shima, et al. Are there reliable indicators predicting post-operative complications in acute appendicitis? *Pediatr Surg Int.* 2015;31(12):1189-93.
144. Obinwa O., C., Peirce, M., Cassidy, T., Fahey, J., Flynn. A model predicting perforation and complications in paediatric appendectomy. *Int J Colorectal Dis.* 2015;30(4):559-65.
145. Ohene-Yeboah M., B., Togbe. An audit of appendicitis and appendectomy in Kumasi,

- 1
2
3 Ghana. West Afr J Med. 2006;25(2):138-43.
4
5
6
7
8 146. Okkyung Suh, 신완균, 강성희, 양대현. Appropriate Duration of Prophylactic
9
10 Antibiotics in Acute Nonperforated Appendicitis Z1 급성 비천공성 충수염 수술시
11
12 예방적 항균제의 사용기간. Korean Journal of Clinical Pharmacy S1 한국임상약학회지.
13
14 2002;12(2):65-70.
15
16
17
18
19
20
21 147. Francisco Gabriel Onieva, Roldán, Sara, Domínguez, José Ramón, Montero, Juan
22 Pedro, Galnares, Alfonso, Peralta, Jordi. Abordaje laparoscópico frente a enfoque clásico
23 en el tratamiento de la apendicitis aguda X1 Laparoscopic approach versus classic open
24 procedure in the treatment of acute appendicitis. Revista Colombiana de Cirugía.
25 2017;32(1):26-31.
26
27
28
29
30
31
32 148. Page A.J., J.D., Pollock, S., Perez, S.S., Davis, E., Lin, J.F., Sweeney. Laparoscopic
33 Versus Open Appendectomy: An Analysis of Outcomes in 17,199 Patients Using
34 ACS/NSQIP. J Gastrointest Surg. 2010;14(12):1955-62.
35
36
37
38
39
40 149. Palesty J.A., X.J., Wang, R.C., Rutland, J., Leighton, S.J., Dudrick, A., Benbrahim.
41 Fifty-five consecutive laparoscopic appendectomy procedures without conversion. JSLS.
42 2004;8(2):141-5.
43
44
45
46
47 150. Pandit R.K. Safe and feasible time limit for early appendectomy in appendiceal mass.
48 Kathmandu Univ Med J. 2016;14(55):210-4.
49
50
51
52
53 151. Parcels J.P., J.P., Mileski, F.T., Gnagy, A.F., Haragan, W.J., Mileski. Using
54 antimicrobial solution for irrigation in appendicitis to lower surgical site infection rates. Am
55 J Surg. 2009;198(6):875-80.
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
152. Park H.-C., M.J., Kim, B.H., Lee. Effect of a Standardized Protocol of Antibiotic Therapy on Surgical Site Infection after Laparoscopic Surgery for Complicated Appendicitis. *Surg Infect.* 2017;18(6):684-8.
153. Seongmun Park, Park, Min-Su, Lee, Kil-Yeon. Relationship between the Hospital Visit-to-Operation Time Interval and the Risk of Appendiceal Perforation and Clinical Outcomes. *Journal of Minimally Invasive Surgery.* 2018;21(1):31-7.
154. Reoyo Pascual J.F., R., León Miranda, C., Cartón Hernández, E., Alonso Alonso, R.M., Martínez Castro, J., Sánchez Manuel. Laparoscopic appendectomy by 'glove port' system: Our first 100 cases. *Apendicectomía laparoscópica por sistema «glove port»: nuestros primeros 100 casos.* 2017;69(6):467-71.
155. Patrice Lemieux., Pascal Rheaume., Isabelle Levesque., Emmanuel Bujold., Gaetan Brochu. Laparoscopic appendectomy in pregnant patients: a review of 45 cases. *Surg Endosc.*2009; 23:1701.
156. Patel S.C., G.F., Jumba, S., Akmal. Laparoscopic appendectomy at the Aga Khan Hospital, Nairobi. *East Afr Med J.* 2003;80(9):447-51.
157. Percy C., K., Almahmoud, T., Jackson, C., Hartline, A., Cahill, L., Spence, et al. Risky business? Investigating outcomes of patients undergoing urgent laparoscopic appendectomy on antithrombotic therapy. *Am J Surg.* 2017;214(6):1012-5.
158. Pishori T., A.R., Siddiqui, M., Ahmed. Surgical wound infection surveillance in general surgery procedures at a teaching hospital in Pakistan. *Am J Infect Control.* 2003;31(5):296-301.
159. Putnam L.R., T.G., Ostovar-Kermani, A., Le Blanc, K.T., Anderson, G., Holzmann-Pazgal, K.P., Lally, et al. Surgical site infection reporting: more than meets the agar. *J*

- 1
2
3 Pediatr Surg. 2017;52(1):156-60.
4
5
6
7
8 160. Al-Qahtani S.M., H.M., Al-Amoudi, S., Al-Jehani, A.S., Ashour, M.R., Abd-Hammad,
9 O.R., Tawfik, et al. Post-appendectomy surgical site infection rate after using an
10 antimicrobial film incise drape: A prospective study. Surg Infect. 2015;16(2):155-8.
11
12
13
14
15 161. Quezada F., N., Quezada, R., Mejia, A., Brañes, O., Padilla, N., Jarufe, et al.
16 Laparoscopic versus open approach in the management of appendicitis complicated
17 exclusively with peritonitis: A single center experience. Int J Surg. 2015;13:80-3.
18
19
20
21
22 162. Raakow J., H.-G., Liesaus, P., Neuhaus, R., Raakow. Single-incision versus multiport
23 laparoscopic appendectomy: a case-matched comparative analysis. Surg Endosc Interv
24 Tech. 2015;29(6):1530-6.
25
26
27
28
29 163. Ríos J., C., Murillo, G., Carrasco, C., Humet. Increase in costs attributable to surgical
30 infection after appendectomy and colectomy. Incremento de costes atribuible a la
31 infección quirúrgica de la apendicectomía y colectomía. 2003;17(3):218-25.
32
33
34
35
36
37 164. Rafiq M.S., M.M., Khan, A., Khan, H., Jan. Evaluation of postoperative antibiotics after
38 non-perforated appendectomy. J Pak Med Assoc. 2015;65(8):815-7.
39
40
41
42
43
44
45 165. Reinisch A., J., Heil, G., Woeste, W., Bechstein, J., Liese. The meteorological influence
46 on seasonal alterations in the course of acute appendicitis. J Surg Res. 2017;217:137-43.
47
48
49
50
51 166. Romano A., P., Parikh, P., Byers, N., Namias. Simple acute appendicitis versus non-
52 perforated gangrenous appendicitis: Is there a difference in the rate of post-operative
53 infectious complications? Surg Infect. 2014;15(5):517-20.
54
55
56
57
58 167. Romel Hilaire, Fernández, Zenén Rodríguez, García, Lázaro Ibrahim Romero, Sánchez,
59 Luis Pablo Rodríguez. Apendicectomía videolaparoscópica frente a apendicectomía
60

- 1
2
3 convencional X1 Laparoscopic versus conventional appendectomy. *Revista Cubana de*
4 *Cirugía*. 2014;53(1):30-40.
5
6
7
8
9 168. Romy S., M.-C., Eisenring, V., Bettschart, C., Petignat, P., Francioli, N., Troillet.
10 Laparoscope use and surgical site infections in digestive surgery. *Ann Surg*.
11 2008;247(4):627-32.
12
13
14
15
16 169. Rooh-ul-Muqim, M., Khan, M., Zarin. Experience of laparoscopic appendectomies
17 versus open appendectomies. *Pak J Med Sci*. 2010;26(2):324-8.
18
19
20
21
22 170. Van Rossem C.C., M.H.F., Schreinemacher, K., Treskes, R.M., Van Hogeand,
23 A.A.W., Van Geloven. Duration of antibiotic treatment after appendectomy for acute
24 complicated appendicitis. *Br J Surg*. 2014;101(6):715-9.
25
26
27
28
29
30 171. Van Rossem C.C., M.D., Bolmers, M.H., Schreinemacher, A.A., van Geloven, W.A.,
31 Bemelman. Prospective nationwide outcome audit of surgery for suspected acute
32 appendicitis. *Br J Surg*. 2016;103(1):144-51.
33
34
35
36
37 172. Viet Hung N., T., Anh Thu, V.D., Rosenthal, D., Tat Thanh, N., Quoc Anh, N., Le Bao
38 Tien, et al. Surgical site infection rates in seven cities in Vietnam: Findings of the
39 international nosocomial infection control consortium. *Surg Infect*. 2016;17(2):243-9.
40
41
42
43
44 173. Rotermann M. Infection after cholecystectomy, hysterectomy or appendectomy. *Health*
45 *Rep*. 2004;15(4):11-23.
46
47
48
49 174. Saar S., P., Talving, J., Laos, T., Põdrämägi, M., Sokirjanski, T., Lustenberger, et al.
50 Delay Between Onset of Symptoms and Surgery in Acute Appendicitis Increases
51 Perioperative Morbidity: A Prospective Study. *World J Surg*. 2016;40(6):1308-14.
52
53
54
55
56 175. Saber A.A., M.H., Elgamal, T.H., El-Ghazaly, A.V., Dewoolkar, A., Akl. Simple
57 technique for single incision transumbilical laparoscopic appendectomy. *Int J Surg*.
58 2010;8(2):128-30.
59
60

- 1
2
3
4
5 176. Sadraei-Moosavi S.-M., N., Nikhbakhsh, A.-A., Darzi. Postoperative antibiotic therapy
6 after appendectomy in patients with non-perforated appendicitis. *Caspian J Int Med.*
7 2017;8(2):104-7.
8
9
10
11
12 177. Saha N., D.K., Saha, M.A., Rahman, M.K., Islam, M.A., Aziz. Comparison of post
13 operative morbidity between laparoscopic and open appendectomy in children.
14 *Mymensingh Med J.* 2010;19(3):348-52.
15
16
17
18 178. Sahm M., R., Kube, S., Schmidt, C., Ritter, M., Pross, H., Lippert. Current analysis of
19 endoloops in appendiceal stump closure. *Surg Endosc Interv Tech.* 2011;25(1):124-9.
20
21
22
23
24 179. Sahm M., M., Pross, R., Otto, A., Koch, I., Gastinger, H., Lippert. Clinical health service
25 research on the surgical therapy of acute appendicitis: Comparison of outcomes based on 3
26 German multicenter quality assurance studies over 21 years. *Ann Surg.* 2015;262(2):338-
27 46.
28
29
30
31
32
33 180. Salö M., E., Järbur, M., Hambraeus, B., Ohlsson, P., Stenström, E., Arnbjörnsson. Two-
34 trocar appendectomy in children - description of technique and comparison with
35 conventional laparoscopic appendectomy. *BMC Surg.* 2016;16(1):52.
36
37
38
39 181. Sánchez-Santana T., J.A., del-Moral-Luque, P., Gil-Yonte, L., Bañuelos-Andrío, M.,
40 Durán-Poveda, G., Rodríguez-Caravaca. Effect of compliance with an antibiotic
41 prophylaxis protocol in surgical site infections in appendectomies. Prospective cohort
42 study. Efecto de la adecuación a protocolo de la profilaxis antibiótica en la incidencia de
43 infección quirúrgica en apendicectomías Estudio de cohortes prospectivo. 2017;85(3):208-
44 13.
45
46
47
48
49
50
51 182. Sauvain M.-O., K., Slankamenac, M.K., Muller, S., Wildi, U., Metzger, W., Schmid, et
52 al. Delaying surgery to perform CT scans for suspected appendicitis decreases the rate of
53 negative appendectomies without increasing the rate of perforation nor postoperative
54 complications. *Langenbeck's Arch Surg.* 2016;401(5):643-9.
55
56
57
58
59
60

- 1
2
3 183. Scarborough J.E., K.M., Bennett, T.N., Pappas. Racial disparities in outcomes after
4 appendectomy for acute appendicitis. *Am J Surg.* 2012;204(1):11-7.
5
6
7
8 184. Seifarth F.G., N., Kundu, A.D., Guerron, M.M., Garland, M.W., Gaffley, S., Worley, et
9 al. Umbilical Negative Pressure Dressing for Transumbilical Appendectomy in Children.
10 *JLS.* 2016;20(4).
11
12
13
14
15 185. Federico G. Seifarth, Kundu, Neilendu, Guerron, Alfredo D., Garland, Mary M.,
16 Gaffley, Michaela W. G., Worley, Sarah, et al. Umbilical Negative Pressure Dressing for
17 Transumbilical Appendectomy in Children. *JLS-JOURNAL OF THE SOCIETY OF*
18 *LAPAROENDOSCOPIC SURGEONS.* 2016;20(4).
19
20
21
22
23
24 186. Senekjian L., R., Nirula. Tailoring the operative approach for appendicitis to the patient:
25 A prediction model from national surgical quality improvement program data. *J Am Coll*
26 *Surg.* 2013;216(1):34-40.
27
28
29
30
31 187. Sesia S.B., M., Frech, F.-M., Häcker, J., Mayr. Laparoscopic "single-port"
32 appendectomy in children. *Laparoskopische "single port"-appendektomie im kindesalter.*
33 2011;136(1):50-5.
34
35
36
37
38 188. Shaikh A.R., S., Khatoon, M., Arif. Evaluation of re-admission after open
39 appendicectomy. *Rawal Med J.* 2011;36(2):100-3.
40
41
42
43
44
45 189. Shang Q., Q., Geng, X., Zhang, C., Guo. The efficacy of combined therapy with
46 metronidazole and broad-spectrum antibiotics on postoperative outcomes for pediatric
47 patients with perforated appendicitis. *Medicine.* 2017;96(47).
48
49
50
51 190. Shindholimath V., K., Thinakaran, T., Rao, Y., Veerappa. Laparoscopic management
52 of appendicular mass. *J Minimal Access Surg.* 2011;7(2):136-40.
53
54
55
56 191. Shimizu T., M., Ishizuka, K., Kubota. The preoperative serum C-reactive protein level
57 is a useful predictor of surgical site infections in patients undergoing appendectomy. *Surg*
58 *Today.* 2015;45(11):1404-10.
59
60

- 1
2
3
4
5 192. Siam B., A., Al-Kurd, N., Simanovsky, H., Awesat, Y., Cohn, B., Helou, et al.
6 Comparison of appendectomy outcomes between senior general surgeons and general
7 surgery residents. *JAMA Surg.* 2017;152(7):679-85.
8
9
10
11
12 193. 서승원, 김신곤. Acute Appendicitis in Pregnant Patients and Non-Pregnant Patients:
13 Recent Clinical Experience of the Tertiary Hospital Z1 임신부와 가임기 여성의 급성
14 충수염의 비교 고찰: 최근 3차 병원의 임상적 경험. *Annals of Surgical Treatment and*
15 *Research S1 대한외과학회지.* 2002;62(6):486-90.
16
17
18
19
20
21
22
23
24
25
26
27
28 194. Alfredo Silva, M, Guido Vargas, A, Amparo Moreno, H, Pablo Becerra. Utilidad del
29 retractor elástico abdominal para disminuir el riesgo de infección de herida operatoria en
30 apendicitis aguda X1 Use of an elasticwall retractor during appendectomy to reduce wound
31 infection. *Revista chilena de cirugía.* 2008;60(6):527-33.
32
33
34
35
36
37 195. Singh V.K., K., Nishant, B., Kharga, A.K., Kalita, P., Bhutia, J., Jain. Randomized
38 controlled trial comparing open, conventional, and single port laparoscopic appendectomy.
39 *J Clin Diagn Res.* 2017;11(10):PC05-PC10.
40
41
42
43
44
45
46 196. Siribumrungwong B., K., Srikuea, A., Thakkinstian. Comparison of superficial surgical
47 site infection between delayed primary and primary wound closures in ruptured
48 appendicitis. *Asian J Surg.* 2014;37(3):120-4.
49
50
51
52
53 197. Sivrikoz E., E., Karamanos, E., Beale, P., Teixeira, K., Inaba, D., Demetriades. The
54 effect of diabetes on outcomes following emergency appendectomy in patients without
55 comorbidities: A propensity score-matched analysis of National Surgical Quality
56 Improvement Program database. *Am J Surg.* 2015;209(1):206-11.
57
58
59
60

- 1
2
3 198. Soll C., P., Wyss, H., Gelpke, D.A., Raptis, S., Breitenstein. Appendiceal stump closure
4 using polymeric clips reduces intra-abdominal abscesses. *Langenbeck's Arch Surg.*
5 2016;401(5):661-6.
6
7
8
9
10 199. Sozutek A., T., Colak, M., Dirlik, K., Ocal, O., Turkmenoglu, A., Dag. A prospective
11 randomized comparison of single-port laparoscopic procedure with open and standard 3-
12 port laparoscopic procedures in the treatment of acute appendicitis. *Surg Laparoscopy*
13 *Endosc Percutaneous Tech.* 2013;23(1):74-8.
14
15
16
17
18 200. Srishewachart P., S., Narksut. Incidence of abnormal preoperative blood testing and
19 postoperative complication in appendectomy patients in Siriraj Hospital. *J Med Assoc*
20 *Thailand.* 2016;99(5):517-24.
21
22
23
24
25 201. Staszewicz W., M.-C., Eisenring, V., Bettschart, S., Harbarth, N., Troillet. Thirteen
26 years of surgical site infection surveillance in Swiss hospitals. *J Hosp Infect.* 2014;88(1):40-
27 7.
28
29
30
31
32 202. Suttie S.A., S., Seth, C.P., Driver, A.A., Mahomed. Outcome after intra- and extra-
33 corporeal laparoscopic appendectomy techniques. *Surg Endosc.* 2004;18(7):1123-5.
34
35
36
37 203. Svensson J.F., B., Patkova, M., Almström, S., Eaton, T., Wester. Outcome after
38 introduction of laparoscopic appendectomy in children: A cohort study. *J Pediatr Surg.*
39 2016;51(3):449-53.
40
41
42
43
44 204. Taguchi Y., S., Komatsu, E., Sakamoto, S., Norimizu, Y., Shingu, H., Hasegawa.
45 Laparoscopic versus open surgery for complicated appendicitis in adults: a randomized
46 controlled trial. *Surg Endosc Interv Tech.* 2016;30(5):1705-12.
47
48
49
50
51 205. Tanaka S., D., Kubota, S.H., Lee, K., Oba, M., Matsuyama. Effectiveness of
52 laparoscopic approach for acute appendicitis. *Osaka City Med J.* 2007;53(1):1-8.
53
54
55
56 206. Tijerina J., R., Velasco-Rodríguez, C., Vásquez, V., Melnikov, S., Rodriguez.
57 Effectiveness of a systemic antibiotic followed by topical ionized solution as surgical site
58 infection prophylaxis. *J Int Med Res.* 2010;38(4):1287-93.
59
60

- 1
2
3
4
5 207. The SCARLESS Study Group. Single port/incision laparoscopic surgery compared with
6 standard three-port laparoscopic surgery for appendectomy: A randomized controlled
7 trial. *Surg Endosc Interv Tech*. 2015;29(1):77-85.
8
9
10
11 208. Juan Pablo Toro, Barrera, Óscar Javier, Morales, Carlos Hernando. Superioridad clínica
12 de la apendicectomía laparoscópica sobre la técnica abierta: ¿adopción lenta de un nuevo
13 estándar de tratamiento? X1 Clinical superiority of laparoscopic appendectomy over the
14 open technique: sluggish adoption of a new standard of treatment? *Revista Colombiana de*
15 *Cirugía*. 2017;32(1):32-9.
16
17
18
19
20
21
22 209. Towfigh S., T., Clarke, W., Yacoub, A.H., Pooli, R.J., Mason, N., Katkhouda, et al.
23 Significant reduction of wound infections with daily probing of contaminated wounds : A
24 prospective randomized clinical trial. *Arch Surg*. 2011;146(4):448-52.
25
26
27
28
29 210. Troillet N., E., Aghayev, M.-C., Eisenring, A.F., Widmer. First Results of the Swiss
30 National Surgical Site Infection Surveillance Program: Who Seeks Shall Find. *Infect*
31 *Control Hosp Epidemiol*. 2017;38(6):697-704.
32
33
34
35
36 211. Tsioplis C., C., Brockschmidt, S., Sander, D., Henne-Bruns, M., Kornmann. Factors
37 influencing the course of acute appendicitis in adults and children. *Langenbeck's Arch Surg*.
38 2013;398(6):857-67.
39
40
41
42
43 212. Vahdad M.R., M., Nissen, A., Semaan, T., Klein, E., Palade, T., Boemers, et al.
44 Experiences with LESS-appendectomy in Children. *Arch Iran Med*. 2016;19(1):57-63.
45
46
47
48 213. Van Rossem C.C., M.H.F., Schreinemacher, A.A.W., Van Geloven, W.A., Bemelman,
49 G.J.D., Van Acker, B., Akkermans, et al. Antibiotic duration after laparoscopic
50 appendectomy for acute complicated appendicitis. *JAMA Surg*. 2016;151(4):323-9.
51
52
53
54
55 214. Van Rossem C.C., M.H.F., Schreinemacher, K., Treskes, R.M., Van Hogeand,
56 A.A.W., Van Geloven. Duration of antibiotic treatment after appendectomy for acute
57 complicated appendicitis. *Br J Surg*. 2014;101(6):715-9.
58
59
60

- 1
2
3 215. Wang-Chan A., F.H., Hetzer, C., Gingert, C., Gingert, E., Angst, E., Angst, et al.
4 Clinical relevance and effect of surgical wound classification in appendicitis: Retrospective
5 evaluation of wound classification discrepancies between surgeons, Swissnoso-trained
6 infection control nurse, and histology as well as surgical site infection rates by wound class.
7
8 J Surg Res. 2017;215:132-9.
9
10
11
12
13 216. Watanabe A., S., Kohnoe, H., Sonoda, K., Shirabe, K., Fukuzawa, S., Maekawa, et al.
14 Effect of intra-abdominal absorbable sutures on surgical site infection. Surg Today.
15 2012;42(1):52-9.
16
17
18
19
20 217. Willis Z.I., E.M., Duggan, B.T., Bucher, J.B., Pietsch, M., Milovancev, W., Wharton,
21 et al. Effect of a clinical practice guideline for pediatric complicated appendicitis. JAMA
22 Surg. 2016;151(5).
23
24
25
26
27 218. Ramírez-Wong F.M., T., Atencio-Espinoza, V.D., Rosenthal, E., Ramirez, S.L., Torres-
28 Zegarra, Z.R., Díaz Tavera, et al. Surgical Site Infections Rates in More Than 13,000
29 Surgical Procedures in Three Cities in Peru: Findings of the International Nosocomial
30 Infection Control Consortium. Surg Infect. 2015;16(5):572-6.
31
32
33
34
35
36 219. Wu J.-M., K.-H., Chen, H.-F., Lin, L.-M., Tseng, S.-H., Tseng, S.-H., Huang.
37 Laparoscopic appendectomy in pregnancy. J Laparoendosc Adv Surg Techn Part A.
38 2005;15(5):447-50.
39
40
41
42
43 220. Wu H.-S., H.-W., Lai, S.-J., Kuo, Y.-T., Lee, D.-R., Chen, C.-W., Chi, et al. Competitive
44 edge of laparoscopic appendectomy versus open appendectomy: A subgroup comparison
45 analysis. J Laparoendosc Adv Surg Techn. 2011;21(3):197-202.
46
47
48
49
50 221. Wu K., L., Yang, A., Wu, J., Wang, S., Xu, H., Zhao, et al. Single-site laparoscopic
51 appendectomy in children using conventional instruments: a prospective, randomized,
52 control trial. Pediatr Surg Int. 2014;31(2):167-71.
53
54
55
56
57 222. Wu T.-C., Q., Lu, Z.-Y., Huang, X.-H., Liang. Efficacy of emergency laparoscopic
58 appendectomy in treating complicated appendicitis for elderly patients. Saudi Med J.
59 2017;38(11):1108-12.
60

- 1
2
3
4
5 223. Yaghoubian A., C., de Virgilio, V., Chiu, S.L., Lee. "July effect" and appendicitis. *J Surg Educ.* 2010;67(3):157-60.
6
7
8
9
10 224. Yagnik V., J., Rathod, A., Phatak. A retrospective study of two-port appendectomy and
11 its comparison with open appendectomy and three-port appendectomy. *Saudi J*
12 *Gastroenterol.* 2010;16(4):268-71.
13
14
15
16
17 225. Yousef Y., F., Youssef, M., Homsy, T., Dinh, K., Pandya, H., Stagg, et al.
18 Standardization of care for pediatric perforated appendicitis improves outcomes. *J Pediatr*
19 *Surg.* 2017;52(12):1916-20.
20
21
22
23
24 226. Zhang Z., Y., Wang, R., Liu, L., Zhao, H., Liu, J., Zhang, et al. Suprapubic single-
25 incision versus conventional laparoscopic appendectomy. *J Surg Res.* 2016;200(1):131-8.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

MOOSE Checklist for Meta-analyses of Observational Studies

Item No	Recommendation	Reported on Page No
Reporting of background should include		
1	Problem definition	5
2	Hypothesis statement	6
3	Description of study outcome(s)	5
4	Type of exposure or intervention used	NA
5	Type of study designs used	5-6
6	Study population	5-6
Reporting of search strategy should include		
7	Qualifications of searchers (eg, librarians and investigators)	13
8	Search strategy, including time period included in the synthesis and key words	6; Suppl. Table 1
9	Effort to include all available studies, including contact with authors	6
10	Databases and registries searched	6
11	Search software used, name and version, including special features used (eg, explosion)	6
12	Use of hand searching (eg, reference lists of obtained articles)	6
13	List of citations located and those excluded, including justification	8, Suppl. Fig 1, Suppl. References
14	Method of addressing articles published in languages other than English	7
15	Method of handling abstracts and unpublished studies	7
16	Description of any contact with authors	6
Reporting of methods should include		
17	Description of relevance or appropriateness of studies assembled for assessing the hypothesis to be tested	6
18	Rationale for the selection and coding of data (eg, sound clinical principles or convenience)	6-7
19	Documentation of how data were classified and coded (eg, multiple raters, blinding and interrater reliability)	6-7
20	Assessment of confounding (eg, comparability of cases and controls in studies where appropriate)	8
21	Assessment of study quality, including blinding of quality assessors, stratification or regression on possible predictors of study results	8
22	Assessment of heterogeneity	7
23	Description of statistical methods (eg, complete description of fixed or random effects models, justification of whether the chosen models account for predictors of study results, dose-response models, or cumulative meta-analysis) in sufficient detail to be replicated	7-8
24	Provision of appropriate tables and graphics	Table 1; Fig 1-3
Reporting of results should include		
25	Graphic summarizing individual study estimates and overall estimate	Fig 1-3
26	Table giving descriptive information for each study included	Suppl. Table 1
27	Results of sensitivity testing (eg, subgroup analysis)	9; Table 1

1	28	Indication of statistical uncertainty of findings	9; Table 1; Fig 1-3
2			
3			
4	Item No	Recommendation	Reported on Page No
5			
6			
7	Reporting of discussion should include		
8			
9	29	Quantitative assessment of bias (eg, publication bias)	12-13
10	30	Justification for exclusion (eg, exclusion of non-English language citations)	12-13
11			
12	31	Assessment of quality of included studies	12-13
13	Reporting of conclusions should include		
14			
15	32	Consideration of alternative explanations for observed results	13
16	33	Generalization of the conclusions (ie, appropriate for the data presented and within the domain of the literature review)	13
17			
18	34	Guidelines for future research	13
19			
20	35	Disclosure of funding source	14

21 From: Stroup DF, Berlin JA, Morton SC, et al, for the Meta-analysis Of Observational Studies in Epidemiology
 22 (MOOSE) Group. Meta-analysis of Observational Studies in Epidemiology. A Proposal for Reporting. *JAMA*.
 23 2000;283(15):2008-2012. doi: 10.1001/jama.283.15.2008.



PRISMA 2009 Checklist

Section/topic	#	Checklist item	Reported on page #
TITLE			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	1
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	2-3
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known.	5-6
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	6
METHODS			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	8
Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	6
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	6
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	6
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	7
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	7
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	7
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	8
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).	7-8
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., I^2) for each meta-analysis.	7-8



PRISMA 2009 Checklist

Section/topic	#	Checklist item	Reported on page #
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	7-8
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	8
RESULTS			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	8
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	8
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	8
Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	8-9
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	8-9
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).	8-9
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).	9
DISCUSSION			
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).	10
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	12
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	12
FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	14

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit: www.prisma-statement.org.