

In the format provided by the authors and unedited.

Accumulation of succinate controls activation of adipose tissue thermogenesis

Evanna L. Mills^{1,2}, Kerry A. Pierce³, Mark P. Jedrychowski^{1,2}, Ryan Garrity¹, Sally Winther^{1,2}, Sara Vidoni^{1,2}, Takeshi Yoneshiro⁴, Jessica B. Spinelli², Gina Z. Lu¹, Lawrence Kazak⁵, Alexander S. Banks⁶, Marcia C. Haigis², Shingo Kajimura⁴, Michael P. Murphy⁷, Steven P. Gygi², Clary B. Clish³ & Edward T. Chouchani^{1,2*}

¹Department of Cancer Biology, Dana–Farber Cancer Institute, Boston, MA, USA. ²Department of Cell Biology, Harvard Medical School, Boston, MA, USA. ³Broad Institute of Harvard and MIT, Cambridge, MA, USA. ⁴Diabetes Center, University of California, San Francisco, San Francisco, CA, USA. ⁵Goodman Cancer Research Centre, McGill University, Montreal, Quebec, Canada. ⁶Division of Endocrinology, Diabetes, and Hypertension, Brigham and Women’s Hospital and Harvard Medical School, Boston, MA, USA. ⁷MRC Mitochondrial Biology Unit, University of Cambridge, Cambridge Biomedical Campus, Cambridge, UK. *e-mail: edwardt_chouchani@dfci.harvard.edu

uncropped blots from Extended Data Fig. 6

