

SUPPLEMENTARY TABLE S2. GLYCEMIC OUTCOMES DURING HYBRID CLOSED-LOOP STRATIFIED BY STANDARD THERAPY MODALITY
(PUMP OR MULTIPLE DAILY INJECTIONS)

	<i>Adults</i>		<i>Adolescents</i>		<i>Children</i>	
	<i>Pump</i> (n=8)	<i>MDI</i> (n=3)	<i>Pump</i> (n=7)	<i>MDI</i> (n=3)	<i>Pump</i> (n=12)	<i>MDI</i> (n=3)
Percentage time in range, %						
<54 mg/dL	0.2±0.3	0.2±0.3	0.4±0.5	0.2±0.2	0.4±0.4	0.2±0.0
<70 mg/dL	2.0±1.4	1.4±1.1	2.7±2.0	2.1±2.1	2.2±2.1	2.0±0.8
70–180 mg/dL	74.7±8.5	71±3.7	79.5±11.2	77.8±18.3	66.9±13.9	78.4±7.3
≥250 mg/dL	3.7±4.1	6.7±4.6	3.5±5.4	3.5±5.1	10.0±9.3	3.0±1.9
Time in HCL (%)	98.4±1.1	95.4±1.5	98.9±0.7	97.6±2.1	98.2±1.7	98.9±0.5
Hypoglycemic events per participant per day ^a	1.1	0.7	1.1	0.9	1.1	0.8

Data are mean ± SD or median [IQR].

^aCapillary BG <70 mg/dL.

BG, blood glucose; HCL, hybrid closed-loop; IQR, interquartile range; MDI, multiple daily injections.