

Article Title: Clinicopathological correlations of mesenteric fibrosis and evaluation of a novel biomarker for fibrosis detection in small bowel neuroendocrine neoplasms

Journal: Endocrine

Authors: Faidon-Marios Laskaratos, Dalvinder Mandair, Andrew Hall, Sarah Alexander, Conrad von Stempel, Josephine Bretherton, TuVinh Luong, Jennifer Watkins, Olagunju Ogunbiyi, Krista Rombouts, Martyn Caplin, Christos Toumpanakis

Corresponding author: Dr Faidon-Marios Laskaratos MD, MSc, MRCP, Royal Free London NHS Foundation Trust, flaskaratos@gmail.com

Online Resource 1. Summary of demographic and clinical characteristics of patients enrolled in the biomarker assessment study

	Patients with midgut NETs who underwent surgery (n=20)
	n (%)
Age (mean±SD)	63±10
Sex	
Male	13 (65%)
Female	7 (35%)
Grade	
1	13 (65%)
2	7 (35%)
Extent of disease	
Localised	2 (10%)
Locoregional	5 (25%)
Metastatic	13 (65%)
Mesenteric mass	18 (90%)
Liver metastases	10 (50%)
Distant extrahepatic metastases	5 (25%)
Macroscopic mesenteric fibrosis	17 (85%)
Medical therapy	
Octreotide LAR	6 (30%)
Lanreotide Autogel	7 (35%)
Surgical therapy	
Small bowel resection	1 (5%)
Gastro-jejunal bypass	1 (5%)

Right hemicolectomy (R0)	10 (50%)
Right hemicolectomy (R1)	8 (40%)