


Cell Reports

Hypoxia Produces Pro-arrhythmic Late Sodium Current in Cardiac Myocytes by SUMOylation of $\text{Na}_v1.5$ Channels

Graphical Abstract



Authors

Leigh D. Plant, Dazhi Xiong,
Jesus Romero, Hui Dai,
Steve A.N. Goldstein

Correspondence

I.plant@northeastern.edu (L.D.P.),
sgoldst2@uci.edu (S.A.N.G.)

In Brief

The cardiac channel $\text{Na}_v1.5$ passes pro-arrhythmic late sodium currents in response to hypoxia. Plant et al. demonstrate the pathophysiological mechanism to be rapid, hypoxia-induced monoSUMOylation of $\text{Na}_v1.5$ channels. Blocking SUMOylation of lysine₄₄₂ prevents hypoxia-induced late currents and attendant prolongation of the action potential in human cardiomyocytes derived from pluripotent stem cells.

Highlights

- Acute cardiac hypoxia elevates late sodium current (I_{LATE}) to pro-arrhythmic levels
- Increased I_{LATE} is due to rapid SUMOylation of $\text{Na}_v1.5$ channels at the cell membrane
- SUMOylation of lysine₄₄₂ reopens $\text{Na}_v1.5$ channels when they are normally inactive
- Blocking SUMOylation prevents increased I_{LATE} and action potential prolongation



Hypoxia Produces Pro-arrhythmic Late Sodium Current in Cardiac Myocytes by SUMOylation of Na_v1.5 Channels

Leigh D. Plant,^{1,*} Dazhi Xiong,² Jesus Romero,² Hui Dai,² and Steve A.N. Goldstein^{2,3,*}

¹Department of Pharmaceutical Sciences, Northeastern University, 360 Huntington Avenue, Boston, MA 02115, USA

²Departments of Pediatrics and Physiology & Biophysics, University of California, Irvine, 1001 Health Sciences Road, Irvine Hall, Irvine, CA 92697, USA

³Lead Contact

*Correspondence: l.plant@northeastern.edu (L.D.P.), sgoldst2@uci.edu (S.A.N.G.)

<https://doi.org/10.1016/j.celrep.2020.01.025>

SUMMARY

Acute cardiac hypoxia produces life-threatening elevations in late sodium current (I_{LATE}) in the human heart. Here, we show the underlying mechanism: hypoxia induces rapid SUMOylation of Na_v1.5 channels so they reopen when normally inactive, late in the action potential. Na_v1.5 is SUMOylated only on lysine 442, and the mutation of that residue, or application of a deSUMOylating enzyme, prevents hypoxic reopenings. The time course of SUMOylation of single channels in response to hypoxia coincides with the increase in I_{LATE} , a reaction that is complete in under 100 s. In human cardiac myocytes derived from pluripotent stem cells, hypoxia-induced I_{LATE} is confirmed to be SUMO-dependent and to produce action potential prolongation, the pro-arrhythmic change observed in patients.

INTRODUCTION

In the human heart, Na_v1.5 voltage-gated sodium channels pass I_{Na} , a rapidly activating and inactivating Na⁺ current that determines the rise, duration, and timing of action potentials (Marbán, 2002; Catterall, 1986). In addition, Na_v1.5 channels pass a small, persistent Na⁺ current, the late sodium current (I_{LATE}), which contributes to maintaining the action potential plateau (Makielski, 2016). In healthy hearts, I_{LATE} is less than 0.5% of the magnitude of peak I_{Na} and is passed by a small number of channels that do not stay closed after inactivation (Makielski, 2016). In patients with an ischemic and failing heart, sudden infant death syndrome (SIDS), and mutations in SCN5A, the gene that encodes the Na_v1.5 channel, which produces long QT syndrome, the size of I_{LATE} can increase to 4%–5% of peak I_{Na} (Bennett et al., 1995; Plant et al., 2006; Belardinelli et al., 2015). Furthermore, acute hypoxia and ischemia have been recognized to increase I_{LATE} in cardiac myocytes (Saint et al., 1992; Ju et al., 1996; Carmeliet, 1999; Belardinelli et al., 2006) prior to slower processes like remodeling (West, 2017), and excess I_{LATE} has been shown to be pro-arrhythmic because it prolongs action potential duration (APD), reducing repolarization reserve, increasing suscepti-

bility to after-depolarizations, and causing a predisposition to torsades de pointes (TdP), a ventricular dysrhythmia that is lethal when sustained (Gaur et al., 2009; Shryock et al., 2013; Chadda et al., 2017). Thus, increases in I_{LATE} above baseline by just 0.3% to 1% predisposition to sudden cardiac death (Bennett et al., 1995; Belardinelli et al., 2015). These observations underpin proposals that inhibiting I_{LATE} has therapeutic potential (Belardinelli et al., 2006; Song et al., 2006).

We recently reported that the rapid influx of Na⁺ flux into neurons in response to acute hypoxic challenge is due to SUMOylation of Na_v1.2, the major voltage-gated sodium channel in the brain; in that channel, SUMO modification of lysine 38 alters opening and closing with changes in voltage and does not effect I_{LATE} (Plant et al., 2016). SUMOylation is the enzyme-mediated linkage of one of three SUMO isoforms to the ε amino group of specific lysine residues in a target protein (Henley et al., 2014). Present in all eukaryotic cells, the SUMO pathway was known to regulate the trafficking and activity of nuclear transcription factors when we described it to operate as well at the plasma membrane by direct SUMOylation of Na⁺ and K⁺ channels to regulate excitability (Rajan et al., 2005; Plant et al., 2010, 2011, 2012, 2016; Xiong et al., 2017). Given the important role of hypoxia-induced increases in I_{LATE} in heart disease, we tested the hypothesis that SUMOylation of Na_v1.5 channels was the underlying mechanism.

We demonstrate that hypoxia induces a rapid increase in I_{LATE} in human cardiac myocytes derived from pluripotent stem cells (iPS-CMs). The response is reproduced by application of cytosolic SUMO1 at ambient levels of oxygen and suppressed by the deSUMOylating enzyme SENP1. SUMOylation of Na_v1.5 on lysine 442, a residue located in the segment between channel domains I and II, is shown to be necessary and sufficient to explain hypoxia-induced changes in I_{LATE} by reconstitution of the response in heterologous cells and by studies using whole-cell and single-channel patch-clamp recording and live cell Förster resonance energy transfer (FRET). The implied mechanism was confirmed using total internal reflection fluorescence microscopy (TIRFM) to study single particles on the surface of live cells in real-time; the time course was the same for hypoxic challenge, the recruitment of SUMO1 to Na_v1.5 channels on cell surface, and the increase in I_{LATE} . Furthermore, hypoxia-induced, SUMO-dependent increases in I_{LATE} were observed to increase APD






Figure 1. Acute Hypoxia and SUMO1 Augment the I_{LATE} in Human iPS-CMs

I_{Na} in human iPS-CMs was studied by whole-cell patch clamp. I_{LATE} was measured at -30 mV and is shown as a mean percentage of the peak current remaining between 50 and 100 ms after the peak current. Hypoxia challenge was a drop in O_2 from 21% to 1.5%. The O_2 level was measured in real time in the recording chamber and fell from 21% to 1.5% O_2 in 11 ± 3 s (not shown). The time course of hypoxic modulation of I_{Na} was studied by steps from -100 mV to -30 mV every 10 s. Whole-cell currents were not studied for more than 600 s to maintain consistent voltage clamp control of I_{Na} . Cells were studied with control solution (black), 1 nM SUMO1 (red), or 1 nM SENP1 (blue) in the recording pipette. Data are mean \pm SEM for 8 to 10 cells per group. Measured values and statistical analyses are listed in Table 1.

(A) Left: example traces showing I_{LATE} at -30 mV increased when control perfusate at 21% O_2 (black) was exchanged with a hypoxic solution at 1.5% O_2

(black dash). Inset: the increase in I_{LATE} shown in response to hypoxia. Right: the time course for changes in I_{LATE} in response to hypoxia and on return to normoxia (21%). Scale bars represent 500 pA and 10 ms.

(B) Left: SUMO1 in the pipette (red) increased I_{LATE} . No further change in I_{LATE} was observed when the control perfusate was exchanged with a hypoxic solution at 1.5% O_2 (red dash). Right: I_{LATE} was not observed with SENP1 in the pipette when cells were studied at 21% O_2 (blue) or 1.5% O_2 (blue dash). In each case, the dashed lines overlap with the solid lines.

(C) A histogram showing the mean I_{LATE} as a percentage of the peak current from cells studied at 21% O_2 (solid color) after 200 s at 1.5% O_2 (hatched color) and after 200 s following a return to 21% O_2 (solid color) with control solution (black), 1 nM SUMO1 (red), or 1 nM SENP1 (blue) in the recording pipette.

in human induced pluripotent stem cells (iPSCs) by recording spontaneous action potentials in current-clamp mode; and the changes were suppressed by ranolazine, a drug that inhibits $Na_v1.5$ channel late current. Incorporating the measured hypoxia-induced increase in I_{LATE} in iPS-CMs into the O'Hara-Rudy model for human cardiac action potentials (O'Hara et al., 2011) was sufficient to reproduce the observed pro-arrhythmic increase in APD. Given the role of I_{LATE} in arrhythmogenesis, SUMOylation of $Na_v1.5$ is identified as a mechanistically defined target for therapeutic intervention.

RESULTS

Hypoxia Rapidly Increases I_{LATE} in Human iPS-CMs

At ambient O_2 levels (21%), I_{Na} in human iPS-CMs activated rapidly to a mean peak of -40 ± 4 pA/pF when measured at -30 mV in whole-cell mode (Figure 1A; Table 1). I_{Na} then inactivated to produce a residual I_{LATE} that was $0.46\% \pm 0.1\%$ of the magnitude of peak current 50 to 100 ms after the onset of inactivation. When O_2 was lowered from ambient levels to 1.5% by the perfusion of cells with a hypoxic solution (Plant et al., 2016), mean peak I_{Na} was unchanged (-42 ± 3 pA/pF), but I_{LATE} increased ~ 10 -fold to $\sim 4.4\%$ (-1.85 pA/pF) of peak I_{Na} after 100 s of hypoxia. The hypoxia-induced increase in I_{LATE} rose to half its maximal level in 40 ± 2 s and remained stable for at least 3 min after iPS-CMs were restored to ambient O_2 . Cells were not studied beyond 600 s to maintain consistent voltage-clamp control of I_{Na} (Figure 1A). These changes are like those observed in isolated rat ventricular myocytes (Ju et al., 1996) and human cardiac muscle (Makielski, 2016; Wei et al., 2016).

SUMO1 Increases I_{LATE} in Human iPS-CMs

We delivered 1 nM purified SUMO1 peptide into iPS-CMs by the patch pipette; a concentration that evoked maximal effects on $Na_v1.2$ channels in cerebellar granule neurons (CGNs) and three different K^+ channels in CGNs, hippocampal neurons, and rat cardiac myocytes (Xiong et al., 2017; Plant et al., 2011, 2012, 2016). SUMO1 induced changes in cardiac I_{Na} like those produced by hypoxia: I_{LATE} increased to $\sim 4.2\%$ of peak I_{Na} without a change in the peak magnitude (Figures 1B and 1C; Table 1). When SUMO1 was in the recording pipette, changing the solution bathing iPS-CMs from 21% to 1.5% O_2 did not further alter the parameters of I_{Na} peak or I_{LATE} . The notion that hypoxic regulation of I_{Na} proceeds by SUMOylation was further supported by the observation that delivering 1 nM SENP1 deSUMOylase into the iPS-CMs by the recording pipette suppressed the increase in I_{LATE} induced by hypoxia (Figures 1B and 1C; Table 1).

I_{LATE} is observed in the "window" between the voltages where Na_v channels activate and inactivate (Figure S1; Table 1). Thus, exposure of the iPS-CMs to 1.5% O_2 caused a leftward shift in the half-maximal activation voltage of I_{Na} ($V_{1/2-act}$) of -6 ± 4 mV and a rightward shift in the steady-state inactivation midpoint ($V_{1/2-SSI}$) of 6 ± 3 mV, increasing the range of voltages over which I_{Na} channels are likely to be activated but not fully inactivated. In ambient O_2 , SUMO1 in the pipette increased the magnitude of the window current in a manner like that observed on exposure to 1.5% O_2 , and hypoxic challenge with SUMO1 in the pipette did not further alter the parameters. Furthermore, inclusion of 1 nM SENP1 prevented the effects of hypoxia, maintaining the window current at control levels.

Table 1. Effects of Hypoxia, SUMO1, and SENP1 on I_{Na} in iPS-CMs and Nav1.5 Channels in CHO Cells

Condition	I_{Na} -iPS-CMs						Nav1.5 CHO-K1						Nav1.5-K442Q CHO-K1						
	Activation			Current, pA/pF			SSI			Activation			SSI			Current, pA/pF			
	$V_{1/2}$, mV	K	$V_{1/2}$, mV	I_{PEAK}	I_{LATE}	I_{PEAK}	$V_{1/2}$, mV	K	$V_{1/2}$, mV	I_{PEAK}	I_{LATE}	I_{PEAK}	$V_{1/2}$, mV	K	$V_{1/2}$, mV	I_{PEAK}	I_{LATE}	I_{PEAK}	I_{LATE}
O ₂	-44.3 ± 2	7.1 ± 1	-87.2 ± 3	6.2 ± 1	-40 ± 4	-0.22 ± 1	-45.1 ± 2	7.8 ± 1	-87.2 ± 1	-87.2 ± 2	6.5 ± 1	-198 ± 6	-0.96 ± 0.2	-47.0 ± 3	7.4 ± 2	-87.4 ± 2	6 ± 2	-200 ± 4	-0.93 ± 0.3
21% to 1.5%	-50.5 ± 1*	5 ± 1*	-81.7 ± 3*	4.6 ± 1*	-1.85 ± 0.5	-50.3 ± 3*	4.4 ± 2*	81 ± 2*	-81 ± 2*	-81 ± 2*	4.5 ± 2*	-201 ± 4	-8.3 ± 0.6	-46 ± 1	7.2 ± 2	-87.2 ± 1	5.8 ± 3	-197 ± 6	-0.91 ± 0.4
SENP1	-44.6 ± 2	7 ± 2	-89 ± 2.5	6.5 ± 2	-0.28 ± 0.7	-44.7 ± 3	7.8 ± 2	-85.2 ± 3	6.2 ± 1	-196 ± 5	-1.1 ± 0.4	-48 ± 2.3	7.3 ± 2	-83.7 ± 2.5	5.9 ± 2	-194 ± 5	-0.88 ± 0.4		
SENP1	-43.7 ± 2	7.2 ± 1	-90 ± 2	6.4 ± 2	0.25 ± 0.8	-45.4 ± 2	7.6 ± 1	-85.3 ± 2	6.7 ± 3	-197 ± 4	-1.3 ± 0.3	-46.7 ± 3.7	7.2 ± 3	-86.2 ± 2	5.9 ± 1	-199 ± 5	-0.93 ± 0.5		
SUMO1	-52 ± 2*	5.2 ± 1*	-81.9 ± 2*	4.3 ± 1*	-2.18 ± 0.9	-51.5 ± 4*	4.5 ± 1*	-82 ± 2*	4.6 ± 2*	-202 ± 5	-8.7 ± 0.6	-47.5 ± 2.4	7.2 ± 1	-82.7 ± 3	5.7 ± 1	-203 ± 5	-0.98 ± 0.6		
SUMO1	-51 ± 3*	5.3 ± 1*	-82 ± 3*	4.2 ± 2*	-2.22 ± 1	-50.6 ± 3*	4.5 ± 2*	-82.5 ± 3*	4.1 ± 1*	-198 ± 5	-9.1 ± 0.5	-46.5 ± 3	7.3 ± 2	-83.5 ± 3.7	6 ± 3	-201 ± 4	-0.94 ± 0.5		

Human cardiac myocytes derived from iPS cells (Figure 1) or cloned channels in CHO-K1 cells (Figure 3) were studied in whole-cell mode. Stimulation protocols are described in the Methods. $V_{1/2}$, the voltage evoking half-maximal conductance and k , for activation or steady-state inactivation (SSI), were obtained by fitting the normalized current plotted against voltage to a Boltzmann function, $I = I_{max}/(1 + \exp[-(V - V_{1/2})/k])$, where I_{max} is maximum current. For comparison between groups, current densities were measured at -30 mV, the potential that gave peak current in all conditions studied. Data are means ± SEM for 8 to 12 cells per group. Changes from values measured from cells studied at 21% oxygen O₂ under control conditions were assessed by ANOVA and are indicated with *p < 0.05.

Of note, the kinetics of recovery of I_{Na} from the fast-inactivated state was insensitive to hypoxia, SUMO1, and SENP1 (Figure S1).

SUMOylation of Nav1.5 on Lys442 Is Necessary and Sufficient to Recapitulate the Hypoxic Response

In cardiomyocytes, Nav1.5 channel complexes contain the 2016 amino-acid-pore-forming subunit (Figure 2A) and an accessory subunit, Navβ1 (Catterall, 2012). A combination of spectroscopic and electrophysiological assays in live CHO cells showed that SUMOylation in response to hypoxia occurred only on Nav1.5 K442 and that modification of the site mediated the response to acute hypoxia.

First, the association of heterologously expressed human Nav1.5 and SUMO proteins at the surface of cells was studied by measuring FRET between subunits tagged with the fluorescent proteins mTFP1 and YFP (Figure 2A). The Navβ1 subunit was expressed in these studies but did not carry a fluorescent protein. Tagging the N terminus of Nav1.5 (mTFP-Nav1.5) did not alter the biophysical properties of the channel nor its responsiveness to hypoxia or SUMO1 or SENP1 in the recording pipette (Figures S2; Table S1). Similarly, tagging the N terminus of SUMO1 (YFP-SUMO1) did not alter its operation with Nav1.5, as previously observed with Nav1.2 and three K⁺ channels (Xiong et al., 2017; Plant et al., 2010, 2011, 2012, 2016).

Both mTFP1-Nav1.5 and YFP-SUMO1 were visualized at the plasma membrane when expressed in CHO cells (Figure 2A). Indicating FRET, and thus the intimate association of the two fluorophores consistent with SUMOylation, the time course of mTFP1 photobleaching under continuous illumination had a time constant (τ) that was prolonged from less than 10 s to 27 s when the channel was expressed with YFP-SUMO1 (Figures 2B and 2C). mTFP-Nav1.5 also showed FRET with YFP-Ubc9, the SUMO conjugating enzyme that binds to target proteins only when a lysine subject to modification is present (Xiong et al., 2017; Plant et al., 2016). In contrast, FRET was not observed when mTFP-Nav1.5 was expressed with linkage-incompetent YFP-SUMO1₉₅ (a variant that lacks the terminal Gly-Gly motif present in SUMO1 and the matured isoform SUMO1₉₇) or soluble YFP; in both cases, τ was less than 10 s.

To identify SUMOylation site(s) in Nav1.5 channels, we analyzed the primary sequence of the subunit with an algorithm that looks for variants of the canonical SUMO motif ψ -K-X-E/D, as before with Nav1.2 and the K⁺ channels (Xiong et al., 2017; Plant et al., 2010, 2011, 2012, 2016). We mutated each of five potential target lysines individually to glutamine in mTFP-Nav1.5 and used FRET with YFP-SUMO1 to assess interaction (Figure S2). Only K442, within the motif L-K₄₄₂-K-E, was required to produce FRET. Thus, mTFP-Nav1.5-K442Q was expressed on the cell surface like wild-type mTFP1-Nav1.5 but did not show FRET with YFP-SUMO1 or YFP-Ubc9 (Figure 2C). The absence of FRET supported the conclusion that K442 was necessary for SUMOylation of the Nav1.5 channel complex and, moreover, that no other lysine residue in the α subunit or in the Navβ1 was subject to SUMOylation.




Figure 2. FRET between Nav1.5 and SUMO1 at the Cell Surface Requires K442

Human Nav1.5 was expressed with Nav β 1 and studied in live CHO-K1 cells per the STAR Methods. FRET was assessed by measuring the time constant (τ) for mTFP photobleaching (donor) in the presence of YFP (acceptor) from 3 regions of 5–7 cells per group. Data are mean $\tau \pm$ SEM, and significant changes in τ compared to free YFP are indicated (* $p < 0.001$, two-way ANOVA).

(A) Top: a cartoon showing the secondary structure of a mTFP-Na_v1.5 subunit and the location of putative SUMOylation sites identified by the SUMOplot algorithm. The Na_v1.5 α -subunit has four homologous domains (I–IV), each with six transmembrane segments that fold to form one central ion conduction pore. Domains I and II are critical to opening the pore (activation) (Catterall, 2017; Cestèle et al., 1998, 2001). Fast inactivation results from pore occlusion by an IFM triplet of residues in the linker between domains III and IV (Vassilev et al., 1988; West et al., 1992; Cha et al., 1999). Bottom: fluorescent photomicrographs showing that mTFP-Na_v1.5 (blue) and YFP-SUMO1 (yellow) reach the cell surface. Scale bar represents 10 μ m. The montage shows photobleaching of mTFP-Na_v1.5 in response to continued excitation.

(B) Exemplar photobleaching studies show the decay of mTFP fluorescence intensity for single cells expressing mTFP-Na_v1.5 (black circle) or mTFP-Na_v1.5-K442Q (green circle) with YFP-SUMO1, fit by an exponential to give τ .

(C) A histogram of mean FRET τ values show the assembly of mTFP-Na_v1.5 (black bars) with YFP-SUMO1 and YFP-Ubc9 ($\tau = 28 \pm 3^*$ and $26 \pm 3.5^*$,

respectively) but not with linkage-incompetent YFP-SUMO1₉₅ or free YFP ($\tau = 9 \pm 2$ and 9 ± 2 , respectively). In contrast, mTFP-Na_v1.5-K442Q (green bars) did not show FRET with YFP-SUMO1, YFP-SUMO1₉₅, YFP-Ubc9, or free YFP ($\tau = 10 \pm 2$; 9.5 ± 2 ; 10 ± 2 ; and 9 ± 2 , respectively).

To confirm that K442 in the Nav1.5 sequence motif was subject to SUMOylation, a mass spectrometry (MS) strategy was used that we had previously developed for K⁺ channels (Plant et al., 2010, 2011). SUMO1₉₇T95K, a variant that leaves a Gly-Gly remnant of SUMO1 on the ϵ amino group of the target Lys after enzyme cleavage, was expressed with Nav1.5 residues 353–502 bearing a His₆ affinity tag and the mammalian E1 and E2 SUMOylation enzymes in bacteria. Affinity purification followed by SDS-PAGE revealed a band of the expected apparent mass for the Nav1.5-SUMO adduct (~35 kDa). The purified protein was treated with trypsin and subjected to MS to yield a sequence coverage for Nav1.5_{353–502} of 42% and was identified uniquely as the fragment carrying the Gly-Gly SUMO1 remnant (Figure S4).

Next, we confirmed that K442 controlled the response to hypoxia by studying human Nav1.5 channels expressed in CHO cells. With the wild-type channel, decreasing O₂ from 21% to 1.5% was associated with a rapid increase in I_{LATE} from ~0.5% (-0.96 ± 0.2 pA/pF) to ~4.1% (-8.3 ± 0.6 pA/pF), without a change in the peak current, ~200 pA/pF at -30 mV (Figure 3A; Table 1), as observed for I_{LATE} in iPS-CMs. As predicted, Nav1.5-K442Q channels were insensitive to hypoxia; I_{LATE} for the point

mutant channels remained less than ~0.5% of the peak current when O₂ was decreased to 1.5%. Furthermore, peak current measured at -30 mV was like wild-type channels (~200 pA/pF) and was also unaltered by hypoxia (Figures 3A and 3B). Of note, the time course for the change in I_{LATE} for Nav1.5 channels heterologously expressed in the CHO cells was like that for native I_{Na} in iPS-CMs, increasing from 0.5 to 4.2% over ~100 s, with half the rise time of 39.7 ± 2 s, and the increase was stable after the cells were restored to ambient levels of O₂ for at least 3 min (Figure 3C).

Nav1.5 channels heterologously expressed in tissue culture cells also responded to SUMO1 and SENP1 like I_{Na} in iPS-CMs. Thus, Nav1.5 channels with SUMO1 in the recording pipette showed I_{LATE} of ~4% of the peak current at both 21% and 1.5% O₂ (Figure 3C; Table 1). Furthermore, the window-current voltages increased in 1.5% O₂, or with 1 nM SUMO1, by -6 mV for $V_{1/2-act}$ and $+6$ mV for $V_{1/2-SSI}$ (Figure S3), as observed for I_{Na} in iPS-CMs. Also like I_{Na} in iPS-CMs, I_{LATE} for Nav1.5 channels was below 0.5%, with 1 nM SENP1 in the pipette at both 21% and 1.5% O₂, with no change in the window current, $V_{1/2-act}$, or $V_{1/2-SSI}$. Finally, the biophysical properties of Nav1.5-K442Q channels were insensitive to hypoxia and




Figure 3. K442 Is Required for Hypoxia and SUMO1-Mediated I_{LATE}

Human $Na_v1.5$ wild-type (WT) or $Na_v1.5$ -K442Q was expressed in CHO-K1 cells with $Na_v\beta1$ and studied under normoxic (21%) and hypoxic (1.5%) O_2 conditions with control solution (black), 1 nM SUMO1 (red), or 1 nM SENP1 (blue) in the recording pipette, as indicated. The late current (I_{LATE}) was studied at -30 mV and is shown as a mean percentage of the peak current remaining between 50 and 100 ms after the peak current. The time course of hypoxic modulation of I_{Na} was studied by steps from -100 mV to -30 mV every 10 s. Data are mean \pm SEM for 8 to 12 cells per group. Single-channel currents were studied in cell-attached mode and elicited every 5 s by a 50-ms depolarizing pulse to -30 mV from a holding potential of -120 mV. Data were recorded at filter and sampling frequencies of 5 and 50 kHz, respectively, and processed offline using a 1.2-kHz Bessel filter for display. Recording pipettes were filled with extracellular buffer. For each cell, null sweeps (with no channel activity) were identified, averaged offline, and subtracted from the data sweeps before analysis.

(A) Example traces from a cell expressing WT $Na_v1.5$ channel activation at -30 mV; I_{LATE} increased when control perfusate at 21% O_2 (black) was exchanged with a hypoxic solution at 1.5% O_2 (black dash). Inset: the increase in I_{LATE} shown in response to hypoxia.

(B) Example traces showing that hypoxia does not induce an increase in I_{LATE} in cells expressing $Na_v1.5$ -K442Q channels.

(C) Left: the time course for changes in I_{LATE} in response to hypoxia and on return to normoxia for cells expressing WT $Na_v1.5$ (closed circles) or $Na_v1.5$ -K442Q channels (open circles) with control solution (black), SUMO1 (red), or SENP1 (blue) in the recording pipette. Right: a histogram showing the mean I_{LATE} as a percentage of the peak current from cells studied at 21% O_2 , after 200 s at 1.5% O_2 and after 200 s following a return to 21% O_2 , with the pipette solutions indicated.

(D) Top: upon depolarization, single $Na_v1.5$ channels opened, inactivated rapidly, and did not

reopen. $Na_v1.5$ channels reopened upon exposure to 1.5% O_2 . Null traces in normoxia and hypoxia were $55\% \pm 5\%$ ($n = 20$ patches; 2,000 sweeps) and $53\% \pm 4\%$ ($n = 20$ patches; 2,000 sweeps), respectively. Bottom: ensemble average traces ($n = 200$ –220 sweeps) for the condition indicated.

(E) Top: single $Na_v1.5$ -K442Q channels did not reopen in ambient or hypoxic conditions. Null traces were $51\% \pm 3\%$ ($n = 20$ patches; 2,000 sweeps) and $50\% \pm 2\%$ ($n = 20$ patches; 2,000 sweeps), respectively. Bottom: ensemble average traces ($n = 200$ –225 sweeps) for the condition indicated.

were not altered by SUMO1 or SENP1, showing less than 0.5% I_{LATE} and without changes in $V_{1/2-act}$ and $V_{1/2-SSI}$.

Single $Na_v1.5$ Channels Are Reopened from the Inactive State by Hypoxia

Single $Na_v1.5$ channels are closed at rest, open briefly upon depolarization, and then inactivate rapidly; the channels remain in the inactive state until the membrane returns to the resting potential and the channels move back to the closed state from which they are available to open once again in response to the next heartbeat. Thus, single $Na_v1.5$ channels in cell-attached patches open just once in response to depolarization and almost always remain inactive (Figure 3D). As predicted by their behavior in whole-cell mode, hypoxia induced abnormal sin-

gle-channel reopenings during sustained depolarization. Ensemble currents generated by the cumulative averaging of 300 repetitive cycles of depolarization demonstrated that the reopening of inactivated $Na_v1.5$ channels in response to hypoxia was sufficient to increase I_{LATE} from $0.4\% \pm 0.1\%$ to $4.8\% \pm 0.3\%$. In contrast, $Na_v1.5$ -K442Q channels opened only once and remained inactive when cells were studied in either ambient levels of O_2 or at 1.5% O_2 (Figure 3E).

Single $Na_v1.5$ Channels Are Reopened from the Inactive State by SUMOylation

To confirm the mechanistic role of SUMO, single $Na_v1.5$ channels were studied in inside-out membrane patches excised from CHO cells, allowing perfusion of the intracellular face of




Figure 4. Hypoxia Recruits One SUMO1 Monomer to Each Cell Surface $\text{Na}_v1.5$ Channel

Single mTFP- $\text{Na}_v1.5$ channels (blue) and SUMO1 tagged with mCherry (m-SUMO1, red) were studied in CHO-K1 cells by TIRFM. Stoichiometric (photobleaching) and pixel-by-pixel analysis for subunit density and co-localization were performed as described in the [STAR Methods](#). Manders' coefficients were assessed post hoc for 3–5 regions per cell, and co-localization was defined as the presence of both fluorophores at more than 30% of the maximum fluorescence level recorded in that stack (and their overlap is represented in the images as white pixels). The time course of hypoxic modulation of $\text{Na}_v1.5 I_{LATE}$ was studied with steps from -100 mV to -30 mV every 10 s and normalized to the peak current. Data represent 5–8 cells, and biophysical parameters and single particle statistical analyses are summarized in [Tables S1](#) and [S3](#).

(A) Left: single co-localized particles with mCherry and TFP fluorescence were observed at the surface of cells expressing $\text{Na}_v1.5$ and SUMO1. The time courses for simultaneous photobleaching of the fluorophores revealed that complexes have one subunit of each type.

(B) Histogram of photobleaching steps showing that hypoxia increased single m-SUMO1 (red) subunits at the cell surface co-localized with mTFP- $\text{Na}_v1.5$ channels (blue), without a change in subunit stoichiometry.

(legend continued on next page)

the channel with experimental solutions. In ambient O_2 , single $Na_v1.5$ channels opened once and remained inactive for the duration of the depolarizing step, and in this configuration reducing O_2 from 21% to 1.5% did not produce channel reopening (Figure S5). This suggested that hypoxia-induced reopening in whole-cell mode depended on the presence of cytoplasmic SUMO, a hypothesis confirmed when patches were perfused with 1 nM SUMO1, and reopening was observed at both 21% and 1.5% O_2 .

Like the changes observed in whole-cell mode with hypoxia, single channels in the patches reopened within 150 s of exposure to SUMO1, with 50% of the patches showing channel reopening at 65 ± 10 s (Figure S5). In agreement with the cell-attached recordings, ensemble currents generated from single $Na_v1.5$ channels by averaging 500 cycles of depolarization in off-cell patches showed that SUMO1 increased I_{LATE} from $0.4\% \pm 0.2\%$ to $4.1\% \pm 0.3\%$ (Figure S5). As anticipated, single $Na_v1.5$ -K442Q channels in inside-out patches opened once and did not reopen even when exposed to SUMO1 or hypoxia (Figure S5). When considered in conjunction with the cell-attached studies, these findings support three conclusions: SUMOylation is necessary and sufficient for aberrant reopening of single $Na_v1.5$ channels, K442 is required to mediate the effect of SUMO1 on $Na_v1.5$, and hypoxia-evoked changes in I_{LATE} are mediated by the SUMO pathway.

Hypoxia Induces Rapid monoSUMOylation of $Na_v1.5$ Channels

To determine the stoichiometry of SUMOylation, we directly counted the number of SUMO1 monomers on individual $Na_v1.5$ channels at the surface of live CHO cells by using TIRFM photobleaching, as before for $Na_v1.2$ and I_{Ks} channels (Xiong et al., 2017; Plant et al., 2014, 2016). When mTFP- $Na_v1.5$ and SUMO1 tagged with mCherry (m-SUMO1) were co-expressed, particles containing only mTFP- $Na_v1.5$ and others with both mTFP- $Na_v1.5$ and m-SUMO1 were observed at the cell membrane (Figures 4A and 4B; Table 2). Approximately 10% of mTFP- $Na_v1.5$ channels were co-localized with one m-SUMO1 in ambient oxygen: continuous, two-color excitation bleached each fluorophore in the complexes in a single step consistent with one subunit of each type. On exposure to hypoxia, m-SUMO1 was recruited to the cell surface but only at sites with mTFP- $Na_v1.5$, increasing the number of pixels that contained both mTFP- $Na_v1.5$ and m-SUMO1 channels from 10% to 87% within ~ 100 s (Figure 4B). Increased co-localization did not alter the 1:1 stoichiometry of $Na_v1.5$ channel and m-SUMO1 complexes, as judged by bleaching.

Evidence that the co-localization represented a covalent interaction of the channel and SUMO1 was provided by two additional experiments (Figure S6). First, m-SUMO1 was not observed at the membrane in cells expressing mTFP- $Na_v1.5$ -K442Q and rarely yielded co-localization of the fluorophores in 21% or 1.5% O_2 . Second, hypoxia did not increase the number of particles at the membrane with m-SUMO1₉₅, the SUMO1 variant that is unable to covalently attach to target lysines. Because $Na_v1.5$ channels contain just one α -subunit, the observed 1:1 stoichiometry with SUMO1 supports three additional conclusions: $Na_v1.5$ subunits are SUMOylated only on K442, the channel is monoSUMOylated and polySUMO1 chains were not observed, and the $Na_v\beta 1$ subunit is not SUMOylated.

Although hypoxia recruited m-SUMO1 to the cell surface, it did not change the number of $Na_v1.5$ channels at the plasma membrane (Figures 4C and 4D; Table 2). This was shown by studying the surface density and localization of mTFP- $Na_v1.5$ and m-SUMO1 subunits by using pixel-by-pixel analysis, a method we have applied before (Xiong et al., 2017; Plant et al., 2014, 2016). In ambient O_2 , about 10% of mTFP- $Na_v1.5$ fluorescent pixels were localized with m-SUMO1 ($33 \pm 3/\mu m^2$) and 90% of the channels were unmodified ($305 \pm 21/\mu m^2$). After 100 s of hypoxia, the ratio was reversed; the number of mTFP- $Na_v1.5$ pixels co-localized with m-SUMO1 increased to about 90% ($301 \pm 13/\mu m^2$), and 10% of the channels remained unmodified by m-SUMO1 ($37 \pm 4/\mu m^2$). Thus, hypoxia increased the surface density of m-SUMO1 ~ 33 -fold (from $10/\mu m^2$ to $334/\mu m^2$), whereas the density of mTFP- $Na_v1.5$ did not change ($338/\mu m^2$).

Consistent with cause and effect, the appearance of m-SUMO1 at the plasma membrane with mTFP- $Na_v1.5$, demonstrated above to require covalent modification of the channel on K442, matched the time course of hypoxia-induced increase in I_{LATE} observed in iPS-CMs and $Na_v1.5$ channels studied in CHO cells (Figures 1 and 3; Table 1). These rates also matched the increase in the mean Manders' coefficient of co-localization of m-SUMO1 and mTFP- $Na_v1.5$ (Figure 4E) as well as Pearson's correlation coefficient (Table S2). Also, as observed for I_{LATE} in iPS-CMs and $Na_v1.5$ channels in CHO cells, the number of mTFP- $Na_v1.5$ surface particles and the fraction of channels localized with m-SUMO1 were stable on restoration to ambient O_2 for 3 min after hypoxia (Figure 4E; Table 2). Supporting the notion that heterologous expression of m-SUMO1 increased basal SUMOylation of mTFP- $Na_v1.5$, about 10% of mTFP- $Na_v1.5$ channels co-localized with m-SUMO1 in these cells and I_{LATE} was 2.5-fold higher ($\sim 1\%$ of peak current) than the control in ambient O_2 , rising to 4.5% with hypoxia (Tables 1 and S1).

(C) Left: the images show that in ambient O_2 , the surface density of m-SUMO1 (top) is low compared to $Na_v1.5$ (middle) with little co-localization (bottom). Right: hypoxia recruits m-SUMO1 to the cell surface within ~ 100 s (top), where it is co-localized with $Na_v1.5$ channels (bottom); surface levels of $Na_v1.5$ were not observed to change (middle). The scale bar represents 10 μm .

(D) Histogram of surface density summarizing six cells studied as described in (A). In ambient O_2 , the density of pixels per μm^2 with SUMO1 alone (red) was 3 ± 1 , and 305 ± 21 for $Na_v1.5$ (blue). The density of pixels with both fluorophores (green) was 33 ± 3 per μm^2 . Hypoxia increased co-localization to 301 ± 13 pixels per μm^2 and decreased the density of free $Na_v1.5$ channels (37 ± 4) without altering the density of free SUMO1 (6 ± 2).

(E) The time course for the hypoxia-induced increase in the co-localization of mTFP- $Na_v1.5$ and m-SUMO1 (Manders' coefficient, black circle) and the magnitude of I_{LATE} , as a percentage of the peak current (open circle) were coincident. A mean Manders' coefficient of 0.10 ± 0.01 measured in ambient O_2 increased to 0.91 ± 0.03 in ~ 100 s of exposure to hypoxia. The mean I_{LATE} rose from $0.46\% \pm 0.1\%$ to $4.4\% \pm 0.8\%$. Increases in the Manders' coefficient and late current versus peak current ratio were unchanged for at least 3 min after cells were restored to ambient O_2 .

Table 2. Co-localization of SUMO1 with Na_v1.5 in Response to Hypoxia

Subunits Expressed	mTFP-Na _v 1.5 + m-SUMO1				mTFP-Na _v 1.5-K442Q + m-SUMO1			
	Ambient O ₂	Hypoxia 50 s	Hypoxia 100 s	Recovery 150 s	Ambient O ₂	Hypoxia 50 s	Hypoxia 100 s	Recovery 150 s
Condition	O ₂	50 s	100 s	150 s	O ₂	50 s	100 s	150 s
Single particle stoichiometry SUMO1: Na _v 1.5	1: 1	1: 1	1: 1	1: 1	0: 1	0: 1	0: 1	0: 1
Free mTFP-Na _v 1.5, pixels/μm ²	305 ± 21	166 ± 18*	37 ± 4**	39 ± 5**	328 ± 26	333 ± 29	326 ± 25	308 ± 19
Free m-SUMO1, pixels/μm ²	3 ± 1	4 ± 2	7 ± 2	5 ± 2	3 ± 1	4 ± 1	4 ± 2	6 ± 2
Co-localized pixels/μm ²	33 ± 3	179 ± 15**	301 ± 13**	302 ± 15**	4 ± 1	4 ± 2	3 ± 1	4 ± 1

Na_v1.5 or Na_v1.5-K442Q subunits tagged with mTFP1 were expressed in CHO-K1 cells with mCherry-SUMO1 (m-SUMO1) and studied by TIRFM and whole-cell patch clamp (Figures 3 and 5). The number of photobleaching steps observed for each fluorophore in each single fluorescent spot reports on the stoichiometry of the channel complex. Na_v1.5 channels are monomers and show no more than one bleaching step when tagged with mTFP1 (Figure 5). No more than one bleaching steps was observed for mCherry-tagged SUMO1 subunits (free or co-localized with the channel). A 1:1 stoichiometry is maintained when cells are exposed to hypoxia (1.5% O₂). SUMO1 was not observed to co-localize with Na_v1.5-K442Q channels. The surface density of subunits was quantified as the mean of four 100- by 100-pixel regions for 6–10 cells per group. Exposure to hypoxia increased the number of SUMO1 monomers observed at the cell surface within 100 s, and almost all were co-localized with Na_v1.5. Data are mean ± SEM for 5 to 8 cells per group; * p < 0.05, **p < 0.01 compared with cells studied in ambient O₂ for each channel type.

Hypoxia-Induced I_{LATE} in iPS-CMs Yields SUMO-Dependent APD Prolongation

To evaluate if the increase in I_{LATE} we observed was sufficient to produce pro-arrhythmic APD prolongation, iPS-CMs were studied in current-clamp mode and the time to repolarize the membrane to resting values during spontaneous action potentials was measured. Under control conditions, the cells had a resting potential of −45 ± 3 mV, and the phase 1 action potential upstroke mediated by I_{Na} peaked at 43 ± 4 mV (n = 6). Hypoxia increased the time for 50% restoration (APD₅₀) by 22% without changing the resting potential or the action potential height (Figure 5A). To confirm that action potential prolongation was due to an increase in I_{LATE}, we applied ranolazine, an open-state blocker of Na_v1.5 channels in clinical use that preferentially inhibits I_{LATE} rather than peak I_{Na} (Maier and Sossalla, 2013). Ranolazine reversed hypoxia-induced lengthening of APD₅₀ to within 5% of control values without altering other action potential biophysical parameters (Table S3).

To verify that SUMOylation mediates hypoxia-induced APD prolongation, 1 nM SUMO1 or SENP1 was included in the recording pipette. As expected, SUMO1 lengthened APD₅₀ to 123% of control values without causing a significant shift in the resting membrane potential or the height of spontaneous action potentials in 21% O₂, and hypoxia evoked only a further 5% increase in the APD₅₀ (Figure 5B). Subsequent application of ranolazine reversed the effects of SUMOylation as it had the effects of hypoxia with control solution in the pipette, reducing APD₅₀ to within 5% of its control value. In contrast, 1 nM SENP1 in the recording pipette yielded APD₅₀, resting membrane potential, and action potential height like control cells studied in 21% O₂, and neither hypoxia nor ranolazine altered the three parameters of excitability (Figure 5C). These observations support the conclusion that hypoxia-induced changes in APD are primarily due to the SUMO-mediated increase in I_{LATE} passed by Na_v1.5 channels. Of note, baseline APD in iPS-CMs is longer than that in native cardiomyocytes, apparently due to a lower expression of K⁺ channels (Yang et al., 2014).

The O'Hara-Rudy model for human cardiac action potentials was used to demonstrate that the increase in I_{LATE} due to

SUMOylation of Na_v1.5 channels observed in reconstituted CHO cells and iPS-CMs was sufficient to meet the theoretical requirements for pro-arrhythmic prolongation of APD. As anticipated, the time to repolarize the membrane from peak to rest by 50% (APD₅₀) was prolonged by 27% when I_{LATE} was increased in the model by 5-fold, half the observed maximal effect in the cells, from 205 ms under control conditions to 260 ms (Figure 5D); also as observed in iPS-CMs, there was no appreciable change in the resting membrane potential or the action potential peak current in the model (Table S3).

DISCUSSION

Myocardial hypoxia increases I_{LATE} leading to increased APD, a common setting for dangerous arrhythmias. Here, we show that hypoxia induces SUMOylation of Na_v1.5 channels on K442 and this is necessary and sufficient to increase I_{LATE} and APD in human cardiomyocytes. Supporting a direct mechanism, SUMOylation of Na_v1.5 produces I_{LATE} in the absence of hypoxia in membrane patches excised from cells, basal levels of Na_v1.5 SUMOylation are low under control conditions in iPS-CMs, and hypoxia leads to SUMOylation of the channels with the same time course as the increase in I_{LATE}. Furthermore, SUMOylation produces changes in the gating of single Na_v1.5 ion channels, macroscopic I_{Na}, and APDs like those described in human heart and in animal models with hypoxia (Chadda et al., 2017; Austen et al., 1963; Thung et al., 1962; Brown et al., 2014); and the effects are suppressed by the mutation of Na_v1.5 K442, the application of SENP1 deSUMOylase, or the application of ranolazine, an inhibitor of I_{LATE}.

Increased I_{LATE} and AP prolongation predispose to arrhythmia because they increase the influx of Na⁺, raising the intracellular Na⁺ concentration and depolarizing the membrane potential. These changes impact the activity of other ion channels and transporters, including the Na/K-ATPase, leading to Ca²⁺ loading by L-type Ca²⁺ channel reactivation and reverse operation of the sodium-calcium exchanger (NCX). These excitatory effects increase the likelihood of early after-depolarizations (EADs), delayed after-depolarizations (DADs), triggered activity, and




Figure 5. Hypoxia-Induced Increase in APD Mediated by SUMO Modulation of I_{LATE}

Spontaneous action potentials from human iPS-CMs were recorded in current-clamp mode with the same sequence of perfusates: normoxic (21% O₂, solid line), hypoxic (1.5% O₂, dashed line), and then hypoxia with 1 μ M ranolazine (purple). Action potentials were studied with control intracellular solution (black) or with 1 nM SUMO1 (red) or SENP1 (blue) in the recording pipette. APD₅₀ was determined by calculating the time required for the membrane potential to return to 50% of the resting value from the peak deflection of the action potential. Hypoxia was induced as described in the STAR Methods. Data are means \pm SEM for 6–9 cells studied per group (* p < 0.01, two-way paired t test). The O’Hara-Rudy model for human action potentials was applied as described in the STAR Methods and Table S3. Scale bars represent 25 mV and 250 ms in (A)–(C) and 30 mV and 150 ms in panel (D).

(A) Exemplar spontaneous action potentials recorded from iPS-CMs studied with control pipette solution under normoxic conditions and hypoxia with and without ranolazine. The APD₅₀ increased with hypoxia by 22%, from 663 \pm 2 ms to 810 \pm 2 ms, and application of ranolazine restored the APD₅₀ to 689 \pm 2 ms. (B) When SUMO1 was included in the pipette solution, APD₅₀ was 822 \pm 2 ms, and this increased to 870 \pm 2 ms with hypoxia; ranolazine with hypoxia produced an APD₅₀ of 699 \pm 3 ms like that observed without SUMO1 in the pipette under normoxic conditions.

(C) APD₅₀ was 663 \pm 2 ms when cells were studied with SENP1 in the recording pipette, and the APD₅₀ increased only to 698 \pm 2 ms with hypoxia and was returned to 663 \pm 4 ms with ranolazine.

(D) When a 5-fold increase in I_{LATE} (half the maximal effect observed with hypoxia/SUMOylation), was simulated using the O’Hara-Rudy model, the APD₅₀ increased by 27%, from 205 ms in normoxia to 260 ms.

(E) Histograms summarizing the mean APD₅₀ of iPS-CMs under the conditions described in (A)–(C).

spatiotemporal changes in repolarization that promote reentrant arrhythmias, including ventricular TdP (Gaur et al., 2009; Shryock et al., 2013; Chadda et al., 2017). Medications like ranolazine that suppress I_{LATE} minimize disruption of Ca²⁺ homeostasis, reducing EADs and DADs (Maier and Sossalla, 2013). The hypoxic increase in I_{LATE} we observe with iPS-CMs modeled in the O’Hara-Rudy algorithm are found sufficient to underlie the increase in APD that induces these downstream effects (Figure 5). In addition to hypoxia and ischemia (Saint, 2006; Hammarström and Gage, 2002), cardiac I_{LATE} is reported to increase with oxidative stress (Ward and Giles, 1997), acidosis (Plant et al., 2006), heart failure (Valdivia et al., 2005; Undrovinas et al., 2006), and inflammation (Ward et al., 2006), suggesting a role for SUMOylation in the transduction of these stimuli that merits study.

Rapid SUMOylation of Na_v1.5 channels on the cardiac plasma membrane in response to hypoxia is reminiscent of our finding that hypoxia mediates SUMOylation of Na_v1.2 channels on the surface of central neurons in under 60 s on K38, leading to an increase peak current (without changes in I_{LATE}) due to changes in the voltage dependence of activation, consistent with early, rapid Na⁺ flux in cerebral ischemia, a prelude to downstream pathology (Plant et al., 2016). In both cases, rapid recruitment of SUMO in the first few minutes of hypoxia initiates deleterious sequelae. In contrast, chronic hypoxia has been associated with cardioprotection by increased SUMOylation of the ATPase SERCA2A in heart failure and ischemia-reperfusion injury (Du et al., 2018) and with neuroprotection by a global increase in SUMOylation, changes in gene expression, and remodeling of protein expression in ground squirrel hibernation torpor (Kho et al., 2011, 2015; Lee et al., 2007) and chronic oxygen-glucose deprivation (Lee et al., 2009).

Na_v1.5 is recognized as the preeminent sodium channel in the heart, although other voltage-gated sodium channels are expressed at lower levels and in specific cardiac regions (Maier et al., 2004), notably purkinje fibers (Haufe et al., 2005). Although non-Na_v1.5 channels (Maier et al., 2004) have been described as contributing only 5%–10% of peak current, a role for hypoxia-induced I_{LATE} is possible because non-Na_v1.5 currents have been implicated in the late currents measured with ambient oxygen in isolated canine and murine myocytes (Biet et al., 2012; Yang et al., 2012). We do not infer roles for Na_v1.2, Na_v1.1, or Na_v1.6 in SUMO-mediated hypoxic late current because we have demonstrated that the stimuli alter peak current and do not induce I_{LATE} with Na_v1.2 in rat CGN and CHO cells and that Na_v1.1 and Na_v1.6 channels are insensitive to SUMOylation in rat hippocampal neurons and rat CGN, respectively (Plant et al., 2016). We cannot rule out contributions from Na_v1.3, Na_v1.4, Na_v1.7, Na_v1.8, or Na_v1.9.

At the single-channel level, the abnormal increases in I_{LATE} we observe with hypoxia are like those recorded with mutations in Na_v1.5 that cause LQT3 that destabilize internal pore occlusion by the IFM inactivation “lid” that resides between channel domains III and IV (Bennett et al., 1995; West et al., 1992). Rather than a single opening followed by inactivation, the LQT3 channels re-open with maintained depolarization, suggesting a lowered energetic barrier for return to the open state. Consistent with the idea that SUMOylated Na_v1.5 channels also experience potentials where the energetic barriers to moving between the

open and inactive states are lowered, we observe hypoxia to increase the window between activation and steady-state inactivation voltage curves (Figure S3). Furthermore, our failure to observe long first openings suggests that the barrier to inactivation is not increased by hypoxia. Of note, destabilization of the inactivated state can rationalize I_{LATE} in association with channels that recover from inactivation too rapidly during repolarization (Belardinelli et al., 2006, 2015; Shryock et al., 2013). It is feasible that hypoxia-induced SUMOylation of Na_v1.5 on K442, a site on the linker between domains I and II, may interfere with the stability of the interaction of the IFM lid and its pore receptor.

Here, we apply SUMO1₁₀₁ to the inner surface of excised membrane patches to SUMOylate Na_v1.5 channels, as we observed before for K2P1, K_v2.1, and Na_v1.2 channels; this indicates that enzymes stably associated with the cytoplasmic face of the plasma membrane mature SUMO1 to SUMO₉₇ and mediate coupling to target lysine (Xiong et al., 2017; Plant et al., 2010, 2011, 2012, 2016). In contrast, when SUMOylation of Na_v1.5 channels is induced by hypoxia, the membrane must be in continuity with the cytosol, indicating cellular components have a role in transducing the signal that oxygen tension is low to the SUMO pathway. Thus far, exogenous SENP1 must be applied to deSUMOylate the ion channels in off-cell mode in all cases we have studied.

It is notable that changes in Ca²⁺ homeostasis are not required for hypoxia-induced SUMOylation of Na_v1.5 in whole cells as the reaction proceeds despite 10 mM EGTA in the recording pipette to suppress changes in cellular Ca²⁺ and 200 μM CdCl₂ in the extracellular bath solution to suppress conductance through voltage-gated calcium channels (Figure 1A). The similar attributes of SUMOylated Na_v1.5 currents in intact cells and single channels in excised patches also support the conclusion that changes in Ca²⁺ at the cytoplasmic face of the channel are not necessary for SUMO-induced increases in I_{LATE} (Figure S5). Those points do not rule out a role for variation in cytoplasmic Ca²⁺ when the ionic concentration is not clamped by chelation and blockade.

Ion channel SUMOylation appears to enhance excitability by decreasing potassium currents (K2P1, K_v2.1, and Kv7.1/KCNE1) (Rajan et al., 2005; Plant et al., 2010, 2012) or increasing currents passed by Na_v1.2 (Plant et al., 2016) and Na_v1.5 (this work). Our prior study showing that acute hypoxia leads to the rapid SUMOylation of Na_v1.2 in CGN and influx of Na⁺ flux due to altered voltage-dependent gating (but not a change in I_{LATE}) (Plant et al., 2016) and this study of Na_v1.5 suggest that hypoxia-mediated SUMOylation is an emerging common mechanism shared by some, but not all, ion channels.

We observed previously that acidosis increases I_{LATE} in Na_v1.5 channels bearing a common polymorphism associated with SIDS and arrhythmia in African American adults, presenting a model for the genetic predisposition to dysrhythmia in the face of an environmental challenge (Plant et al., 2006). Inherited ion channel mutations that predispose to arrhythmia by increasing APD also yield life-threatening events in response to specific stimuli, for example, concurrent drug blockade (Sesti and Goldstein, 1998). Here, we present the mechanism for the sensitivity of wild-type Na_v1.5 channels to the environmental challenge of

hypoxia. Identifying the role of the SUMO pathway in the increase in I_{LATE} in response to hypoxia supports targeting the pathway for therapeutic intervention.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- KEY RESOURCES TABLE
- LEAD CONTACT AND MATERIALS AVAILABILITY
- EXPERIMENTAL MODEL AND SUBJECT DETAILS
- METHOD DETAILS
 - Molecular Biology and Reagents
- QUANTIFICATION AND STATISTICAL ANALYSIS
- DATA AND CODE AVAILABILITY

SUPPLEMENTAL INFORMATION

Supplemental Information can be found online at <https://doi.org/10.1016/j.celrep.2020.01.025>.

ACKNOWLEDGMENTS

The authors thank Anthony F. Arena, Jordie M. Kamuene, and Meghan Masotti for technical support. We are grateful to the late R.Y. Tsien (UCSD) for the mCherry clone and A. Periasamy (University of Virginia) for mTFP1. The work was funded by National Institutes of Health grants R01HL10549 (to S.A.N.G.) and R01HL144615 (to L.D.P.).

AUTHOR CONTRIBUTIONS

L.D.P., D.X., J.R., and H.D. performed the research and analyzed data; L.D.P. and S.A.N.G. designed the research and wrote the paper.

DECLARATION OF INTERESTS

The authors report no competing financial interests.

Received: May 13, 2019

Revised: December 3, 2019

Accepted: January 7, 2020

Published: February 11, 2020

REFERENCES

- Austen, W.G., Ebert, P.A., and Greenfield, L.J. (1963). Mechanism of cardiac arrest in acute hypoxia. *Surgery* 53, 784–791.
- Belardinelli, L., Shryock, J.C., and Fraser, H. (2006). Inhibition of the late sodium current as a potential cardioprotective principle: effects of the late sodium current inhibitor ranolazine. *Heart* 92, iv6–iv14.
- Belardinelli, L., Giles, W.R., Rajamani, S., Karagueuzian, H.S., and Shryock, J.C. (2015). Cardiac late Na⁺ current: proarrhythmic effects, roles in long QT syndromes, and pathological relationship to CaMKII and oxidative stress. *Heart Rhythm* 12, 440–448.
- Bennett, P.B., Yazawa, K., Makita, N., and George, A.L., Jr. (1995). Molecular mechanism for an inherited cardiac arrhythmia. *Nature* 376, 683–685.
- Biet, M., Barajas-Martínez, H., Ton, A.T., Delabre, J.F., Morin, N., and Dumaine, R. (2012). About half of the late sodium current in cardiac myocytes from dog ventricle is due to non-cardiac-type Na(+) channels. *J. Mol. Cell. Cardiol.* 53, 593–598.

- Brown, S.J., Barnes, M.J., and Mündel, T. (2014). Effects of hypoxia and hypercapnia on human HRV and respiratory sinus arrhythmia. *Acta Physiol. Hung.* *101*, 263–272.
- Carmeliet, E. (1999). Cardiac ionic currents and acute ischemia: from channels to arrhythmias. *Physiol. Rev.* *79*, 917–1017.
- Catterall, W.A. (1986). Molecular properties of voltage-sensitive sodium channels. *Annu. Rev. Biochem.* *55*, 953–985.
- Catterall, W.A. (2012). Voltage-gated sodium channels at 60: structure, function and pathophysiology. *J. Physiol.* *590*, 2577–2589.
- Catterall, W.A. (2017). Forty Years of Sodium Channels: Structure, Function, Pharmacology, and Epilepsy. *Neurochem. Res.* *42*, 2495–2504.
- Cestèle, S., Qu, Y., Rogers, J.C., Rochat, H., Scheuer, T., and Catterall, W.A. (1998). Voltage sensor-trapping: enhanced activation of sodium channels by beta-scorpion toxin bound to the S3-S4 loop in domain II. *Neuron* *21*, 919–931.
- Cestèle, S., Scheuer, T., Mantegazza, M., Rochat, H., and Catterall, W.A. (2001). Neutralization of gating charges in domain II of the sodium channel alpha subunit enhances voltage-sensor trapping by a beta-scorpion toxin. *J. Gen. Physiol.* *118*, 291–302.
- Cha, A., Ruben, P.C., George, A.L., Jr., Fujimoto, E., and Bezanilla, F. (1999). Voltage sensors in domains III and IV, but not I and II, are immobilized by Na⁺ channel fast inactivation. *Neuron* *22*, 73–87.
- Chadda, K.R., Jeevaratnam, K., Lei, M., and Huang, C.L. (2017). Sodium channel biophysics, late sodium current and genetic arrhythmic syndromes. *Pflugers Arch.* *469*, 629–641.
- Du, Y., Liu, P., Xu, T., Pan, D., Zhu, H., Zhai, N., Zhang, Y., and Li, D. (2018). Luteolin Modulates SERCA2a Leading to Attenuation of Myocardial Ischemia/Reperfusion Injury via Sumoylation at Lysine 585 in Mice. *Cell. Physiol. Biochem.* *45*, 883–898.
- Gaur, N., Rudy, Y., and Hool, L. (2009). Contributions of ion channel currents to ventricular action potential changes and induction of early afterdepolarizations during acute hypoxia. *Circ. Res.* *105*, 1196–1203.
- Hammarström, A.K., and Gage, P.W. (2002). Hypoxia and persistent sodium current. *Eur. Biophys. J.* *31*, 323–330.
- Haufe, V., Cordeiro, J.M., Zimmer, T., Wu, Y.S., Schiccitano, S., Benndorf, K., and Dumaine, R. (2005). Contribution of neuronal sodium channels to the cardiac fast sodium current I_{Na} is greater in dog heart Purkinje fibers than in ventricles. *Cardiovasc. Res.* *65*, 117–127.
- Henley, J.M., Craig, T.J., and Wilkinson, K.A. (2014). Neuronal SUMOylation: mechanisms, physiology, and roles in neuronal dysfunction. *Physiol. Rev.* *94*, 1249–1285.
- Ju, Y.K., Saint, D.A., and Gage, P.W. (1996). Hypoxia increases persistent sodium current in rat ventricular myocytes. *J. Physiol.* *497*, 337–347.
- Kho, C., Lee, A., Jeong, D., Oh, J.G., Chaanine, A.H., Kizana, E., Park, W.J., and Hajjar, R.J. (2011). SUMO1-dependent modulation of SERCA2a in heart failure. *Nature* *477*, 601–605.
- Kho, C., Lee, A., Jeong, D., Oh, J.G., Gorski, P.A., Fish, K., Sanchez, R., DeVita, R.J., Christensen, G., Dahl, R., and Hajjar, R.J. (2015). Small-molecule activation of SERCA2a SUMOylation for the treatment of heart failure. *Nat. Commun.* *6*, 7229.
- Lee, Y.J., Miyake, S., Wakita, H., McMullen, D.C., Azuma, Y., Auh, S., and Hallenbeck, J.M. (2007). Protein SUMOylation is massively increased in hibernation torpor and is critical for the cytoprotection provided by ischemic preconditioning and hypothermia in SHSY5Y cells. *J. Cereb. Blood Flow Metab.* *27*, 950–962.
- Lee, Y.J., Castri, P., Bembry, J., Maric, D., Auh, S., and Hallenbeck, J.M. (2009). SUMOylation participates in induction of ischemic tolerance. *J. Neurochem.* *109*, 257–267.
- Ma, J., Guo, L., Fiene, S.J., Anson, B.D., Thomson, J.A., Kamp, T.J., Kolaja, K.L., Swanson, B.J., and January, C.T. (2011). High purity human-induced pluripotent stem cell-derived cardiomyocytes: electrophysiological properties of action potentials and ionic currents. *Am. J. Physiol. Heart Circ. Physiol.* *301*, H2006–H2017.
- Maier, L.S., and Sossalla, S. (2013). The late Na current as a therapeutic target: where are we? *J. Mol. Cell. Cardiol.* *61*, 44–50.
- Maier, S.K., Westenbroek, R.E., McCormick, K.A., Curtis, R., Scheuer, T., and Catterall, W.A. (2004). Distinct subcellular localization of different sodium channel alpha and beta subunits in single ventricular myocytes from mouse heart. *Circulation* *109*, 1421–1427.
- Makielski, J.C. (2016). Late sodium current: A mechanism for angina, heart failure, and arrhythmia. *Trends Cardiovasc. Med.* *26*, 115–122.
- Marbán, E. (2002). Cardiac channelopathies. *Nature* *415*, 213–218.
- O'Hara, T., Virág, L., Varró, A., and Rudy, Y. (2011). Simulation of the undiseased human cardiac ventricular action potential: model formulation and experimental validation. *PLoS Comput. Biol.* *7*, e1002061.
- Pearson, H.A., Sutton, K.G., Scott, R.H., and Dolphin, A.C. (1993). Ca²⁺ currents in cerebellar granule neurones: role of internal Mg²⁺ in altering characteristics and antagonist effects. *Neuropharmacology* *32*, 1171–1183.
- Plant, L.D., Kemp, P.J., Peers, C., Henderson, Z., and Pearson, H.A. (2002). Hypoxic depolarization of cerebellar granule neurons by specific inhibition of TASK-1. *Stroke* *33*, 2324–2328.
- Plant, L.D., Bowers, P.N., Liu, Q., Morgan, T., Zhang, T., State, M.W., Chen, W., Kittles, R.A., and Goldstein, S.A. (2006). A common cardiac sodium channel variant associated with sudden infant death in African Americans, SCN5A S1103Y. *J. Clin. Invest.* *116*, 430–435.
- Plant, L.D., Dementieva, I.S., Kollewe, A., Olikara, S., Marks, J.D., and Goldstein, S.A. (2010). One SUMO is sufficient to silence the dimeric potassium channel K2P1. *Proc. Natl. Acad. Sci. USA* *107*, 10743–10748.
- Plant, L.D., Dowdell, E.J., Dementieva, I.S., Marks, J.D., and Goldstein, S.A. (2011). SUMO modification of cell surface Kv2.1 potassium channels regulates the activity of rat hippocampal neurons. *J. Gen. Physiol.* *137*, 441–454.
- Plant, L.D., Zuniga, L., Araki, D., Marks, J.D., and Goldstein, S.A. (2012). SUMOylation silences heterodimeric TASK potassium channels containing K2P1 subunits in cerebellar granule neurons. *Sci. Signal.* *5*, ra84.
- Plant, L.D., Xiong, D., Dai, H., and Goldstein, S.A. (2014). Individual IKs channels at the surface of mammalian cells contain two KCNE1 accessory subunits. *Proc. Natl. Acad. Sci. USA* *111*, E1438–E1446.
- Plant, L.D., Marks, J.D., and Goldstein, S.A. (2016). SUMOylation of Na_v1.2 channels mediates the early response to acute hypoxia in central neurons. *eLife* *5*, e20054.
- Rajan, S., Plant, L.D., Rabin, M.L., Butler, M.H., and Goldstein, S.A. (2005). SUMOylation silences the plasma membrane leak K⁺ channel K2P1. *Cell* *121*, 37–47.
- Saint, D.A. (2006). The role of the persistent Na⁽⁺⁾ current during cardiac ischemia and hypoxia. *J. Cardiovasc. Electrophysiol.* *17*, S96–S103.
- Saint, K.M., Abraham, S., MacLeod, B.A., McGough, J., Yoshida, N., and Walker, M.J. (1992). Ischemic but not reperfusion arrhythmias depend upon serum potassium concentration. *J. Mol. Cell. Cardiol.* *24*, 701–709.
- Schindelin, J., Argana-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibish, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-image analysis. *Nat. Methods* *9*, 676–682.
- Sesti, F., and Goldstein, S.A. (1998). Single-channel characteristics of wild-type IKs channels and channels formed with two minK mutants that cause long QT syndrome. *J. Gen. Physiol.* *112*, 651–663.
- Shryock, J.C., Song, Y., Rajamani, S., Antzelevitch, C., and Belardinelli, L. (2013). The arrhythmogenic consequences of increasing late I_{Na} in the cardiomyocyte. *Cardiovasc. Res.* *99*, 600–611.
- Song, Y., Shryock, J.C., Wagner, S., Maier, L.S., and Belardinelli, L. (2006). Blocking late sodium current reduces hydrogen peroxide-induced arrhythmogenic activity and contractile dysfunction. *J. Pharmacol. Exp. Ther.* *318*, 214–222.
- Thung, N., Dammann, J.F., Jr., Diaz-Perez, R., Thompson, W.M., Jr., Sanmarco, M., and Mehegan, C. (1962). Hypoxia as the cause of hemorrhage into the cardiac conduction system, arrhythmia, and sudden death. *J. Thorac. Cardiovasc. Surg.* *44*, 687–695.

- Undrovinas, A.I., Belardinelli, L., Undrovinas, N.A., and Sabbah, H.N. (2006). Ranolazine improves abnormal repolarization and contraction in left ventricular myocytes of dogs with heart failure by inhibiting late sodium current. *J. Cardiovasc. Electrophysiol.* *17*, S169–S177.
- Valdivia, C.R., Chu, W.W., Pu, J., Foell, J.D., Haworth, R.A., Wolff, M.R., Kamp, T.J., and Makielski, J.C. (2005). Increased late sodium current in myocytes from a canine heart failure model and from failing human heart. *J. Mol. Cell. Cardiol.* *38*, 475–483.
- Vassilev, P.M., Scheuer, T., and Catterall, W.A. (1988). Identification of an intracellular peptide segment involved in sodium channel inactivation. *Science* *241*, 1658–1661.
- Ward, C.A., and Giles, W.R. (1997). Ionic mechanism of the effects of hydrogen peroxide in rat ventricular myocytes. *J. Physiol.* *500*, 631–642.
- Ward, C.A., Bazzazi, H., Clark, R.B., Nygren, A., and Giles, W.R. (2006). Actions of emigrated neutrophils on Na(+) and K(+) currents in rat ventricular myocytes. *Prog. Biophys. Mol. Biol.* *90*, 249–269.
- Wei, X., Zhu, A., Zhang, Y., Yao, S., and Mao, W. (2016). Pre- and Delayed Treatments With Ranolazine Ameliorate Ventricular Arrhythmias and Nav1.5 Downregulation in Ischemic/Reperused Rat Hearts. *J. Cardiovasc. Pharmacol.* *68*, 269–279.
- West, J.B. (2017). Physiological Effects of Chronic Hypoxia. *N. Engl. J. Med.* *376*, 1965–1971.
- West, J.W., Patton, D.E., Scheuer, T., Wang, Y., Goldin, A.L., and Catterall, W.A. (1992). A cluster of hydrophobic amino acid residues required for fast Na(+)-channel inactivation. *Proc. Natl. Acad. Sci. USA* *89*, 10910–10914.
- Xiong, D., Li, T., Dai, H., Arena, A.F., Plant, L.D., and Goldstein, S.A.N. (2017). SUMOylation determines the voltage required to activate cardiac I_{Ks} channels. *Proc. Natl. Acad. Sci. USA* *114*, E6686–E6694.
- Yang, T., Atack, T.C., Stroud, D.M., Zhang, W., Hall, L., and Roden, D.M. (2012). Blocking Scn10a channels in heart reduces late sodium current and is antiarrhythmic. *Circ. Res.* *111*, 322–332.
- Yang, X., Pabon, L., and Murry, C.E. (2014). Engineering adolescence: maturation of human pluripotent stem cell-derived cardiomyocytes. *Circ. Res.* *114*, 511–523.

STAR★METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Bacterial and Virus Strains		
BL21(DE3) E. Coli	Thermo Fisher	Cat#: EC0114
Chemicals, Peptides, and Recombinant Proteins		
Human: SUMO	Boston Biochem	Cat #: UL-712
Human: His6-SEN1 catalytic domain	Boston Biochem	Cat#: E-700
Ranolazine dihydrochloride	Sigma-Aldrich	Cat#: R6152
Experimental Models: Cell Lines		
Hamster: CHO-K1 cells	ATCC	Cat#: CCL-61; RRID: CVCL_0214
Human: iPS-ventricular cardiomyocytes-2	Cellular Dynamics	Cat# CMC-100-012-000.5
Recombinant DNA		
hNav _v 1.5-pcDNA1	Plant et al., 2006	DOI: 10.1172/JCI25618
GFP-SCN1B-pcDNA1	Plant et al., 2006	DOI: 10.1172/JCI25618
mCherry-SUMO1-pMAX	Plant et al., 2016	https://doi.org/10.7554/eLife.20054
mTFP1-hNav1.5-pMAX	This paper	N/A
eYFP-SUMO1-pMAX	Plant et al., 2010	https://doi.org/10.1073/pnas.1004712107
eYFP-Ubc9-pMAX	Plant et al., 2010	https://doi.org/10.1073/pnas.1004712107
Plasmid for MS with SUMO1(T95K) ₁₋₉₇	Plant et al., 2010	https://doi.org/10.1073/pnas.1004712107
Plasmid for MS with Human Nav _v 1.5 residues 353-502	Plant et al., 2010 , modified as described	https://doi.org/10.1073/pnas.1004712107
Software and Algorithms		
pClamp version 10	Molecular Devices	N/A
Origin version 6	Microcal	N/A
Prism version 8	GraphPad	N/A
ImageJ-FIJI	Schindelin et al., 2012 ; https://imagej.net/Fiji	https://doi.org/10.1038/nmeth.2019
MSCConvert	http://proteowizard.sourceforge.net/tools.shtml	
Batch-Tag in Protein Prospector v. 6.1.0	http://prospector.ucsf.edu/	
O'Hara Rudy Model (ORD)	O'Hara et al., 2011	https://doi.org/10.1371/journal.pcbi.1002061

LEAD CONTACT AND MATERIALS AVAILABILITY

Further Information and requests for resources and reagents should be directed to the Lead Contact, S.A.N. Goldstein (sgoldst2@uci.edu). Materials generated through this work are available from the Lead Contact upon reasonable request.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

CHO-K1 cells (RRID: CVCL_0214) were obtained from ATCC and used for heterologous expression, as detailed below. Human induced pluripotent stem cell cardiomyocytes² were purchased from Cellular Dynamics and cultured on #1.5 glass coverslips coated with gelatin (0.1% w/v in water) according to the manufacturer's instructions. Cells were maintained in iCell maintenance medium (Cellular Dynamics) and incubated at 37°C in a humidified atmosphere containing 5% CO₂ / 95% air (21% O₂). The iPS cells were studied between days 5 - 10 in culture at which time both isolated cells and cells in syncytia exhibited rhythmic beating behavior, a criterium for mature myocyte attributes shown to correspond with expression of troponin-T and sarcomeric α -actinin ([Ma et al., 2011](#)). To further verify that the currents under study in iPS cells were mediated by Nav_v1.5 channels, we employed protocols and

solutions to minimize conductance through potassium and calcium channels (per Figures 1, S1, and 5); showed that the currents had biophysical hallmarks like those of human $\text{Na}_v1.5$ in tissue culture cells (per Table 1 and Figures 3 and S3); and demonstrated blockade with ranolazine, a drug that inhibits I_{LATE} mediated by $\text{Na}_v1.5$ (per Table S3 and Figure 5).

METHOD DETAILS

Molecular Biology and Reagents

Human $\text{Na}_v1.5$ (NM_198056.1) was handled in pcDNA1, as previously described (Plant et al., 2006) and was transiently cotransfected with GFP-tagged SCN1B (the β subunit of cardiac SCN5A channel, isoform b; GenBank: NM_001037). Human SUMO1₁₀₁ (GenBank: NM_003352.8) and Ubc9 (GenBank: NM_003345.5) were amplified from a brain cDNA library (Clontech) and inserted into pMAX, as described before (Rajan et al., 2005). Sequences encoding mCherry or mTFP1 were inserted as described (Plant et al., 2010) at the N terminus of SUMO1 or $\text{Na}_v1.5$ respectively. SUMO1 and Ubc9 were tagged with eYFP using the same strategy. Mutations were introduced with Pfu Quikchange PCR (Agilent). Purified SUMO1₉₇, SUMO2₉₃, SENP1 and SENP2 were purchased from Boston Biochemical.

Heterologous Expression

CHO-K1 cells (RRID: CVCL_0214) were maintained in F12K medium supplemented with 10% FBS. Plasmids were transfected into cells with Lipofectamine 2000 according to the manufacturer's instructions (Life Technologies). Experiments were performed 24 to 48 hours post transfection at room temperature. The human $\beta 1$ accessory subunit was co-expressed with $\text{Na}_v1.5$ subunits unless otherwise indicated.

Electrophysiology

I_{Na} and action potentials in iPS-CMs, and whole-cell $\text{Na}_v1.5$ channel currents in heterologous cells were recorded using an Axopatch 200B amplifier and pCLAMP software (Molecular Devices) at filter and sampling frequencies of 10 and 50 kHz respectively. For voltage-clamp studies, cells were superfused with a solution comprising, in mM: NaCl 130, CsCl 4, CaCl_2 2, MgCl_2 1.2, glucose 5.5, HEPES 10 and, 200 μM CdCl_2 , a concentration previously shown to block >98% of voltage-gated calcium channel current in primary cells (Pearson et al., 1993). The pH was adjusted to 7.4 with NaOH and HCl. Cells were studied at room temperature with borosilicate glass pipettes (Clark Kent) with a resistance of 2–3 $\text{M}\Omega$ when filled with a solution comprising, in mM: CsCl 60, CsF 80, CaCl_2 1, MgCl_2 1, Na_2ATP 5, EGTA 10, HEPES 10, pH 7.4 with CsOH. Pipettes were coated with Sylgard (Dow Corning) prior to use. Capacitance artifacts were subtracted online, series resistance was compensated to 70% and cells with a series resistance of less than 10 $\text{M}\Omega$ were studied. Once whole-cell mode was established, cells were not studied for more than 600 s in order to maintain consistent membrane seals and voltage-clamp control of I_{Na} . Current-voltage relationships were evoked from a holding potential of -100 mV by 100 ms test pulses between -100 and 0 mV, in 10 mV increments. Steady-state inactivation was studied by holding cells at -140 mV and then comparing currents evoked by 50 ms test pulses between -140 mV and -20 mV to those evoked by a 100 ms prepulse to 0 mV. A 10 s interpulse interval was used in both cases. Normalized peak current values are plotted against prepulse potential (mV). A Boltzmann function, $I = I_{\text{max}} / (1 + \exp[(V - V_{1/2})/k])$, where I_{max} is the maximum current and k is slope factor, was used to fit normalized activation-voltage relationships. Recovery from fast inactivation was studied by holding cells at -100 mV and comparing currents evoked by a pair of 50 ms test step to -30 mV separated by an interpulse interval that increased in duration by 5 ms increments per sweep. The time constant for recovery from inactivation (τ) was obtained from mono-exponential fits of the normalized current amplitude to the recovery time using $I_t = I_{\text{max}} + A[e^{-t/\tau}]$, where A is the amplitude of components and t is time. Whole-cell currents were normalized to cell capacitance. Mean \pm SEM capacitance values were 28 ± 5 pF for iPS cells and 10 ± 2 pF for CHO cells.

For current-clamp studies, iPS-CMs were superfused with a solution comprising, in mM: NaCl 135, KCl 3.5, CaCl_2 2, MgCl_2 1.2, glucose 8, HEPES 10 and the pH was adjusted to 7.4. The intracellular solution contained, in mM: KCl 140, MgCl_2 2, Na_2ATP 5, EGTA 1, HEPES 5 and the pH was adjusted to 7.4.

Acute Hypoxia for Electrophysiology

Acute hypoxia was achieved as described previously (Plant et al., 2002). Briefly, the cells were made hypoxic (1.5% O_2) by switching the perfusate with one that had been bubbled with nitrogen for at least 30 minutes prior to perfusion. Oxygen tension was measured at the cell by a polarized carbon fiber electrode; solution exchange occurred in less than 10 s. Ambient O_2 , or normoxia recording conditions, were achieved by allowing recording solutions to equilibrate with room air for at least 30 minutes prior to perfusion. The ambient O_2 solution was measured to be 21% O_2 at the cell. All solutions were equilibrated, measured, and perfused at room temperature (18°C).

Two-Color TIRFM

Single protein complexes at the surface of live CHO-K1 cells were identified using TIRFM as previously described (Plant et al., 2014). Cells were studied in a solution comprising (in mM): NaCl 130, KCl 4, MgCl_2 1.2, CaCl_2 2, HEPES 10, pH was adjusted to 7.4 with NaOH. The critical angle for TIRF was adjusted using a CellTIRF illuminator (Olympus) and a high numerical aperture apochromat objective (150x, 1.45 NA) mounted on an automated fluorescence microscope (IX81) controlled by Metamorph software (Molecular Devices). For simultaneous illumination of two fluorophores, CellTIRF software (Olympus) was used to adjust the critical angle for each excitation wavelength to generate evanescent waves of equivalent depth (100 nm). mCherry was excited by a 561-nm laser line and mTFP1 was excited a 445-nm laser line. When mTFP1 was studied with mCherry the emitted light signals were split using

a 520-nm dichroic mirror mounted in a DualView adaptor (Photometrics) and each wavelength was directed to one half of an EM-CCD. The dichroic mirror was disengaged when single fluorophores were studied.

To assess stoichiometry, fluorophores were photobleached by continual excitation and data were captured as movies of 100–400 frames acquired at 1 Hz using an EM-CCD camera (Hamamatsu). When mTFP1 was studied with mCherry in the same cell, the data for each fluorophore were saved as separate stacks and processed in an identical manner. Images were background corrected by subtracting the mean of 5 fully bleached frames from the end of each stack analyzed. Misalignment of the data between stacks was corrected in ImageJ using StackReg. Fluorescent spots were defined as a discrete 3 × 3-pixel region around a pixel of maximum intensity, as before (Plant et al., 2010, 2014). Fluorescence is reported as the change in fluorescence intensity normalized by the initial fluorescence for each trace. The density of co-localized and single fluor-fluorescent particles was determined following thresholding and watershed separation in ImageJ. The particle number was counted in 4, separate 100 × 100-pixel regions of interest for 6–10 cells per group using the Analyze particles plugin.

Manders' coefficient of colocalization was assessed from live-cell simultaneous two-color TIRF images captured at 5 s intervals to minimize photobleaching. Data stacks were background subtracted and aligned prepared for each fluorophore post hoc, as above. Co-localization of partner pixels from the two stacks of images was defined as the presence of both fluorophores with at least 30% of maximum fluorescence levels recorded in that region of interest. Mean Manders' coefficients were calculated for 3–5 separate 100 × 100-pixel regions of interest per cell.

Live Cell FRET Microscopy

Donor-decay time-course was studied as before (Plant et al., 2010), using an Olympus inverted epi-fluorescence microscope. Cells were studied in a solution comprising (in mM): NaCl 130, KCl 4, MgCl₂ 1.2, CaCl₂ 2, HEPES 10, pH was adjusted to 7.4 with NaOH. mTFP1 was excited at 445 nm and the emission collected through a 470–500 nm bandpass filter, YFP was excited at 514 nm and the emission collected through a 525–575 nm filter. Images were captured using a scientific camera controlled by μ manager or Metamorph software (Molecular Devices) and were analyzed with ImageJ.

Mass Spectrometry

Human Na_v1.5 residues 353–502 were cloned into pET28a vector with six-His residues and a tobacco etch virus (TEV) cleavage site replacing the thrombin site and expressed in BL21(DE3) *E. Coli* with a vector carrying mouse E1 (as a linear fusion product of Aox1 and Uba2), E2 (Ubc-9) and SUMO1_{1–97} with a T95K mutation so that trypsin digestion before MS leaves a Gly-Gly tag on the ϵ -amino group of target lysines as before (Plant et al., 2010). Protein was purified with Ni-NTA affinity columns. The eluate was reduced with TCEP and alkylated using iodoacetamide prior to tryptic digestion via filter-assisted digestion. LC MS/MS analysis was performed using a Thermo Scientific Dionex UltiMate 3000 system coupled in-line to an Orbitrap Fusion Lumos MS. A 25 cm × 75 μ m PepMap EASY-Spray Acclaim PepMap C18 Column was used to separate peptides over acetonitrile gradients of 4% to 28% at a flow rate of 300 nL/min. Each duty cycle consists of one MS scan in FT mode (350–1500 m/z, resolution of 120,000 at m/z 400) followed by data-dependent tandem mass scans in IT mode for 3 s at top speed, utilizing HCD with NCE 30% on the most intense ions with charge states +2 or higher. The RAW data file acquired was extracted to MGF format using MSConvert (<http://proteowizard.sourceforge.net/tools.shtml>) and subjected to database searching using Batch-Tag within a developmental version of Protein Prospector (v. 6.1.0) against a decoy database consisting of a normal *Homo sapiens* Swissprot database concatenated with a randomized version (SwissProt.2019.4.8.random.concat, total 20,418 protein entries). The mass accuracy for parent ions and fragment ions were set as \pm 10 ppm and 0.2 Da, respectively. Trypsin was set as the enzyme and maximum of two missed cleavages were allowed. Protein N-terminal acetylation, methionine oxidation, Gly-Gly, and N-terminal conversion of glutamine to pyroglutamic acid were selected as variable modifications. Minimum protein and peptide scores were set as 22.0 and 5.0 respectively, while maximum E values for proteins and peptides were set to 0.01 and 0.1. A list of identified precursor ions corresponding to peptides carrying GlyGly modifications were generated for targeted sequencing for further validation. All MS/MS spectra for potential SUMOylated peptides were further inspected manually and evaluated based on spectral quality.

Simulations

Action potentials were simulated using the O'Hara Rudy (ORD) model (O'Hara et al., 2011) under normal condition or by scaling the conductance of I_{Late} . To simulate the effect of SUMO (or hypoxia-induced SUMOylation) a scaling factor named "SUMO" that varies from 1 (No SUMO) to 10 (maximum effect at 1 nM SUMO1) was added to the model to increase the conductance of I_{Late} : $G_{NaLate}^{SUMO} = G_{NaLate} * SUMO$, where G_{NaLate} is the nominal conductance value from the O'Hara-Rudy ventricular myocyte model and G_{NaLate}^{SUMO} is the conductance used for simulation in the modified ORD. Other variables and all equations of the model were otherwise as described in the published model, except that the external ionic concentrations were set to match the experimental conditions used here to study the iPS-CMs (in mM): $[Na^+]_o = 135$; $[Ca^{2+}]_o = 2.0$; and $[K^+]_o = 3.5$. A pacing cycle length of 800 ms was used and the numbers of beats was set to 1000. The duration of the APD₅₀ was measured from the time of maximum dV/dt to the time that membrane voltage reached 50% of complete repolarization; resting potential was measured immediately before each beat.

QUANTIFICATION AND STATISTICAL ANALYSIS

Data were analyzed using pClamp, Origin, GraphPad and Excel software. Quantification and analysis specific in each experimental technique is described in the [Method Details](#) section, above. Data were assessed for statistical differences between groups by

one-way analysis of variance with Bonferroni post hoc analysis to test differences within pairs of group means for all dataset with an F-value of $p < 0.05$. Data are presented, where indicated as the mean \pm standard error of the mean (SEM). The number of replicates for each study are described in the legends.

DATA AND CODE AVAILABILITY

The study did not generate any unique datasets or codes.

Cell Reports, Volume 30

Supplemental Information

**Hypoxia Produces Pro-arrhythmic
Late Sodium Current in Cardiac Myocytes
by SUMOylation of Na_v1.5 Channels**

Leigh D. Plant, Dazhi Xiong, Jesus Romero, Hui Dai, and Steve A.N. Goldstein

SUPPLEMENTAL TABLES

Table S1. Biophysical parameters for mTFP-Nav_v1.5 variants. Related to Figure 2.

Table S2. Hypoxia-induced changes in I_{LATE} correlate with SUMOylation. Related to Figure 4.

Table S3. Hypoxia yields a SUMO-dependent increase in APD in iPS-CMs. Related to Figure 5.

SUPPLEMENTAL FIGURES

Fig. S1. Hypoxia, or SUMO1 in ambient oxygen, augment Nav_v1.5 window-current in human iPS-CMs. Related to Figure 1.

Fig. S2. FRET shows that SUMOylation of Nav_v1.5 K442 and no other site. Related to Figure 2.

Fig. S3. Acute hypoxia and SUMO1 augment the Nav_v1.5 channel window current in CHO cells. Related to Figure 3.

Fig. S4. Mass spectroscopic demonstration of SUMOylation of Nav_v1.5 K442. Related to Figures 2 and 3.

Fig. S5. Reopening of single Nav_v1.5 channels requires SUMO1. Related to Figure 3.

Fig. S6. Hypoxia does not recruit SUMO1 to Nav_v1.5-K442Q channels at the cell surface. Related to Figure 4.

Table S1. Biophysical parameters for mTFP-Nav_v1.5 variants. Related to Figure 2.

Channel variant	I_{PEAK} , 21% O ₂ pA/pF	I_{LATE} , 21% O ₂ pA/pF	I_{PEAK} , 1.5% O ₂ pA/pF	I_{LATE} , 1.5% O ₂ pA/pF
mTFP-Nav _v 1.5	-198 ± 6	-0.96 ± 0.2	-201 ± 4	-8.3 ± 0.6
mTFP-Nav _v 1.5-K442Q	-200 ± 4	-0.93 ± 0.3	-197 ± 6	-0.91 ± 0.4*
mTFP-Nav _v 1.5-K496Q	-198 ± 3	-0.94 ± 0.2	-197 ± 3	-8.3 ± 0.3
mTFP-Nav _v 1.5-K1120Q	-193 ± 4	-0.95 ± 0.2	-196 ± 3	-8.7 ± 0.3
mTFP-Nav _v 1.5-K1162Q	-199 ± 4	-0.91 ± 0.2	-194 ± 5	-8.7 ± 0.4
mTFP-Nav _v 1.5-K1800Q	-195 ± 3	-0.97 ± 0.3	-193 ± 4	-8.8 ± 0.5

Legend. Human Nav_v1.5 tagged with mTFP1 was expressed with Nav_vβ1 and studied in live CHO-K1 cells. As observed with untagged channels, mTFP-Nav_v1.5-K442Q channels are insensitive to acute hypoxia. The effect of 1.5% O₂ on the magnitude of the peak (I_{PEAK}) and late (I_{LATE}) sodium current was assessed at -30 mV, as described in Fig. 3. Data are mean ± s.e.m. for 8 – 12 cells per group. Statistical significance between the condition studied and the normoxic condition for each cell was determined using a unpaired t-test, where P<0.05 is represented by an asterisk (*).

Table S2. Hypoxia-induced changes in I_{LATE} correlate with SUMOylation. Related to Figure 4.

Time after start of hypoxia, s	I_{LATE} (% I_{PEAK})	Manders' coefficient	Pearson's correlation coefficient
0	0.05 ± 0.07	0.10 ± 0.01	0.075 ± 0.01
50	2.8 ± 0.30	0.66 ± 0.02	0.62 ± 0.02
100	4.1 ± 0.44	0.91 ± 0.03	0.89 ± 0.015
300	4.0 ± 0.50	0.88 ± 0.03	0.86 ± 0.02
500	4.3 ± 0.49	0.89 ± 0.02	0.87 ± 0.03


Legend. The late current was determined as the %-change in peak current as per **Fig 5** and **Table 2** for Nav1.5 channels expressed in CHO cells and studied in whole-cell mode. Briefly, cells were subject to hypoxia (1.5% O₂ at time = 0 s), and returned to 21% oxygen at 300 s; cells were studied for up to a total of 500 s. Manders' coefficient and Pearson's correlation coefficient were determined for the hypoxia-induced co-localization of mTFP-tagged Nav1.5 channels and mCherry-tagged SUMO1 in TIRFM studies, as described in **Fig 4** using the Coloc2 algorithm in FIJI (ImageJ). Data are mean \pm S.E.M. from 7-10 regions studied in 8 cells.

Table S3. Hypoxia yields a SUMO-dependent increase in APD in iPS-CMs. Related to Figure 5.

Parameter	Control			SUMO1			SENP1		
	21% O ₂	1.5% O ₂	1.5% O ₂ + ranolazine	21% O ₂	1.5% O ₂	1.5% O ₂ + ranolazine	21% O ₂	1.5% O ₂	1.5% O ₂ + ranolazine
Resting potential, mV	-60.2 ± 3	-59.5 ± 3	-59.7 ± 3	-58.8 ± 2	-58.2 ± 3	-58.5 ± 3	-60.0 ± 1	-61.5 ± 4	-61 ± 2
dV/dt , V/s	15.4 ± 2	16.1 ± 3	15.8 ± 3	16.1 ± 2	15.8 ± 3	16.5 ± 1	16.6 ± 2	15.9 ± 4	15.5 ± 3
AP peak, mV	49.8 ± 3	50.5 ± 4	50.3 ± 3	51.2 ± 3	51.8 ± 4	51.5 ± 2	50 ± 3	48.5 ± 1	49 ± 2
APD ₅₀ , ms	663 ± 28	$810 \pm 19^*$	690 ± 17	822 ± 22	869 ± 27	$699 \pm 20^*$	663 ± 26	699 ± 24	663 ± 19
APD ₉₀ , ms	918 ± 32	$1122 \pm 26^*$	945 ± 28	1127 ± 24	1184 ± 27	$905.4 \pm 30^*$	909 ± 32	947 ± 21	905.4 ± 29

Legend. Human cardiac myocytes derived from iPS cells (**Fig. 1 & 5**) were studied by current-clamp in whole-cell mode. Spontaneous action potentials were studied with control intracellular solution of with 1 nM SUMO1 or SENP1 in the recording pipette with the same sequence of perfusates: normoxic (21% O₂), hypoxic (1.5% O₂), then hypoxia with 1 μ M ranolazine. dV/dt is the velocity of the upstroke of the action potential. The action potential duration (APD₅₀ and APD₉₀) was determined by calculating the time required for the membrane potential to return to 50% and 90% of the resting value from the peak deflection of the action potential. Data are means \pm S.E.M. for 6-9 cells studied per group; statistical significance between the condition studied and the normoxic condition for each cell was determined using a two-way paired t-test, * indicates P<0.01.

Figure S1. Hypoxia, or SUMO1 in ambient oxygen, augment I_{Na} window-current in human iPS-CMs. Related to Figure 1.




Legend. I_{LATE} is observed in the ‘window’ between the voltages where Na_v channels activate and inactivate. The window-current in iPS-CMs was studied by measuring the voltage-dependence of I_{Na} activation (circles) and steady-state inactivation (SSI, squares). Activation was assessed using current-voltage relationships evoked from a holding potential of -100 mV by 100 ms test pulses between -100 and 0 mV, in 10 mV increments. Steady-state inactivation was studied by holding cells at -140 mV and then comparing currents evoked by 50 ms test pulses between -140 mV and -20 mV to those evoked by a 100 ms prepulse to 0 ms. A 10 s interpulse interval was used in both cases. Normalized peak currents were plotted against prepulse potential (mV). Normalized activation and inactivation relationships were fit with a Boltzmann function, $I = I_{max}/(1 + \exp[(V - V_{1/2})/k])$, where I_{max} is the maximum current and k is slope factor. Data are mean \pm S.E.M. for 8 to 10 cells per group.

- A.** *Upper*, when cells were studied with a control pipette solution, exchanging the normoxic perfusate (**black line**) with a hypoxic perfusate (**black dash**) left shifted the $V_{1/2}$ of activation and right shifted $V_{1/2}$ -SSI. *Lower*, the changes in activation and SSI increased the magnitude of the window current.
- B.** *Upper*, when cells were studied with 1 nM SUMO1 in the pipette (**red line**) the $V_{1/2}$ -activation was left shifted compared to control (**A**) and $V_{1/2}$ -SSI was right shifted. The currents were insensitive to hypoxia (**red dash**). *Lower*, SUMO1 increased the magnitude of the window current compared to control (**A**).
- C.** *Upper*, 1 nM SENP1 in the pipette (**blue line**) did not alter the $V_{1/2}$ of activation or SSI compared to control (**A**) and currents were insensitive to hypoxia (**blue dash**). *Lower*, SENP1 did not change the window current.
- D.** The recovery of I_{Na} from fast inactivation is not altered by hypoxia, SUMO1 or SENP1. The rate at which I_{Na} recovered from fast inactivation was determined using the protocol detailed in the methods. The time-constant for recovery τ , was obtained by fitting the normalized current amplitude to the recovery time using the mono-exponential function: $I_t = I_{max} + A[e^{-t/\tau}]$. The mean $\tau \pm$ s.e.m. was not different for 6-8 cells

studied under each of the experimental conditions indicated. For iPS-CMs studied with control pipette solution, τ was 13.6 ± 0.6 ms under ambient conditions and 12.9 ± 0.3 ms when the cells were subjected to hypoxia. The exemplar record, inset, shows the recovery of $\text{Na}_v1.5$ channels studied in ambient levels of O_2 . When cells were studied with 1 nM SUMO1 in the pipette, τ was 14.8 ± 1 ms in normoxia and 13.7 ± 0.8 ms in hypoxia. When 1 nM SENP1 was included in the pipette, τ changed from 13.2 ± 0.3 ms in normoxia and 12.4 ± 0.5 ms in hypoxia. Similar results were obtained for $\text{Na}_v1.5$ and $\text{Na}_v1.5\text{-K442Q}$ channels expressed in CHO-K1 cells with $\text{Na}_v\beta1$.

Figure S2. FRET shows SUMOylation of $\text{Na}_v1.5$ K442 and no other site. Related to Figure 2.



Legend. Human $\text{Na}_v1.5$ tagged with mTFP1 was expressed with $\text{Na}_v\beta1$ and studied in live CHO-K1 cells. FRET between mTFP1 (donor) and YFP-tagged subunits (acceptor) was assessed by donor photobleaching, as described in Fig. 2. Data are mean $\tau \pm$ s.e.m. for 3 regions of 5-7 cells per group. The effect of 1.5% O_2 on the magnitude of the late sodium current was assessed at -30 mV, as described in Fig. 3.


A. A cartoon showing the secondary structure of mTFP- $\text{Na}_v1.5$ and the location of putative SUMOylation sites identified by the SUMOplot algorithm. K442, the only site where SUMO1 was found to interact with $\text{Na}_v1.5$ (Figs. 2-4) is in red.

B. *Left*, A histogram of mean FRET τ -values show assembly of mTFP- $\text{Na}_v1.5\text{-K496Q}$ channels with YFP-SUMO1 and YFP-Ubc9 but not linkage-incompetent YFP-SUMO1₉₅ or free YFP. *Right*, Exemplar currents show the development of late current when cells expressing mTFP- $\text{Na}_v1.5\text{-K496Q}$ were studied in ambient (black) and hypoxic conditions (red).

C. **E.** FRET studies show that mTFP- $\text{Na}_v1.5\text{-K1120Q}$, mTFP- $\text{Na}_v1.5\text{-K1162Q}$ and mTFP- $\text{Na}_v1.5\text{-K1800Q}$ channels interact with YFP-SUMO1 and Ubc9 like wild type and whole-cell current recording shows that hypoxia induced I_{LATE} in each of the channels.

F. A histogram showing the mean I_{LATE} as % of peak current from cells studied at 21% O_2 (open), after 200 s at 1.5% O_2 (hatched) and after 200 s following a return to 21% O_2 with the pipette solutions indicated (grey).


Figure S3. Acute hypoxia and SUMO1 augment the $\text{Na}_v1.5$ channel window current in CHO cells. Related to Figure 3.



Legend. $\text{Na}_v1.5$ or $\text{Na}_v1.5\text{-K442Q}$ were expressed in CHO-K1 cells with $\text{Na}_v\beta1$ and studied in whole-cell mode to determine normalized activation (circles) and steady-state inactivation (SSI; squares) relationships for peak currents, as described in the STAR Methods and for iPSC-CMs in **Fig. S1**. Measured values are noted in the text and **Table 1**. Hypoxia was 1.5% O_2 . Cells were studied with control solution (**black**), 1 nM SUMO1 (**red**), or 1 nM SENP1 (**blue**) in the recording pipette. Data are mean \pm S.E.M. for 8 to 10 cells per group.

- A.** *Upper*, when $\text{Na}_v1.5$ channels were studied with a control pipette solution, exchanging the normoxic perfusate (**black line**) with a hypoxic perfusate (**black dash**) left shifted the $V_{1/2}$ of activation and right shifted $V_{1/2}\text{-SSI}$. *Lower*, the changes in activation and SSI increased the magnitude of the window current.
- B.** *Upper*, studying $\text{Na}_v1.5$ channels with 1 nM SUMO1 in the pipette (**red line**) left-shifted the $V_{1/2}$ -activation compared to control (**A**) and right shifted $V_{1/2}\text{-SSI}$. The currents were insensitive to hypoxia (**red dash**). *Lower*, SUMO1 increased the magnitude of the window current compared to control (**A**).
- C.** *Upper*, 1 nM SENP1 in the pipette (**blue line**) did not alter the $V_{1/2}$ of activation or SSI compared to control (**A**) and the currents were insensitive to hypoxia (**blue dash**). *Lower*, SENP1 did not change the window current.
- D. F.** The $V_{1/2}\text{-act}$ and the $V_{1/2}\text{-SSI}$ of cells expressing $\text{Na}_v1.5\text{-K44Q}$ was not altered by hypoxia, or when the cells were studied with 1 nM SUMO1 or SENP1 in the recording pipette.


Figure S4. Mass spectroscopic demonstration of SUMOylation of Na_v1.5 K442. Related to Figures 2 and 3.



Legend. For MS analysis, Na_v1.5₃₅₃₋₅₀₂ and SUMO1₉₇T95K were expressed with SUMOylation enzymes in *E. coli*, purified, subjected to trypsin cleavage, and analyzed by MS as in the STAR Methods.

- A.** *Top*, schematic showing the product Na_v1.5₃₅₃₋₅₀₂-SUMO (box) and the trypsin fragment of Na_v1.5 carrying the Gly-Gly remnant of SUMO1₉₇T95K. *Bottom left*, the sequence of the three-ended fragment with K442 and the Gly-Gly remnant. *Bottom right*, a Coomassie blue-stained SDS-PAGE gel of the purified products. Unmodified Na_v1.5₃₅₃₋₅₀₂ migrates at ~20 kDa (arrow). Expression of SUMO1₉₇T95K (SUMO) and the SUMO enzymes yields SUMO on the overexpressed target as well as native protein and the Na_v1.5₃₅₃₋₅₀₂-SUMO conjugate migrates at ~35 kDa (star). Molecular weight markers are shown.
- B.** Tandem MS sequence analysis of the fragment with Lys442 and the Gly-Gly remnant indicating *b* and *y* ion species as annotated in (a).

Figure S5. Reopening of single Nav1.5 channels requires SUMO1. Related to Figure 3.




Legend. Human Nav_v1.5 and Nav_vβ1 were studied in inside-out patches excised from CHO-K1 cells, as described in the STAR Methods. Single-channel currents were stimulated every 5 s by a 50-ms depolarizing pulse to -30 mV from a holding potential of -120 mV. Data were recorded at filter and sampling frequencies of 5 and 50 kHz, respectively but were refiltered offline using a 1 kHz Bessel filter for display purposes. For each cell, null sweeps (with no channel activity) were identified, averaged offline, and subtracted from data sweeps before analysis. Recording pipettes were filled with extracellular buffer and the inside of patches was perfused with the

intracellular buffer, with or without 1 nM SUMO1 or 1 nM SENP1, as indicated. Hypoxia, 1.5% O₂.

- Upon depolarization, single Nav_v1.5 channels opened, inactivated rapidly, and did not reopen when studied in normoxia or when exposed to hypoxia. The addition of 1 nM SUMO1 induced channel reopening in patches studied in 1.5% O₂ (upper), or 21% O₂ (lower). Null traces were 50 ± 4% (n = 10 patches; 1,000 sweeps) for both configurations of the experiment.
- Upon depolarization, single Nav_v1.5-K442Q channels opened, inactivated rapidly, and did not reopen when exposed to hypoxia or 1 nM SUMO1. Null traces were 52 ± 6% (n = 10 patches; 1,000 sweeps) for both configurations of the experiment.
- A scatter plot showing the portion of patches with single Nav_v1.5 channels that showed reopening plotted against time after the transition to hypoxia. Nav_v1.5 channels were studied in 21% O₂ (●), 21% O₂ + SUMO1 (●), 1.5% O₂ (○), 1.5% O₂ + SUMO1 (○). Reopening was observed with SUMO1 in both 21% and 1.5% O₂, reached at maximum at 155 ± 7 s in each case, with 50% of the patches showing channel reopening at 65 ± 10 s and 67 ± 8 s respectively. Nav_v1.5-K442Q channels were studied with 1.5% O₂ + SUMO1 (◇) and did not reopen.
- Ensemble average traces (n = 200-220 sweeps per channel) showing that hypoxia-induced reopening of Nav_v1.5 channels leads to I_{LATE} only in the presence of SUMO1.
- Left*, A histogram showing the mean I_{LATE} as a % of the peak current from ensemble average currents obtained from off-cell patches studied at 21% O₂, and after 200 s at 1.5% O₂ with the intracellular solutions indicated. *Right*, A histogram showing the mean I_{LATE} as a % of the peak current from ensemble average currents obtained from off-cell patches studied at 21% O₂, and after 200 s at 1.5% O₂ with the intracellular solutions indicated.

Figure S6. Hypoxia does not recruit SUMO1 to Nav1.5-K442Q channels at the cell surface.

Related to Figure 4.



Legend. Single mTFP-Nav_v1.5 and mTFP-Nav_v1.5-K442Q channels and mCherry-SUMO1 (m-SUMO1) were studied in CHO-K1 cells by TIRFM. Stoichiometric (photobleaching) and pixel-by-pixel analysis for subunit density and colocalization were studied as per **Fig. 4**. The surface-density of fluorescent pixels was assessed post-hoc for 3–5 regions per cell. Stoichiometric analysis represents 5–8 cells and is summarized in **Table 2**.

- A.** Histogram of photobleaching steps showing that m-SUMO1₉₅, lacking the C-terminal diglycine motif required for conjugation is not observed at the cells surface, despite the expression of mTFP-Nav_v1.5 channels (teal, single bleaching step per particle). Hypoxia does not increase the surface presence of m-SUMO1₉₅ and does not alter the channel subunit stoichiometry (**Table 2**).
- B.** Histogram of photobleaching steps showing that mTFP-Nav_v1.5-K442Q channels contain one subunit in normoxic and hypoxia conditions and do not associate with m-SUMO1.
- C.** *Left*, in ambient O₂ (0 seconds of hypoxia), the surface density of m-SUMO1 (top) is low compared to mTFP-Nav_v1.5-K442Q channels (middle) with little co-localization (bottom). *Right*, hypoxia does not recruit m-SUMO1 to the cell surface and m-SUMO1 is not observed to colocalize with mTFP-Nav_v1.5-K442Q channels (bottom); surface levels of mTFP-Nav_v1.5-K442Q were not observed to change (middle).
- D.** Histogram of surface density summarizing six cells studied as described in (A) and **Table 2**. At ambient O₂ (0 seconds of hypoxia), the density of pixels per μm² with SUMO1 alone (red) was 2.8 ± 1.0, and 328 ± 26 for Nav_v1.5-K442Q (blue). The density of pixels with both fluors was 3.6 ± 1 per μm² (green). This distribution of subunits was not altered by 100 s hypoxia. Cells were observed to have 3.3 ± 1 colocalized pixels per μm², 326 ± 25 pixels with free Nav_v1.5 channels per μm² and 4 ± 2 pixels per μm² with free SUMO1.