

Supplementary Table 1. Serum triglycerides levels by current smoking status and smoked pack-years (stratified for alcohol drinking status)

Smoking status	Log TG								
	Non-drinkers (n = 97)		Non-high risk drinkers (n = 1,175)		High risk drinkers (n = 660)				
	β	95% CI	p value	β	95% CI	p value	β	95% CI	p value
Never-smoker (n = 29)	Reference			Reference			Reference		
Past light smoker (n = 38) ^a	0.07	-0.14 to 0.28	0.516	0.04	-0.03 to 0.12	0.234	0.03	-0.11 to 0.17	0.651
Past heavy smoker (n = 20) ^a	0.30	0.04 to 0.56	0.026	0.05	-0.05 to 0.14	0.329	0.13	-0.03 to 0.29	0.123
Current light smoker (n = 6) ^a	0.32	-0.08 to 0.71	0.113	0.14	0.05 to 0.22	0.002	0.14	0.001 to 0.28	0.048
Current heavy smoker (n = 4) ^a	-0.01	-0.44 to 0.43	0.977	0.11	0.02 to 0.21	0.017	0.27	0.13 to 0.41	< 0.001

Values are the results from multiple linear regression models adjusted for age, body mass index, systolic blood pressure, fasting serum glucose, physical activity and energy, carbohydrate and fat intake.

TG, triglyceride; CI, confidence interval.

^aPast/current light smokers are past/current smokers with less than 20 pack-year, past/current heavy smokers are past/current smokers with more than 20 pack-year.