

SUPPLEMENTARY TABLE S2. PARTICIPANT CHARACTERISTICS: 2010–2012 VERSUS 2016–2018

	2010–2012, N=25,529	2016–2018, <sup>a</sup> N=22,697
Age, years, mean ± SD	22 ± 17	26 ± 18
Female, <i>n</i> (%)	12,706 (50)	11,394 (50)
Non-Hispanic white, <i>n</i> (%)	20,954 (82)	18,368 (82)
Private insurance, <i>n</i> (%)	17,081 (75)	16,028 (74)
Duration of diabetes, median (IQR)	6 (2–14)	10 (6–19)
BMI z-score, mean ± SD	0.4 ± 1.0	0.4 ± 1.0
Pump use, <i>n</i> (%)	12,637 (57)	14,047 (63)
CGM use, <i>n</i> (%)	1591 (7)	6547 (30)
Self-monitoring of blood glucose, <sup>b</sup> mean ± SD	4.7 ± 3.0	4.1 ± 2.4
Noninsulin medications for blood glucose control, <i>n</i> (%)		
Metformin	388 (2)	811 (4)
GLP-1 agonist	34 (<1)	300 (1)
DPP-IVi	9 (<1)	9 (<1)
SGLT2i	0	232 (1)
Pramlintide	124 (<1)	131 (<1)
Other <sup>c</sup>	24 (<1)	28 (<1)

<sup>a</sup>Twelve thousand seven hundred five (56%) participants in the 2016–2018 cohort, also in the 2010–2012 cohort.

<sup>b</sup>Calculation excludes participants using CGM.

<sup>c</sup>Includes thiazolidinediones and sulfonylureas.