

SUPPLEMENTAL MATERIAL

Table S1. Baseline Characteristics.

	HPR patients	Clopidogrel	Ticagrelor/Prasugrel	p value
	n=196	n=114	n=82	
Age (years)	70.5±10.1	69.9±11.2	71.4±10.0	0.322
≥75 years	77 (39)	44 (39)	33 (40)	0.466
Male sex, (%)	160 (82)	91(80)	69 (84)	0.281
Hypertension, (%)	131 (67)	68 (60)	63 (77)	0.008
Hypercholesterolemia, (%)	127 (65)	73 (64)	54 (66)	0.457
Diabetes, (%)	66 (34)	38 (33)	28 (34)	0.513
CKD, (%)	24 (12)	16 (14)	8 (10)	0.250
Previous MI, (%)	103 (53)	64 (56)	39 (48)	0.149
Previous PCI, (%)	79 (40)	42 (37)	37 (45)	0.154
Previous CABG, (%)	35 (18)	21 (18)	14 (17)	0.481
ACS, (%)	49 (25)	29 (25)	20 (24)	0.867
LVEF < 0.40, (%)	67 (34)	45 (39)	22 (27)	0.045
Multivessel disease, (%)	174 (89)	99 (87)	75 (91)	0.219
Three-vessel disease, (%)	106 (54)	58 (51)	48 (58)	0.180
LAD CTO (%)	59 (30)	29 (25)	30 (37)	0.065

Values are mean ± SD and number of patients (%).

ACS = acute coronary syndrome; CABG = coronary artery bypass grafting; CKD = chronic kidney disease; CTO = chronic total occlusion; DES = drug-eluting stent; HPR= high platelet reactivity; LAD = left anterior descending artery; LVEF = left ventricular ejection fraction; MI = myocardial infarction; PCI = percutaneous coronary intervention; SD = standard deviation.