

Supplementary Information

^1H NMR Metabolic profile of scyphomedusa *Rhizostoma pulmo* (Scyphozoa, Cnidaria) in female gonads and somatic tissues: preliminary results

Angilè, F.¹, Del Coco L.¹, Girelli, C.R.¹, Basso, L.^{1,2}, Rizzo, L.^{2,3}, Piraino, S.^{1,2}, Stabili, L.^{1,4}, Fanizzi, F.P.^{1*}

¹ Department of Biological and Environmental Sciences and Technologies (Di.S.Te.B.A.), University of Salento, via Monteroni, 73100- Lecce (Italy); federica.angile@unisalento.it; laura.delcoco@unisalento.it; chiara.girelli@unisalento.it; fp.fanizzi@unisalento.it

² CoNISMa, Piazzale Flaminio, 9, Roma (Italy); lorena.basso@unisalento.it; stefano.piraino@unisalento.it

³ Integrative Marine Ecology, Stazione Zoologica Anton Dohrn, Villa Comunale, 80121 Napoli, (Italy); lucia.rizzo@szn.it

⁴ Water Research Institute of the National Research Council, (IRSA-CNR), Via Roma 3, Taranto, (Italy); loredana.stabili@irsa.cnr.it

* Correspondence: fp.fanizzi@unisalento.it; tel.: +39-0832-299265




Figure S1. a) t[1]/t[2] PCA scores plot for lipid extracts ($R^2X=0.95$, $Q^2=0.92$) of three body compartments of *R. pulmo*. b) t[1]/t[2] PCA scores plot for aqueous extracts ($R^2X=0.73$, $Q^2=0.53$) of three body compartments of *R. pulmo*. Red circle, female gonads; blue square, oral arms; green triangle, umbrella.