

Supplementary Materials

Physiological and Biochemical Mechanisms Mediated by Allelochemical Isoliquiritigenin on the Growth of Lettuce Seedlings

Shuang Zhang[†], Shi-Wei Sun[†], Hai-Lin Shi, Ke Zhao, Jin Wang, Yang Liu, Xiao-Hong Liu and Wei Wang*

Department of Natural Medicine and Pharmacognosy, School of Pharmacy, Qingdao University, Qingdao 266071, China; qdeduzhangshuang@163.com (S.Z.); sunsw@qdu.edu.cn (S.-W.S.); shihailinjin@163.com (H.-L.S.); Qingdao_zhaoke@163.com (K.Z.); Qingdao_wangjin@163.com (J.W.); buckuper@163.com (Y.L.); liuxiaohong1043@163.com (X.-H.L.)

* Correspondence: w.w.wangwei@263.net; Tel./Fax: +86-532-8699-1172 (W.W.)

[†] These authors contribute equally to this work and joint first authors.

Supporting information

Figure S-1. ¹H NMR spectrum of isoliquiritigenin.

Figure S-2. ¹³C NMR spectrum of isoliquiritigenin.




Figure S-1. ^1H NMR spectrum of isoliquiritigenin.




Figure S-2. ^{13}C NMR spectrum of isoliquiritigenin.