

S3 Table: Distribution of individuals exhibiting GIs in the Terry Collection by affected cranial bones (considering the left and right greater wings of the sphenoid bone as two separate bones), extent, and number of lesions (L = left, R = right). Number of individuals in the A) TB group and B) NTB group.

A)		Frontal bone	Parietal bone (L)	Parietal bone (R)	Temporal bone (L)	Temporal bone (R)	Sphenoid bone (L)	Sphenoid bone (R)	Occipital bone
TB group (Σ=234)		32/68 (47.06%)	6/68 (8.82%)	10/68 (14.71%)	20/68 (29.41%)	20/68 (29.41%)	10/68 (14.71%)	19/68 (27.94%)	62/68 (91.18%)
Extent (x) of lesions	x < 25%	31/32 (96.88%)	6/6 (100.00%)	10/10 (100.00%)	18/20 (90.00%)	14/20 (70.00%)	9/10 (90.00%)	11/19 (57.89%)	50/62 (80.65%)
	25% ≤ x < 50%	1/32 (3.13%)	–	–	2/20 (10.00%)	2/20 (10.00%)	1/10 (10.00%)	7/19 (36.84%)	11/62 (17.74%)
	50% ≤ x < 75%	–	–	–	–	4/20 (20.00%)	–	1/19 (5.26%)	1/62 (1.61%)
	75% ≤ x	–	–	–	–	–	–	–	–
Number of lesions	Unifocal	8/32 (25.00%)	4/6 (66.67%)	6/10 (60.00%)	9/20 (45.00%)	9/20 (45.00%)	9/10 (90.00%)	14/19 (73.68%)	9/62 (14.52%)
	Multifocal	24/32 (75.00%)	2/6 (33.33%)	4/10 (40.00%)	11/20 (55.00%)	11/20 (55.00%)	1/10 (10.00%)	5/19 (26.32%)	53/62 (85.48%)

B)		Frontal bone	Parietal bone (L)	Parietal bone (R)	Temporal bone (L)	Temporal bone (R)	Sphenoid bone (L)	Sphenoid bone (R)	Occipital bone
NTB group (Σ=193)		4/6 (66.67%)	1/6 (16.67%)	1/6 (16.67%)	2/6 (33.33%)	3/6 (50.00%)	1/6 (16.67%)	1/6 (16.67%)	6/6 (100.00%)
Extent (x) of lesions	x < 25%	4/4 (100.00%)	1/1 (100.00%)	1/1 (100.00%)	2/2 (100.00%)	3/3 (100.00%)	1/1 (100.00%)	1/1 (100.00%)	5/6 (83.33%)
	25% ≤ x < 50%	–	–	–	–	–	–	–	–
	50% ≤ x < 75%	–	–	–	–	–	–	–	1/6 (16.67%)
	75% ≤ x	–	–	–	–	–	–	–	–
Number of lesions	Unifocal	–	–	–	–	–	1/1 (100.00%)	1/1 (100.00%)	1/6 (16.67%)
	Multifocal	4/4 (100.00%)	1/1 (100.00%)	1/1 (100.00%)	2/2 (100.00%)	3/3 (100.00%)	–	–	5/6 (83.33%)