

1 S3 Table. Relationship between the detection of *Staphylococcus haemolyticus* and/or  
 2 *Ralstonia pickettii* and antibiotics used during pretransplant conditioning.

	Detection of bacterial taxa	
	<i>Staphylococcus haemolyticus</i> (n= 8)	<i>Ralstonia pickettii</i> (n= 13)
Prophylactic use (n= 19)	0 (0)	2 (15.3)
Additional use for treatment (n= 26)	8 (30.7)	11 (84.6)
CFPM (n= 14)	2 (14.2)	4 (28.4)
CZOP (n= 1)	0 (0)	1 (100.0)
Other antibiotics (n= 11)	6 (54.5)	6 (54.5)
DRPM (n=1)	1 (100)	0 (0)
CFPM+TAZ/PIPC (n= 2)	1 (50)	1 (50)
CZOP+TAZ/PIPC (n= 1)	1 (100)	1 (100)
DRPM+TAZ/PIPC (n= 1)	0 (0)	0 (0)
CFPM+TAZ/PIPC+DRPM (n= 1)	0 (0)	1 (100)
CFPM+TAZ/PIPC+TEIC (n= 1)	1 (100)	1 (100)
MEPM+TEIC (n= 1)	0 (0)	1 (100)
TAZ/PIPC+MEPM+TEIC (n= 2)	2 (100)	1 (50)
CFPM+DRPM+VCM (n= 1)	0 (0)	0 (0)

CFPM, cefepime; CZOP, cefozopran; DRPM, doripenem; MEPM; meropenem; TAZ/PIPC; tazobactam/piperacillin; TEIC; teicoplanin; VCM; vancomycin.