

Supporting Information

Highly efficient gold nano-flowered optical biosensor doped in sol-gel/PEG matrix for the determination of calcitonin biomarker in different serum samples

Walaa. E. Omer¹, Maged A. El-Kemary¹, Mostafa M. Elsaady², Mona N. Abou-Omar³, Ahmed O. Youssef², Ali A. Sayqal⁴, Ayman A. Gouda⁵, Mohamed S. Attia^{2*}

¹Institute of Nanoscience and Nanotechnology, Kafrelsheikh University, 33516 Egypt. ²Department of Chemistry, Faculty of Science, Ain Shams University, 11566, Cairo, Egypt.

³Department of Chemistry, Faculty of Women for Arts, Science and Education, Ain Shams University, Cairo, Egypt.

⁴Chemistry Department, Faculty of Applied Sciences, Umm Al-Qura University, Makkah, Saudi Arabia.

⁵Faculty of Public Health and Health Informatics, Umm Al-Qura University, Makkah, Saudi Arabia.

Corresponding author: (M. S. Attia) Mohamed_sam@yahoo.com,

mohd_mostafa@sci.asu.edu.eg, Tel: +(202) 1229867311, +(202) 1060818922

S1




Figure S.1. Structure of calcitonin

S2




Figure S2. Stern–Völmer plot.

S3




Figure S3. Mechanism of AuNFs formation in HEPES

S4




Figure S4. Mechanism of synthesis of Au doped in sol gel

S1

Table S1. Sensitivity and regression parameters for the proposed method

Parameter	Values
λ_{em} (nm)	360
Linear range, pg/mL	0.01 -1000
Limit of detection (LOD), pg/mL	0.707
Limit of quantification (LOQ), pg/mL	2.14
Intercept (a) $\times 10^{-4}$	5.80
Slope (b)	0.16
Standard deviation	0.03
Regression coefficient (r)	0.99