

Optimization of Docetaxel Loading Conditions in Liposomes: proposing potential products for metastatic breast carcinoma chemotherapy

Roghayyeh Vakili-Ghartavol¹, Seyed Mahdi Rezayat^{1,2}, Reza Faridi-Majidi¹, Kayvan Sadri³, Mahmoud Reza Jaafari*^{4,5}

*Corresponding author: Mahmoud Reza Jaafari; Email: Jafarimr@mums.ac.ir

¹Department of Medical Nanotechnology, School of Advanced Technologies in Medicine, Tehran University of Medical Sciences, Tehran, Iran.

²Department of Pharmacology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran.

³Nuclear Medicine Research Center, Mashhad University of Medical Sciences, Mashhad 98451-3546, Iran

⁴Nanotechnology Research Center, Pharmaceutical Technology Institute, Mashhad University of Medical Sciences, Mashhad, Iran

⁵Department of Pharmaceutical Nanotechnology, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran

Supplementary Table S3

HPLC gradient program based on deionized water and acetonitrile for DTX analysis

Time (min)	F.R (ml/min)	Water (%)	Acetonitrile (%)
0	1	72	28
9	1	72	28
39	1	28	72
39.1	1	72	28
45	1	72	28