

## Optimization of Docetaxel Loading Conditions in Liposomes: proposing potential products for metastatic breast carcinoma chemotherapy

Roghayyeh Vakili-Ghartavol<sup>1</sup>, Seyed Mahdi Rezayat<sup>1,2</sup>, Reza Faridi-Majidi<sup>1</sup>, Kayvan Sadri<sup>3</sup>, Mahmoud Reza Jaafari\*<sup>4,5</sup>

\*Corresponding author: Mahmoud Reza Jaafari; Email: Jafarimr@mums.ac.ir

<sup>1</sup>Department of Medical Nanotechnology, School of Advanced Technologies in Medicine, Tehran University of Medical Sciences, Tehran, Iran.

<sup>2</sup>Department of Pharmacology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran.

<sup>3</sup>Nuclear Medicine Research Center, Mashhad University of Medical Sciences, Mashhad 98451-3546, Iran

<sup>4</sup>Nanotechnology Research Center, Pharmaceutical Technology Institute, Mashhad University of Medical Sciences, Mashhad, Iran

<sup>5</sup>Department of Pharmaceutical Nanotechnology, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran

### Supplementary Table S4

DTX concentrations and correspond Area Under the DTX concentration – time Curves (AUC) obtained from HPLC analysis.

Sample number	AUC	DTX concentration
1	63577833.9	15.6
2	113169183	31.2
3	201391383	62.5
4	359055883	125
5	680722583	250
6	2554465681	1000