

**Pentacyclic triterpenoid-rich fraction of the Hardy kiwi (*Actinidia arguta*)
improves brain dysfunction in high fat diet-induced obese mice**

Jeong Su Ha^{1,#}, Jin Yong Kang^{1,#}, Jeong Eun Kang¹, Seon Kyeong Park¹, Jong Min Kim¹,
Chul-Woo Kim², Sung-Il Oh², Uk Lee², Dae-Ok Kim³ and Ho Jin Heo^{1,*}

***Corresponding author:** Division of Applied Life Science (BK21 plus), Institute of Agriculture and Life Science, Gyeongsang National University, Jinju 52828, Republic of Korea. Tel: +82 55 772 1907; Fax: +82 55 772 1909. E-mail address: hjher@gnu.ac.kr (H. J. Heo).

¹*Division of Applied Life Science (BK21 plus), Institute of Agriculture and Life Science, Gyeongsang National University, Jinju 52828, Republic of Korea*

²*Division of Special Forest Products, National Institute of Forest Science, Suwon 16631, Republic of Korea*

³*Department of Food Science and Biotechnology, Kyung Hee University, Yongin, 17104, Republic of Korea*

JSH: orengewater@naver.com, JYK: kangjy2132@naver.com, JEK: kjd940225@naver.com, SKP: tjrsrud2510@naver.com, JMK: myrock201@naver.com, CWK: futuretree@korea.kr, SIO: osi0919@korea.kr, UL: rich26@korea.kr, DOK: dokim05@khu.ac.kr, HJH: hjher@gnu.ac.kr

[#]Jeong Su Ha and Jin Yong Kang are co-first authors

Supplementary information




Figure S1. Effect of chloroform fraction from *Actinidia arguta* (CFAA) on α-glucosidase activity (A) and α-amylase activity (B). Data were analyzed using ANOVA with Duncan's SAS and expressed as mean ± SD ($n=6$). Data were statistically considered at $p<0.05$, and different small letters represent statistical difference.