

**PERSEVERE Biomarkers Predict Severe Acute Kidney Injury and Renal Recovery
in Pediatric Septic Shock**

Natalja L. Stanski, MD, Erin K. Stenson, MD, Natalie Z. Cvijanovich, MD, Scott L. Weiss, MD, Julie C. Fitzgerald, MD, PhD, Michael T. Bigham, MD, Parag N. Jain, MD, Adam Schwarz, MD, Riad Lutfi, MD, Jeffrey Nowak, MD, Geoffrey L. Allen, MD, Neal J. Thomas, MD, Jocelyn R. Grunwell MD, PhD, Torrey Baines, MD, Michael Quasney, MD, Bereketiab Haileselassie, MD, and Hector R. Wong, MD

Online Data Supplement

Table E1: Patient outcomes by PERSEVERE-II mortality probability risk category.

	PERSEVERE-II Low Risk	PERSEVERE-II Intermediate Risk	PERSEVERE-II High Risk	P value
Total	241 (64)	85 (22)	53 (14)	--
Age (years)*	6.3 (1.9-12.4)	8.5 (3.1-14.3)	4.8 (1.6-11)	0.27
Gender, n (% male)	123 (51)	45 (53)	27 (51)	0.952
Mortality	5 (2)	15 (18)	22 (42)	<0.001
Day 3 Severe AKI	24 (10)	16 (19)	24 (45)	<0.001
RRT Use	9 (4)	12 (14)	17 (32)	<0.001
Day 1-2 AKI	60 (25)	38 (45)	33 (62)	<0.001
Day 3 Renal Recovery	29 (48)	17 (45)	8 (24)	0.039

*Categorical variables are reported as n (%) and continuous variables as median (IQR)