

Supplementary Table 4

Characteristics of the patients lost to follow-up

	Patients lost to follow up	Patients not lost to follow-up	p value
Total	18	1117	
Female n (%)	13 (72.2%)	660 (59.1%)	0.260
Age mean (SD)	40 (18)	39 (17)	0.568
HIV positive/HIV status known (%)	8/16 (50.0%)	359/830 (43.4%)	0.590
Had surgery in hospital n (%)	3 (16.7%)	262 (23.5%)	0.499
Specialty n (%)			0.018
Medicine	8 (44.4%)	465 (41.6%)	
Surgery	2 (11.1%)	317 (28.4%)	
Obstetrics & Gynaecology	6 (33.3%)	286 (25.6%)	
Ophthalmology	2 (11.1%)	18 (1.6%)	
Other	0	31 (2.8%)	
Length of hospital stay			
Median (IQR) in days			
Before data collection	3 (1-7)	5 (2-12)	0.499
After data collection ¹ Total	2 (2-2)	4 (2-10)	0.483
Length of Stay	9 (9-9)	11 (5-22)	0.833
Infection diagnosis n (%)	6 (33.3%)	413 (37.0%)	0.751
qSOFA score ≥2 n (%)	1 (5.6%)	141 (12.6%)	0.369
Sepsis* n (%)	1 (5.6%)	80 (7.2%)	0.793

* sepsis defined as infection diagnosis and qSOFA score ≥2

¹ censored at 30 days