

Supplementary Online Content

Baghdadi JD, Brook RH, Uslan DZ, et al. Association of a care bundle for early sepsis management with mortality among patients with hospital-onset or community-onset sepsis. *JAMA Intern Med*. Published online April 6, 2020.
doi:10.1001/jamainternmed.2020.0183

eTable 1. Pairwise Correlation Between Individual SEP-1 Sepsis Bundle Components for Patients with Community-onset Sepsis and Hospital-onset Sepsis

eTable 2. Relative Risk of In-hospital Mortality Associated with Clinical Variables in the Raw Cohorts with Community-onset and Hospital-onset Sepsis

eTable 3. Characteristics of the Total Sample Before and After Being Rebalanced on Propensity for Treatment and Observable Characteristics

eTable 4. Relative Risk of In-hospital Mortality Associated with Clinical Variables in the Raw Total Sample

eTable 5. Results from Endogenous Treatment Effects Models Compared with Unweighted/Unbalanced Regression

eTable 6. Prespecified Sensitivity Analyses Evaluating Treatment Effect of the Complete SEP-1 Bundle in Different Cohorts

eTable 7. Treatment Effects of SEP-1 Bundle Components in Different Cohorts

eTable 8. Treatment Effects of SEP-1 Sepsis Bundle and Components on 30-Day In-hospital Mortality in Total Sample and Rebalanced Cohorts with Community-onset and Hospital-onset Sepsis

eTable 9. Treatment Effects of SEP-1 Sepsis Bundle and Components on Hospital Length of Stay (Measured from Time Zero of Sepsis until Discharge)

eTable 10. Treatment Effects of SEP-1 Sepsis Bundle and Components on Composite Outcome of In-hospital Mortality or Discharge to Hospice

eTable 11. Treatment Effects of SEP-1 Sepsis Bundle and Components in Rebalanced Cohorts with Sepsis Presenting on the Inpatient Ward or Intensive Care Unit

eTable 12. Treatment Effects of SEP-1 Sepsis Bundle and Components in Rebalanced Cohorts with Time Zero of Sepsis <48 or >48 Hours After Arrival

eTable 13. Post Hoc Sensitivity Analyses

This supplementary material has been provided by the authors to give readers additional information about their work.

1 **eTable 1.** Pairwise Correlation Between Individual SEP-1 Sepsis Bundle Components for
 2 Patients with Community-onset Sepsis and Hospital-onset Sepsis
 3

Community-Onset Sepsis					
	Blood cultures	Lactate	Antibiotics	IV Fluids	2 nd Lactate
Lactate	0.57				
Antibiotics	0.61	0.42			
IV Fluids	-0.037	-0.045	-0.022		
2 nd Lactate	0.065	0.045	0.029	-0.12	
Vasopressors	0.01	0.012	-0.015	0.012	0.024
Hospital-Onset Sepsis					
	Blood cultures	Lactate	Antibiotics	IV Fluids	2 nd Lactate
Lactate	0.33				
Antibiotics	0.66	0.28			
IV Fluids	-0.038	-0.10	-0.0396		
2 nd Lactate	0.068	0.036	0.098	-0.031	
Vasopressors	0.005	0.015	0.082	0.082	-0.001

4
 5 **Legend:** Community-onset sepsis was defined by time zero of sepsis occurring in the emergency
 6 department. Hospital-onset sepsis was defined by time zero of sepsis occurring on an inpatient
 7 unit even if signs or symptoms of sepsis preceded admission.

8 **eTable 2.** Relative Risk of In-hospital Mortality Associated with Clinical Variables in the Raw
 9 Cohorts with Community-onset and Hospital-onset Sepsis
 10

	Community-Onset Sepsis n = 4108		Hospital-Onset Sepsis n = 2296	
	Relative Risk	p-value	Relative Risk	p-value
Age in years	1.01	< 0.001	1.01	0.001
ECI	1.06	< 0.001	1.08	< 0.001
Female	1.03	0.60	1.08	0.36
Postoperative	1.21	0.35	1.05	0.69
Pneumonia POA	1.41	< 0.001	1.19	0.09
UTI POA	0.66	< 0.001	0.68	0.002
SSTI POA	0.83	0.21	0.93	0.70
New Bacteremia	1.04	0.66	1.28	0.032
Immunosuppressed	1.25	0.005	1.26	0.008
Fever	0.47	< 0.001	0.64	< 0.001
Septic Shock	1.84	< 0.001	1.27	0.005
- Hypotension	0.92	0.26	0.97	0.73
- Elevated Lactate	0.83	0.033	1.25	0.10

11 **Legend:** POA, present on admission. UTI, urinary tract infection. SSTI, skin and/or soft tissue
 12 infection. Bold font denotes a statistically significant standardized difference for an $\alpha = 0.05$.
 13

14
15 **eTable 3.** Characteristics of the Total Sample Before and After Being Rebalanced on Propensity
16 for Treatment and Observable Characteristics
17

	Raw Total Sample			Rebalanced Total Sample		
	Treated	Untreated	SD	Treated	Untreated	SD
Time Zero of Sepsis (Hours*)	8.97	34.3	-0.33	6.55	8.45	-0.025
Age	65.6	63.3	0.13	65.9	66.1	-0.011
Female (%)	43.7	45.3	-0.033	43.6	44.68	-0.003
Elixhauser Comorbidities	4.91	5.14	-0.076	4.84	4.85	-0.003
Postoperative (%)	3.01	9.67	-0.28	1.98	1.98	0
Pneumonia POA† (%)	34.0	25.3	0.19	34.0	34.0	0
Urinary tract infection (%)	25.4	21.2	0.10	26.0	26.0	0
Skin / soft tissue infection (%)	7.99	6.03	0.077	7.88	6.53	0.053
New Bacteremia‡ (%)	14.6	14.5	0.003	14.6	15.8	-0.033
Immunosuppressed (%)	19.2	24.7	-0.13	18.3	18.3	0
Fever (%)	36.8	22.7	0.31	35.5	35.5	0
Septic Shock (%)	27.5	42.7	-0.32	27.2	27.2	0
- Hypotension	35.5	55.0	-0.40	35.8	35.8	0
- Elevated Serum Lactate***	28.7	17.2	0.28	29.1	27.8	0.029
Hospital 1 (%)	9.85	11.6	-0.055	9.49	9.49	0
Hospital 2 (%)	13.5	17.3	-0.10	13.2	13.2	0
Hospital 3 (%)	32.7	41.3	-0.18	32.9	32.9	0
Hospital 4 (%)	43.9	29.9	0.29	44.5	44.5	0

18
19 **Legend:** SD, standardized difference. POA, present on admission. Bold font denotes a
20 statistically significant standardized difference for an $\alpha = 0.05$. * Hours from admission. †
21 Pneumonia POA was determined by diagnosis codes. ‡ New bacteremia was defined by positive
22 blood culture after time zero of sepsis in a patient who did not have a positive blood culture in
23 the preceding week. Possible skin contaminants were excluded. *** Elevated serum lactate was
24 defined as a value of >20 mmol/L at time of onset of sepsis or in the preceding 6 hours. To meet
25 the definition of shock, serum lactate needed to be 4 mmol/L or greater.

26 **eTable 4.** Relative Risk of In-hospital Mortality Associated with Clinical Variables in the Raw
 27 Total Sample
 28

	Relative Risk	95% Confidence Interval	p-value
SEP-1 Sepsis Bundle	0.99	(0.88, 1.12)	0.88
Hospital-Onset of Sepsis	1.05	(0.94, 1.17)	0.381
Age in years*	1.01	(1.01, 1.01)	< 0.001
Female	1.06	(0.96, 1.17)	0.28
Elixhauser Comorbidities*	1.07	(1.05, 1.08)	< 0.001
Postoperative	1.35	(0.88, 1.30)	0.48
Pneumonia POA	1.35	(1.21, 1.50)	< 0.001
Urinary Tract infection POA	0.67	(0.58, 0.77)	< 0.001
Skin / soft tissue infection POA	0.85	(0.68, 1.06)	0.16
New Bacteremia	1.12	(0.98, 1.28)	0.10
Immunosuppressed	1.25	(1.11, 1.40)	< 0.001
Fever	0.51	(0.44, 0.59)	< 0.001
Septic Shock	1.55	(1.40, 1.72)	< 0.001

29
 30 **Legend:** POA, present on admission. * Relative risk is reported for incremental increase in 1
 31 year of age or 1 additional Elixhauser comorbidity.

32 **eTable 5.** Results from Endogenous Treatment Effects Models Compared with
 33 Unweighted/Unbalanced Regression

34

Treatment		ATET	NATE
SEP-1	MD	+0.10	-0.23
	VD	+0.26	-0.60
Blood Cultures	MD	-5.01	+0.97
	VD	-0.44	-0.28
Serum Lactate	MD	-2.68	+1.26
	VD	-0.004	+0.28
Antibiotics	MD	-3.83	-1.03
	VD	-0.45	-0.27
IV Fluids	MD	-0.57	+2.64
	VD	-0.23	-0.10

35

36 **Legend:** ATET, average treatment effect on the treated. NATE, naïve (i.e., unweighted /
 37 unbalanced) treatment effect. MD, Mortality difference calculated as the average difference in
 38 probabilities between the treated and untreated (unit is percentage points); VD, vasopressor days;
 39 IV, intravenous.

40 **eTable 6.** Prespecified Sensitivity Analyses Evaluating Treatment Effect of the Complete SEP-1
 41 Bundle in Different Cohorts*
 42

Subgroup	Mortality Difference[†]	Vasopressor Days
Age \geq 65	-0.01% (-3.88, 3.86)	+0.37 (0.14, 0.60)
Age < 65	+1.94% (-1.37, 5.26)	+0.48 (0.21, 0.75)
Pneumonia POA	-0.96% (-6.70, 4.77)	+0.59 (0.26, 0.92)
No Pneumonia POA	+1.30% (-1.40, 4.00)	+0.32 (0.11, 0.52)
Immunosuppressed	+4.63% (-1.53, 10.8)	+0.86 (0.45, 1.28)
Not Immunosuppressed	-0.61% (-3.42, 2.21)	+0.26 (0.07, 0.45)
New Bacteremia	+1.35% (-5.03, 7.72)	+0.45 (-0.09, 0.99)
No New Bacteremia	+0.23% (-2.62, 3.07)	+0.38 (0.20, 0.57)
Fever	+2.30% (-0.97, 5.56)	+0.33 (0.07, 0.59)
Afebrile	+0.53% (-2.93, 3.99)	+0.40 (0.18, 0.62)
Septic Shock	+2.19% (-3.05, 7.43)	+0.55 (0.14, 0.97)
Sepsis without Shock	-0.38% (-3.58, 2.82)	0.01 (-0.16, 0.19)

43
 44 **Legend:** SEP-1 refers to the Early Management Bundle for Severe Sepsis / Septic Shock
 45 (National Quality Forum #0500). * Dedicated propensity and outcomes models were run within
 46 the subgroup of interest. Subgroup analyses excluded all encounters from outside the subgroup. †
 47 Difference in probability of in-hospital death was estimated as the average treatment effect on
 48 the treated and is expressed in percentage points.

49 **eTable 7.** Treatment Effects of SEP-1 Bundle Components in Different Cohorts*
 50

		Blood Cultures	Lactate	Antibiotics	IV Fluids†
Age ≥ 65 n = 2459	MD	-9.58 (-200, 0.86)	-3.51 (-9.06, 2.03)	-3.84 (-8.81, 1.14)	-3.83 (-12.2, 4.51)
	VD	-0.88 (-1.69, -0.07)	0.15 (-0.28, 0.59)	-0.42 (-0.82, -0.01)	-0.18 (-0.86, 0.49)
Age < 65 n = 2854	MD	-2.37 (-8.19, 3.45)	-1.18 (-5.73, 3.38)	1.40 (-2.60, 5.40)	5.61 (-1.02, 12.2)
	VD	-0.48 (-1.00, 0.04)	-0.06 (-0.47, 0.35)	-0.18 (-0.54, 0.18)	-0.13 (-0.83, 0.57)
Pneumonia POA n = 1613	MD	-10.3 (-25.5, 4.91)	-2.41 (-11.0, 6.18)	-3.38 (-10.9, 4.13)	1.95 (-15.9, 19.8)
	VD	-0.26 (-0.90, 0.85)	0.31 (-0.31, 0.94)	-0.08 (-0.63, 0.46)	-0.14 (-1.72, 1.44)
No Pneumonia POA n = 3975	MD	-2.75 (-8.01, 2.50)	-2.27 (-6.43, 1.88)	-3.60 (-8.00, 0.82)	-2.61 (-8.83, 3.62)
	VD	-0.58 (-1.08, -0.07)	-0.09 (-0.44, 0.27)	-0.61 (-0.99, -0.22)	-0.35 (-0.90, 0.21)
Immunosuppressed n = 1376	MD	-3.60 (-14.7, 7.50)	-1.21 (-10.3, 7.85)	0.08 (-8.32, 8.47)	12.4 (-3.59, 28.4)
	VD	-0.24 (-0.99, 0.50)	0.25 (-0.40, 0.89)	-0.75 (-1.50, -0.001)	0.22 (-1.46, 1.91)
Non-immunosuppressed n = 4352	MD	-5.76 (-13.0, 1.47)	-3.01 (-7.39, 1.36)	-4.01 (-8.02, 0.01)	-1.97 (-8.18, 4.24)
	VD	-0.79 (-1.42, -0.16)	-0.03 (-0.38, 0.31)	-0.38 (-0.70, -0.06)	-0.43 (-0.97, 0.11)
New Bacteremia n = 738	MD	-13.9 (-28.4, 0.53)	-4.98 (-18.1, 8.12)	-4.33 (-12.5, 3.81)	-1.43 (-19.5, 16.6)
	VD	-2.11 (-3.53, -0.69)	-0.24 (-1.22, 0.74)	-1.34 (-2.08, -0.60)	-0.07 (-1.71, 1.58)
No New Bacteremia n = 5476	MD	-2.34 (-8.40, 3.73)	-2.11 (-6.13, 1.92)	-2.80 (-6.38, 0.77)	-0.04 (-6.22, 6.14)
	VD	-0.28 (-0.76, 0.21)	0.18 (-0.15, 0.51)	-0.20 (-0.49, 0.09)	-0.16 (-0.71, 0.39)
Fever n = 1722	MD	0.23 (-17.3, 17.7)	-4.62 (-14.0, 4.72)	-2.03 (-8.54, 4.48)	3.60 (-3.70, 10.9)
	VD	0.81 (-0.26, 1.87)	0.02 (-0.66, 0.69)	-0.25 (-0.77, 0.28)	0.28 (-0.68, 1.24)
Afebrile n = 4225	MD	-5.22 (-11.1, 0.64)	-3.47 (-7.92, 0.98)	-2.77 (-6.69, 1.14)	0.47 (-7.05, 7.98)
	VD	-0.61 (-1.08, -0.15)	0.01 (-0.31, 0.34)	-0.38 (-0.69, -0.06)	-0.36 (-1.03, 0.32)
Septic Shock n = 2248	MD	-4.58 (-14.5, 5.29)	-3.39 (-12.9, 6.20)	-6.54 (-13.7, 0.61)	-6.30 (-14.5, 1.94)
	VD	-0.55 (-1.50, 0.40)	0.11 (-0.62, 0.83)	-0.45 (-1.00, 0.10)	-0.61 (-1.32, 0.11)
Sepsis without Shock n = 3164	MD	-6.42 (-13.0, 0.11)	-3.63 (-7.95, 0.68)	-2.45 (-5.99, 1.10)	2.66 (-6.58, 11.9)
	VD	-0.36 (-0.71, -0.003)	-0.17 (-0.47, 0.13)	-0.11 (-0.31, 0.08)	0.19 (-0.40, 0.79)

51
 52 **Legend:** MD, Mortality difference calculated as the average difference in probabilities between
 53 the treated and untreated (unit is percentage points); VD, vasopressor days; 95% CI, 95%
 54 confidence interval. POA, present on admission. * Dedicated propensity and outcomes models
 55 were run within the subgroup of interest. Subgroup analyses excluded all encounters from
 56 outside the subgroup. † Only measured among individuals in whom intravenous fluids were
 57 indicated based on SEP-1 guidance

58 **eTable 8.** Treatment Effects of SEP-1 Sepsis Bundle and Components on 30-day In-hospital
 59 Mortality in Total Sample and Rebalanced Cohorts with Community-onset & Hospital-onset
 60 Sepsis
 61

	Total Sample	Community-Onset Sepsis	Hospital-Onset Sepsis
	Risk Difference* (95% CI)	Risk Difference* (95% CI)	Risk Difference* (95% CI)
SEP-1 Bundle	+0.49 (-1.93, 2.90)	+0.67 (-2.16, 3.49)	+0.91 (-5.29, 7.11)
Blood cultures	-5.13 (-11.0, 0.74)	-5.98 (-15.2, 3.25)	-2.43 (-8.31, 3.45)
Serum Lactate	-1.86 (-5.63, 1.92)	-5.76 (-12.4, 0.88)	+0.44 (-4.94, 4.06)
Antibiotics	-3.78 (-7.67, 0.11)	-3.72 (-8.90, 1.47)	-4.12 (-8.99, 0.74)
Intravenous Crystalloid	+1.14 (-4.07, 6.36)	+0.84 (-5.91, 7.59)	-4.63 (-15.1, 5.84)

62
 63 **Legend:** 95% CI, 95% confidence interval. SEP-1 refers to the Early Management Bundle for
 64 Severe Sepsis / Septic Shock (National Quality Forum #0500). * Mortality is expressed as an
 65 absolute probability difference between the average of the treated and the average of the
 66 untreated, expressed in terms of percentage points.

67 **eTable 9.** Treatment Effects of SEP-1 Sepsis Bundle and Components on Hospital Length of
 68 Stay (Measured from Time Zero of Sepsis Until Discharge)
 69

	Total Sample	Community-Onset Sepsis	Hospital-Onset Sepsis
	Length of Stay* (95% CI)	Length of Stay* (95% CI)	Length of Stay* (95% CI)
SEP-1 Bundle	-0.54 (-1.41, 0.33)	-0.79 (-1.93, 0.35)	-1.08 (-3.42, 1.27)
Blood cultures	-0.05 (-1.78, 1.69)	+1.43 (-0.72, 3.58)	-1.35 (-3.99, 1.28)
Serum Lactate	-0.28 (-1.43, 0.87)	-0.47 (-2.63, 1.69)	-0.99 (-2.84, 0.85)
Antibiotics	+0.26 (-0.65, 1.17)	+0.78 (-0.38, 1.93)	+0.42 (-1.28, 2.12)
Intravenous Crystalloid	-1.15 (-2.86, 0.55)	-0.71 (-2.73, 1.30)	+1.49 (-1.41, 4.38)

70
 71 **Legend:** 95% CI, 95% confidence interval. SEP-1 refers to the Early Management Bundle for
 72 Severe Sepsis / Septic Shock (National Quality Forum #0500). * Length of stay is expressed in
 73 days from time zero of sepsis until discharge.

74 **eTable 10.** Treatment Effects of SEP-1 Sepsis Bundle and Components on Composite Outcome
 75 of In-hospital Mortality or Discharge to Hospice
 76

	Total Sample	Community-Onset Sepsis	Hospital-Onset Sepsis
	Risk Difference* (95% CI)	Risk Difference* (95% CI)	Risk Difference* (95% CI)
SEP-1 Bundle	-0.43 (-3.37, 2.52)	+0.08 (-3.39, 3.23)	-2.97% (-9.91, 3.96)
Blood cultures	-5.00 (-11.0, 1.04)	-10.2% (-19.3, -1.01)	+0.31% (-5.88, 6.49)
Serum Lactate	-1.35 (-5.28, 2.58)	-6.14 (-13.3, 1.01)	-1.01% (-5.96, 3.94)
Antibiotics	-3.58 (-7.52, 0.36)	-3.16 (-8.40, 2.08)	-4.78% (-9.70, 0.14)
Intravenous Crystalloid	-2.18 (-7.74, 3.38)	-4.23 (-11.6, 3.10)	-6.08 (-17.4, 5.25)

77
 78 **Legend:** 95% CI, 95% confidence interval. SEP-1 refers to the Early Management Bundle for
 79 Severe Sepsis / Septic Shock (National Quality Forum #0500). * Risk differences were estimated
 80 as the average treatment effect on the treated and are expressed in terms of percentage points.

81 **eTable 11.** Treatment Effects of SEP-1 Sepsis Bundle and Components in Rebalanced Cohorts
 82 with Sepsis Presenting on the Inpatient Ward or Intensive Care Unit
 83

	Ward-Onset Sepsis (n = 987)		ICU-Onset Sepsis (n = 1,128)	
	Mortality Risk Difference*	95% CI	Mortality Risk Difference*	95% CI
SEP-1 Bundle	+1.70	(-15.2, 18.6)	-0.50	(-11.2, 10.2)
Blood cultures	-8.11	(-19.3, 3.04)	-1.78	(-12.6, 9.11)
Serum Lactate	+3.45	(-5.75, 12.6)	-1.30	(-10.2, 7.5)
Antibiotics	-5.79	(-14.1, 2.54)	-5.81	(-14.4, 2.83)
Intravenous Crystalloid	Utd	Utd	-4.76	(-21.6, 12.1)
	Vasopressor Days*	95% CI	Vasopressor Days*	95% CI
SEP-1 Bundle	+0.53	(-0.16, 1.22)	1.07	(0.14, 2.00)
Blood cultures	+0.29	(-0.22, 0.81)	-0.33	(-1.23, 0.57)
Serum Lactate	+0.70	(0.01, 1.40)	+0.34	(-0.34, 1.02)
Antibiotics	+0.29	(-0.11, 0.70)	-0.22	(-0.93, 0.49)
Intravenous Crystalloid	Utd	Utd	+0.35	(-0.93, 1.63)

84
 85 **Legend:** ICU, intensive care unit. Utd, unable to determine. 95% CI, 95% confidence interval.
 86 SEP-1 refers to the Early Management Bundle for Severe Sepsis / Septic Shock (National
 87 Quality Forum #0500). * Differences in probability of in-hospital mortality and days spent on
 88 vasopressors were estimated as the average treatment effect on the treated. Mortality risk
 89 difference was expressed in terms of percentage points.

90 **eTable 12.** Treatment Effects of SEP-1 Sepsis Bundle and Components in Rebalanced Cohorts
 91 with Time Zero of Sepsis <48 or >48 Hours After Arrival
 92

	Time Zero < 48 Hours After Arrival		Time Zero > 48 Hours After Arrival	
	Mortality Risk Difference*	95% CI	Mortality Risk Difference*	95% CI
SEP-1 Bundle	+0.16	(-2.39, 2.72)	1.26	(-22.3, 24.8)
Blood cultures	-6.08	(-13.0, 0.85)	-4.16	(-15.1, 6.81)
Serum Lactate	-5.26	(-9.82, -0.70)	+7.77	(-4.80, 20.3)
Antibiotics	-4.33	(-8.33, -0.32)	-5.67	(-16.1, 4.79)
Intravenous Crystalloid	-0.92	(-7.29, 5.46)	+34.2	(-15.2, 84.4)
	Vasopressor Days*	95% CI	Vasopressor Days*	95% CI
SEP-1 Bundle	+0.37	(-0.02, 0.03)	0.13	(-1.38, 1.64)
Blood cultures	-0.64	(-1.19, -0.09)	-0.28	(-1.20, 0.64)
Serum Lactate	-0.10	(-0.45, 0.25)	0.82	(-0.04, 1.69)
Antibiotics	-0.45	(-0.77, -0.12)	-0.48	(-1.44, 0.48)
Intravenous Crystalloid	-0.48	(-1.00, 0.05)	0	(0, 0)

93
 94 **Legend:** 95% CI, 95% confidence interval. SEP-1 refers to the Early Management Bundle for
 95 Severe Sepsis / Septic Shock (National Quality Forum #0500). * Differences in probability of in-
 96 hospital mortality and days spent on vasopressors were estimated as the average treatment effect
 97 on the treated. Mortality risk difference was expressed in terms of percentage points.

98 **eTable 13.** Post Hoc Sensitivity Analyses

99

	Admitted from Home (n = 3732 matched)		Source Other than Home (n = 2077 matched)	
	Mortality Risk Difference*	95% CI	Mortality Risk Difference*	95% CI
SEP-1 Bundle	-0.04	(-3.13, 3.05)	1.33	(-3.31, 5.97)
Blood cultures	-0.59	(-5.99, 4.82)	-7.63	(-16.4, 1.14)
Serum Lactate	-4.27	(-9.70, 1.17)	-4.46	(-11.6, 2.68)
Antibiotics	-2.66	(-6.93, 1.62)	-5.28	(-11.5, 0.91)
Intravenous Crystalloid	-3.02	(-10.2, 4.21)	0.32	(-6.23, 6.87)
	Vasopressor Days	95% CI	Vasopressor Days	95% CI
SEP-1 Bundle	0.33	(0.13, 0.54)	0.59	(0.27, 0.90)
Blood cultures	-0.54	(-1.12, 0.04)	-0.40	(-1.03, 0.23)
Serum Lactate	0.02	(-0.38, 0.41)	0.01	(-0.55, 0.58)
Antibiotics	-0.45	(-0.79, -0.10)	-0.09	(-0.51, 0.33)
Intravenous Crystalloid	-0.51	(-1.15, 0.14)	-0.10	(-0.65, 0.45)
	Malignancy Absent (n = 4521 matched)		Malignancy Present (n = 978 matched)	
	Mortality Risk Difference*	95% CI	Mortality Risk Difference*	95% CI
SEP-1 Bundle	1.11	(-1.60, 3.83)	-1.23	(-8.81, 6.35)
Blood cultures	-6.01	(-12.6, 0.60)	-8.46	(-21.8, 4.92)
Serum Lactate	-0.66	(-4.59, 3.28)	-10.0	(-22.0, 0.10)
Antibiotics	-3.89	(-8.04, 0.26)	0.69	(-8.58, 9.96)
Intravenous Crystalloid	0.32	(-6.23, 6.87)	-2.01	(-16.1, 12.0)
	Vasopressor Days*	95% CI	Vasopressor Days*	95% CI
SEP-1 Bundle	0.46	(0.27, 0.65)	0.17	(-0.32, 0.66)
Blood cultures	-0.40	(-0.86, 0.06)	-0.78	(-1.84, 0.29)
Serum Lactate	0.09	(-0.25, 0.43)	-0.22	(-0.96, 0.51)
Antibiotics	-0.48	(-0.83, -0.14)	-0.11	(-0.88, 0.67)
Intravenous Crystalloid	-0.10	(-0.65, 0.45)	-0.39	(-1.65, 0.87)

100

101 **Legend:** 95% CI, 95% confidence interval. SEP-1 refers to the Early Management Bundle for
102 Severe Sepsis / Septic Shock (National Quality Forum #0500). * Differences in probability of in-
103 hospital mortality and days spent on vasopressors were estimated as the average treatment effect
104 on the treated. Mortality risk difference was expressed in terms of percentage points.