

Supporting information for

Rapid and Efficient Isolation of Exosomes by Clustering and Scattering

Jinhyun Kim^{1‡}, Hoyoon Lee^{1‡}, KyongHwa Park^{2,3}, and Sehyun Shin^{1,3,*}

¹ School of Mechanical engineering, Korea University, Seoul, Korea; mimoz742@korea.ac.kr (J.K.); leehoyoon@korea.ac.kr (H.L.)

² Department of Internal Medicine, Anam Hospital, Korea University, Seoul, 02841, Korea; khpark@korea.ac.kr (K.H.P.)

³ Engineering Research Center for Biofluid Biopsy, Seoul, 02841, Korea; lexerdshin@korea.ac.kr (S.S.)

‡ Jinhyun Kim and Hoyoon Lee contributed equally to the article as first authors

* Correspondence: lexerdshin@korea.ac.kr; Tel.: +82-10-4506-2825









Incubation Time	10 min	30 min	60 min	90 min
24°C				
4°C				

Fig. S1 EV cluster formation according to incubation time and temperature; The input PLL concentration was fixed at 0.5 mg/mL.




Fig. S2 Comparison of normal plasma and EV depleted plasma via pellet volume




Fig. S3 Comparison of zeta potential results of representative plasma proteins: fibrinogen, globulin, and albumin