





Novel coronavirus infection in newborn babies aged <28 days in China

Zhi-Jiang Zhang^{1,5}, Xue-Jie Yu^{1,5}, Tao Fu ^{2,5}, Yu Liu³, Yan Jiang², Bing Xiang Yang ⁴ and Yongyi Bi¹

Affiliations: ¹Dept of Preventive Medicine, School of Health Sciences, Wuhan University, Wuhan, China. ²Renmin Hospital of Wuhan University, Wuhan, China. ³Dept of Statistics, School of Management, Wuhan Institute of Technology, Wuhan, China. ⁴Dept of Nursing, School of Health Sciences, Wuhan University, Wuhan, China. ⁵Contributed equally.

Correspondence: Zhi-Jiang Zhang, Dept of Preventive Medicine, School of Health Sciences, Wuhan University, No. 185 Donghu Road, Wuhan 430071, China. E-mail: zhang22968@163.com

 @ERSpublications

Neonates are susceptible to SARS-CoV-2 infection. The symptoms in neonates were milder and outcomes were less severe compared to adults. Intrauterine vertical transmission is not impossible but direct evidence is still lacking. <https://bit.ly/342jGK9>

Cite this article as: Zhang Z-J, Yu X-J, Fu T, *et al.* Novel coronavirus infection in newborn babies aged <28 days in China. *Eur Respir J* 2020; 55: 2000697 [<https://doi.org/10.1183/13993003.00697-2020>].

This single-page version can be shared freely online.

To the Editor:

The outbreak of coronavirus disease 2019 (COVID-19) is spreading rapidly [1–8]. On 31 January 2020, the World Health Organization declared COVID-19 a public health emergency of international concern. By 13 March 2020, COVID-19 had been confirmed in 127 countries, with 145 166 cases and 5428 deaths worldwide.