

The supplemental files

Table S1. Crystal data and structure refinement for Rsg-Met.

Empirical formula	C ₂₂ H ₃₀ N ₈ O ₃ S
Formula weight	486.60
Crystal system	triclinic
Space group	P-1
a/Å	6.9482
b/Å	11.3147
c/Å	11.3147
α/°	72.375
β/°	82.370
γ/°	74.153
Volume/Å ³	1240.82
Z	2
D _{calc} g/cm ³	1.302
μ/mm ⁻¹	1.493
2θ range for data collection/°	8.46 to 134.146
Reflections collected	16699
Independent reflections	4433 [<i>R</i> _{int} = 0.0390, <i>R</i> _{sigma} = 0.0305]
Data/restraints/parameters	4433/18/334
Goodness-of-fit on F ²	1.026
Final R indexes [<i>I</i> > 2σ (<i>I</i>)]	<i>R</i> ₁ = 0.0523, <i>wR</i> ₂ = 0.1428
Final R indexes [all data]	<i>R</i> ₁ = 0.0629, <i>wR</i> ₂ = 0.1550
Largest diff. peak/hole/e Å ⁻³	0.43/-0.28
CCDC no.	1962489




Figure S1. The FT-IR spectrum of Rsg




Figure S2. The FT-IR spectrum of Met




Figure S3. The FT-IR spectrum of Rsg-Met