

Salinomycin-Loaded Iron Oxide Nanoparticles for Glioblastoma Therapy

Mohammad Norouzi ^{1,2}, Vinith Yathindranath ², James A. Thliveris ³ and Donald. W Miller ^{1,2,*†}

¹ Department of Biomedical Engineering, University of Manitoba, Winnipeg, MB R3T 5V6 , Canada; norouzim@myumanitoba.ca


² Department of Pharmacology and Therapeutics, University of Manitoba, Winnipeg, MB R3E 0Z3, Canada; Vinith.Yathindranath@umanitoba.ca

³ Department of Human Anatomy and Cell Science University of Manitoba, Winnipeg, MB R3E 0J9, Canada; james.thliveris@gmail.com

* Correspondence: donald.miller@umanitoba.ca; Tel.: +1-204-789-3278

† Current address: Department of Pharmacology & Therapeutics, University of Manitoba, A205 Chown Bldg., 753 McDermot Avenue, Winnipeg, MB R3E 0Z3, Canada

Received: 21 January 2020; Accepted: 03 March 2020; Published: date



Element	Weight %	Atomic %	Error %
C K	10.48	22.30	11.46
N K	5.04	9.20	14.62
O K	24.73	39.49	8.11
AlK	0.69	0.65	22.42
SiK	0.95	0.86	15.47
ClK	3.47	2.50	10.13
FeK	54.63	24.99	3.72

Figure S1. Energy-dispersive X-ray spectroscopy of PEI-PEG-IONPs.





Figure S2. DLS graph of PEI-PEG-IONPs measured in D.I. water (pH 7.2).





Figure S3. TEM images of IONPs (top) and Sali-IONPs (bottom) uptake by bEnd.3 cells after 4 h of the treatment.