

SUPPLEMENTARY TABLE S1. DEMOGRAPHIC AND CLINICAL CHARACTERISTICS OF SAMPLE (N=171)

Mean (SD) age in years (SD)	8.44 (1.26)
Female, <i>n</i> (%)	49 (29)
Race/ethnicity ^a , <i>n</i> (%)	
White, non-Hispanic	139 (81)
African American, Hispanic, other	28 (16) ^a
ADHD subtype, <i>n</i> (%)	
Inattentive type	126 (74)
Combined type	45 (26)
Met criteria for any anxiety disorder ^b on the DISC, <i>n</i> (%)	21 (12)
Met criteria for a mood disorder ^c on the DISC, <i>n</i> (%)	2 (1)
Met criteria for a disruptive behavior disorder ^d on the DISC, <i>n</i> (%)	45 (26)
Titration dose schedules (by weight), <i>n</i> (%)	
Low weight (≤25 kg)	137 (80)
High weight (>25 kg)	34 (20)
VADPRS anxiety/depression score, mean (SD) [range: 0–21]	5.25 (4.70)
VADPRS oppositional defiant score, mean (SD) [range: 0–24]	7.95 (6.67)

^aReported by a parent/caregiver, four participants were missing information about race/ethnicity.

^bSocial phobia, separation anxiety, panic agoraphobia, generalized anxiety disorder, obsessive compulsive disorder, and/or posttraumatic stress disorder.

^cMajor depressive episode/dysthymia.

^dOppositional defiant or conduct disorder.

ADHD, attention-deficit/hyperactivity disorder; DISC, diagnostic interview schedule for children; SD, standard deviation; VADPRS, Vanderbilt ADHD Diagnostic Parent Rating Scale.