

H5N1 infection of the respiratory tract and beyond: a molecular pathology study

Correspondence to:
Prof Jiang Gu, Infectious Disease
Centre, Peking University,
Beijing, 100083 China
jianggu@bjmu.edu.cn

	Supplier	Specificity	Dilution	Incubation temperature	Incubation period
CD3	Zymed Laboratories, San Francisco, CA, USA	T cell	1:100	4°C	Overnight
CD20	Zymed Laboratories	B cell	1:100	4°C	Overnight
CD68	Zymed Laboratories	Macrophage, monocyte	1:100	4°C	Overnight
CD8	DakoCytomation, Glostrup, Denmark	Cytotoxic T cell	1:50	4°C	Overnight
Cytokeratin (AE1/AE3)	Zymed Laboratories	Epithelial cell	1:100	4°C	Overnight
Surfactant protein A	DakoCytomation	Type II pneumocyte	1:100	4°C	Overnight
Tubulin-β	Zymed Laboratories	Ciliated cell	1:100	4°C	Overnight
Factor VIII	Zymed Laboratories	Endothelial cell	1:100	4°C	Overnight
PLAP	Zymed Laboratories	Placental syncytiotrophoblast	1:100	4°C	Overnight
E-cadherin	Zymed Laboratories	Placental cytotrophoblast	1:100	4°C	Overnight
Haemagglutinin	Beijing perfect Biotechnology, Beijing, China	Human H5N1 (haemagglutinin)	1:200	4°C	Overnight
Nucleoprotein	VivoStat-Inc, Portland, ME, USA	Influenza A virus (nucleoprotein)	1:20	4°C	Overnight
Neurofilament	Zymed Laboratories	Neuron	1:100	4°C	Overnight
Neuron-specific enolase	Zymed Laboratories	Neuron	1:100	4°C	Overnight
Gilal fibrillary acid protein	DakoCytomation	Glial cell	1:100	4°C	Overnight

Webtable: Sources, dilutions, and incubation times of primary monoclonal antibodies used in immunohistochemistry