

Supplemental Figures for:  
Postoperative Radiotherapy in Pathological T2-3N0M0 Thoracic Esophageal Squamous Cell Carcinoma: Interim Report of a  
Prospective, Phase III, Randomized Controlled Study  
Wei Deng et al.

Figure S1. Schoenfeld residuals plotted over time for each of the covariates in Table 1.






Figure S2A. Disease-free survival curves for surgery alone and postoperative radiotherapy groups with different T categories



Number at risk

S alone, T2	20	16	13	10	9	4
S + RT, T2	14	13	9	7	5	0
S alone, T3	57	44	30	23	15	5
S + RT, T3	66	57	45	31	20	9


Figure S2B. Overall survival curves for surgery alone and postoperative radiotherapy groups with different T categories



Number at risk

S alone, T2	20	18	15	13	9	4
S + RT, T2	14	13	10	8	5	0
S alone, T3	57	56	46	33	19	5
S + RT, T3	66	64	53	37	22	9


Figure S3A. Disease-free survival curves for surgery alone and postoperative radiotherapy groups with different tumor lengths



Number at risk

S alone, <5cm	27	21	13	9	8	4
S + RT, <5cm	32	29	24	19	12	5
S alone, >=5cm	50	39	30	24	16	5
S + RT, >=5cm	48	41	30	19	13	4

Figure S3B. Overall survival curves for surgery alone and postoperative radiotherapy groups with different tumor lengths



Number at risk

S alone, <5cm	27	26	22	16	11	4
S + RT, <5cm	32	30	25	19	12	5
S alone, >=5cm	50	48	39	30	17	5
S + RT, >=5cm	48	47	38	26	15	4