

Table SI. Primer sequences employed in reverse transcription-quantitative PCR.

Primer	Primer sequence (5'-3')	Supplier
GAPDH forward	GGAAGGTGAAGGTCGGAGTCA	Invitrogen ^a
GAPDH reverse	GTCATTGATGGCAACAATATCCACT	Invitrogen
B-Raf forward	AGAAAGCACTGATGATGAGAGG	Invitrogen
B-Raf reverse	GGAAATATCAGTGTCCAACCA	Invitrogen
E-cadherin forward	CCAGACCCACTCAGATGTCAAG	Takara Bio, Inc.
E-cadherin reverse	GGGCAGGTATGGAGAGGAAGA	Takara Bio, Inc.
N-cadherin forward	AACCTGGCGAGGACATCA	Takara Bio, Inc.
N-cadherin reverse	TCAAGGTCAAGACGTGCCAGA	Takara Bio, Inc.
Integrin α 2 forward	ACTTTGTTGCTGGTGCTCCT	Takara Bio, Inc.
Integrin α 2 reverse	CAAGAGCACGTCTGTAATGG	Takara Bio, Inc.
Integrin β 1 forward	GACGCCGCGCGAAAAGATG	Takara Bio, Inc.
Integrin β 1 reverse	GCACCACCCACAATTGGCCC	Takara Bio, Inc.
MMP2 forward	CTCATCGCAGATGCCTGGAA	Takara Bio, Inc.
MMP2 reverse	TTCAGGTAATAGGCACCCTGAAGA	Takara Bio, Inc.

^aInvitrogen, Thermo Fisher Scientific, Inc. MMP2, matrix metalloprotease 2.