## S3 Table. PTLD patients' characteristics.

Summary	Gender	Number of subjects	Age (years)		
			Mean ± SD	Median [Range]	
	F	7	2.29 ± 0.95	2.42 [1.21 – 3.51]	
	М	6	4.41 ± 4.31	3.05 [0.75 – 12.6]	
Total	F:M ratio <b>1.17</b>	13	3.27 ± 3.07	2.42 [0.75 – 12.6]	

## Details

No.	Type of Transplant	Type of PTLD	Acute Time-Point*	Recovery Time-Point*	Analysis
1	Liver	IM-like PTLD	0m	48m	a, b
2	Liver	IM-like PTLD	0.5m	36m	a, b
3	Liver	PTLD, plasmacytic hyperplasia	0m	36m	a, b
4	Liver	Monomorphic PTLD	0.25m	48m	a, b
5	Liver	IM-like PTLD	0.25m	45m	a, b
6	Liver	IM-like PTLD	0m	42m	a, b
7	Liver	Monomorphic PTLD (Burkitt lymphoma)	0m	45m	a, b
8	Liver and stem cells	Atypical PTLD; lymphoid infiltrate with a majority of EBER+ CD3+ CD8+ cells	0m	45m	a, b
9	Liver	IM-like PTLD	0.25m	48m	а
10	Liver	IM-like PTLD	0m	36m	а
11	Liver	IM-like PTLD	1m	36m	а
12	Liver	IM-like PTLD	3m	42m	а
13	Liver	IM-like PTLD	2m	42m	а

<sup>\*</sup> months post diagnosis a sCD30 serum concentration b PRR4 and cytokine serum concentration