

BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email info.bmjopen@bmj.com

BMJ Open

Androgen Deprivation Therapy and Iron-Deficiency Anemia among Patients with Prostate Cancer

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2019-034202
Article Type:	Original research
Date Submitted by the Author:	10-Sep-2019
Complete List of Authors:	Wu, Fang-Jen ; West Garden Hospital Li, I-Hsun; Tri-Service General Hospital Chien, Wu-Chien ; National Defense Medical Center Shih, Jui-Hu; Tri-Service General Hospital, Department of Pharmacy Practice Chuang, Chin-Min ; China Medical University Hospital Cheng, Yih-Dih ; China Medical University Hospital, Department of Pharmacy Kao, Li-Ting; Tri-Service General Hospital, Department of Pharmacy Practice
Keywords:	androgen deprivation therapy, iron-deficiency anemia, prostate cancer, anemia, drug safety

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

1
2
3
4 **Androgen Deprivation Therapy and Iron-Deficiency Anemia among**
5
6 **Patients with Prostate Cancer**
7
8
9

10
11 Fang-Jen Wu ¹ I-Hsun Li ^{2,3} Wu-Chien Chien ^{4,5,6} Jui-Hu Shih ^{2,3}
12
13 Chin-Min Chuang ⁷ Yih-Dih Cheng ^{8*} Li-Ting Kao ^{2,3,4*}
14
15

16
17 ¹ Department of Pharmacy, West Garden Hospital, Taipei, Taiwan

18
19 ² Department of Pharmacy Practice, Tri-Service General Hospital, Taipei, Taiwan

20
21 ³ School of Pharmacy, National Defense Medical Center, Taipei, Taiwan

22
23 ⁴ Graduate Institute of Life Sciences, National Defense Medical Center, Taipei, Taiwan

24
25 ⁵ Department of Medical Research, Tri-Service General Hospital, National Defense
26 Medical Center, Taipei, Taiwan.

27
28 ⁶ School of Public Health, National Defense Medical Center, Taipei, Taiwan.

29
30 ⁷ Emergency Department, China Medical University Hospital

31
32 ⁸ Department of Pharmacy, China Medical University Hospital, Taichung City, Taiwan
33
34

35
36 **Corresponding author:** Li-Ting Kao, Department of Pharmacy Practice, Tri-Service
37 General Hospital, No.325, Sec.2, Chenggong Rd., Neihu District, Taipei City 11490,
38 Taiwan. Tel: 886-2-8792-3311 ext. 17629, Fax: 886-2-8792-3171, E-mail:
39 kaoliting@gmail.com
40
41
42

43
44
45 * Li-Ting Kao and Yih-Dih Cheng have equal contributions to this study
46
47

48 **Running title:** Androgen Deprivation Therapy and Iron-Deficiency Anemia

49 **Disclosure:** No conflict of financial interest to declare.

50
51 No funding source.

52
53 Word count for text: 2855, number of tables/figures: 2/1, number of references: 21
54
55
56
57
58
59
60

Abstract

Objectives: Administration of androgen deprivation therapy (ADT) for metastatic prostate cancer might be associated with some adverse effects, such as anemia, but very few studies have been performed in East Asian populations. This study aimed to investigate the association between ADT and iron-deficiency anemia (IDA) among prostate cancer patients in a population-based nationwide cohort.

Design: Cohort study.

Setting: Taiwan.

Participants: Data for the cohort study was retrieved from the Taiwan National Health Insurance Research Dataset. Propensity score matching was used to select 8,474 prostate cancer patients who received ADT as study group and 4,237 prostate cancer patients who did not receive ADT as control group.

Primary and secondary outcome measures: This study individually tracked each patient for a 3-year study period and identify those who were subsequently diagnosed with IDA following the index date.

Results: This study revealed a 3.68 (95% CI 3.29-4.13) and 2.26 (95% CI 1.85-2.77) incidence rate of IDA per 100,000 person-years for prostate cancer patients receiving and not receiving ADT, respectively. Furthermore, proportional Cox regression showed

1
2
3
4
5 an hazard ratio (HR) for IDA in prostate cancer patients receiving ADT of 1.61 (95%
6
7
8 CI 1.28–2.03) after adjusting for patients' age, monthly income, geographic location,
9
10
11 residential urbanization level, hyperlipidemia, diabetes, hypertension, coronary heart
12
13
14 disease, and inflammatory bowel disease.

15
16
17
18 **Conclusion:** ADT use was found to be associated with a higher risk of IDA compared
19
20
21 with androgen deprivation therapy non-use among patients with prostate cancer.
22
23
24
25
26
27
28

29 **Key words:** androgen deprivation therapy, iron-deficiency anemia, prostate cancer,
30
31
32 anemia, drug safety.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Strengths and limitations of this study

1. Data for this cohort study was retrieved from a large, Asian population-based dataset in Taiwan.
2. This study used the propensity score-matched strategy to eliminate the potential bias such as patients' demographic characteristics and comorbidities between the study cohort and the comparison cohort.
3. The sample size and statistical power were sufficient in this study.
4. Selection bias could be eliminated due to the use of population-based dataset.
5. Information on personal history and lifestyle was unavailable when data was retrieved.

Funding source: This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors

Competing interests: None declared.

INTRODUCTION

Prostate cancer (PC) is the leading cause of male cancer and accounts for 293,000 deaths globally.^{1,2} In 2018, an estimated 164,690 new male diagnoses and 29,430 deaths from the disease were reported in the United States³. Androgen deprivation therapy (ADT), also known as hormone therapy, has been a fundamental component of metastatic PC management for more than half a century.⁴ Although androgen deprivation treatment may improve long-term survival for many patients,^{5,6} however, decreased serum levels of endogenous androgen in men with PC which have been linked to potential adverse effects, including decreased muscle mass, increased insulin resistance and fall risk.⁷⁻¹⁰

Iron-deficiency anemia (IDA) is a common type of anemia and approximately 5% females and 2% males would experience this clinical symptoms in the United States¹¹. More and more studies have recognized as that anemia may have a potential detrimental impact on quality of life and survival among PC patients.¹²⁻¹⁵ Previous study (InCHIANTI study) further suggested that androgen deficiency may be a factor that contribute to the development of anemia in men with PC¹⁶. Some studies had proposed that androgens may stimulate erythropoiesis through directly effect on erythroid progenitor cells and indirectly inhibit on hepcidin.¹⁷⁻¹⁹ Therefore, it is plausible that ADT use may be associated with the incidence of IDA.

1
2
3
4
5 Accordingly, the association between ADT and anemia has been reported in several
6
7
8 Western studies with Caucasian population.²⁰⁻²² To date, very few studies have been
9
10
11 conducted in East Asian populations, even though the hereditary of prostate cancer was
12
13
14 substantially vary by ethnicity and geography.²³ To fill the gap, this study proposed to
15
16
17 examine whether ADT is associated with a subsequent risk of anemia in PC patients by
18
19
20 employing the propensity score-matched strategy with an Asian population-based
21
22
23 dataset in Taiwan.
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

METHODS

Database

Data for the retrospective cohort study was retrieved from the Taiwan National Health Insurance Research Dataset (NHIRD). The NHIRD is derived from the Taiwan Bureau of National Health Insurance. It consists of inpatient and ambulatory medical benefit claims from approximately 22 million enrollees, representing over 99% of Taiwan's population. Therefore, this database provides scientists in Taiwan with an exclusive opportunity to analyze and keep track of medical service use by enrollees since the beginning of the Taiwan National Health Insurance program in 1995. This study was exempt from full revision by the Tri-Service General Hospital Institutional Review Board (TSGHIRB No. 2-105-05-082), since the dataset used was blinded to the public for research purposes.

Study Sample

This cohort study included a study and a comparison group. For the study group, 28,367 patients who received a first-time diagnosis of PC (ICD-9-CM code 185) between January 1, 2001 to December 31, 2010, were selected from the Registry of Catastrophic Illness Patient Database in NHIRD. A total of 454 patients aged less than 40 years were excluded. The date of ADT administration (including gonadotropin-releasing hormone (GnRH) agonists, antiandrogens, and estrogens) was assigned as the

1
2
3
4
5 index date for PC patients who subsequently received ADT treatment. Furthermore, the
6
7
8 date of the first outpatient visit in which patients received the diagnosis of PC was set
9
10
11 as the index date for PC patients who did not receive ADT. A total of 681 patients who
12
13
14 had never received a diagnosis of IDA (ICD-9-CM codes 280, 280.0, 280.1, 280.8, and
15
16
17 280.9) prior to their index date were further excluded, as well as 2,730 patients who had
18
19
20 received an orchiectomy. As a result, 24,502 PC patients remained in the study, 20,265
21
22
23 of which received ADT and 4,237 did not receive ADT during the 3-year study period.
24
25

26 Propensity score matching was used to identify 8,474 PC patients who received
27
28
29 ADT (two for every PC patient who did not receive ADT) from the remaining ADT
30
31
32 group. Matching variables included age, monthly income, geographic location,
33
34
35 urbanization level of patients' residence (that were divided into 5 levels, with 1 being
36
37
38 the most urbanized and 5 being the least urbanized),²⁴ hyperlipidemia, diabetes,
39
40
41 hypertension, coronary heart disease, and inflammatory bowel disease. Ultimately,
42
43
44 12,711 patients were enrolled in this study, including 8,474 who received ADT and
45
46
47 4,237 who did not receive ADT. Each patient was subsequently followed individually
48
49
50 for a 3-year study period to identify those who received an IDA diagnosis following
51
52
53 index date.
54
55
56
57
58
59
60

Statistical Analysis

All statistical analyses were performed with the SAS system (SAS System for Windows, vers. 9.4 SAS Institute). Chi-squared tests were used to investigate differences in sociodemographic characteristics and medical comorbidities between PC patients receiving and not receiving ADT. Additionally, Cox proportional hazard regressions were conducted to examine the relationship between ADT and IDA risk during the 3-year study period. A total of 2,716 patients who died during the 3-year study period (2,020 ADT use and 796 non - ADT users) were censored. The two-sided p value=0.05 was used to determine statistical significance.

Patient and Public Involvement

Patients or public were not involved in the development of the research question, outcome measures, study design, and recruitment/ conducting of the present study.

RESULTS

This study included 8,474 PC patients receiving ADT and 4,237 propensity score matching PC patients who did not receive ADT. Patients' sociodemographic characteristics and medical comorbidities were shown in Table 1. No statistically significant differences were found in age, monthly income, geographic location, urbanization level of patients' residence, hyperlipidemia, diabetes, hypertension, coronary heart disease, and inflammatory bowel disease between PC patients receiving and not receiving ADT.

IDA incidence during the 3-year study period is presented in Table 2. IDA incidence rate per 100,000 person-years for the 12,711 PC patients was 3.20 (95% CI 2.89-3.53). IDA incidence rate per 100,000 person-years for PC patients receiving and not receiving ADT was 3.68 (95% CI 3.29-4.13) and 2.26 (95% CI 1.85-2.77), respectively. Log-rank test revealed that PC patients receiving ADT had a significantly lower 3-year IDA-free survival rate than patients not receiving ADT ($p < 0.001$). Figure 1 shows the 3-year IDA-free survival curves for the study and comparison groups.

Table 2 further shows that hazard ratio (HR) for IDA in PC patients receiving ADT compared with those not receiving ADT was 1.62 (95% CI 1.29-2.05). After

1
2
3
4
5 adjusting for patients' age, monthly income, geographic location, residential
6
7
8 urbanization level, hyperlipidemia, diabetes, hypertension, coronary heart disease, and
9
10
11 inflammatory bowel disease, HR for PC patients receiving ADT compared with
12
13
14 nonusers was 1.61 (95% CI 1.28-2.03).
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

DISCUSSION

As far as the authors are aware, this is the first study that proposed to investigate the relationship between men receiving ADT for PC and 3-year IDA risk in East Asian populations. This study only included PC patients for relevant analyses in order to avoid the potential effects due to cancer. To mitigate selection bias, propensity score-matched strategy (by age, monthly income, geographical location, residential urbanization level, and medical comorbidities) was used to selected comparisons with PC patients not receiving ADT. In this population-based retrospective cohort study of PC patients, a significant increase in IDA risk was seen with men receiving ADT. The study reported an adjusted HR for IDA among PC patients receiving ADT of 1.61 (95% CI 1.28-2.03).

The findings in our study were consistent with some prior western studies. For instance, among 110 PC patients receiving ADT, one cohort study reported an average hemoglobin drop from 14.8 g/dL at baseline to 12.9 g/dL at evaluation²⁰. One Canadian clinical study including 250 PC patients found that the average decline in hemoglobin level of ADT users was a mean of 8.9 g/L²¹. In the United States, Strum et al. demonstrated the low hemoglobin levels in 90% of PC patients receiving ADT²⁵. Another study have reported PC patients receiving adjuvant radiotherapy plus ADT, with a decrease in hemoglobin level of 14.8 g/dL from the baseline to a minimum of 10.5 g/dL²⁶. In addition, one cohort study using the United Kingdom Clinical Practice

1
2
3
4
5 Research Database linked to the Hospital Episode Statistics repository found that
6
7
8 prostate patients with ADT had almost three-fold greater risk of anemia than nonusers
9
10 (HR 2.90, 95% CI: 2.67-3.16)²². Meanwhile, the clinical studies were designed with
11
12
13 small sample sizes and short duration which were mostly performed with Western
14
15
16 population²².
17

18
19 Our study indicated the association between ADT and 3-year IDA risk among PC
20
21
22 patients; however, anemia that caused by ADT has remained elusive. To date, increasing
23
24
25 biological evidences suggested a potential connection between men receiving ADT
26
27
28 and subsequent IDA risk.²⁷ A possible explanation for the underlying mechanism may
29
30
31 be that androgen significantly increases the bone marrow erythroid precursors and
32
33
34 enhances the differentiation of bone marrow erythroid stem cells to erythrocytes.^{19 28}
35
36
37 Furthermore, testosterone may have an influence on the erythropoietin secretion via
38
39
40 peritubular fibroblast-like cells level in kidney.²⁹ In animal experiments, testosterone
41
42
43 was able to induce erythropoiesis-stimulating factor by a bioassay using a polycythemic
44
45
46 mouse model.³⁰ Guo et al. found that the sensitivity of erythropoietin was stimulated
47
48
49 and the stress erythropoiesis could be increased, as result of testosterone upregulated the
50
51
52 expression of GATA binding protein 1 and GATA-dependent genes in the study of aging
53
54
55 male mice³¹. In addition, Bachman et al. indicated that men treated with testosterone
56
57
58 was related to serum hepcidin suppression in a randomized double-blind clinical study.¹⁷
59
60

1
2
3
4
5 Testosterone directly interact with bone morphogenetic protein-Smad signaling pathway
6
7
8 in hepatocytes and lead to the downregulation of hepcidin transcription.³² Furthermore,
9
10 this effects of androgen on hepatic hepcidin and renal erythropoietin gene expression
11
12
13 could increase systemic iron transport and erythropoietin levels.³²
14
15

16
17 The strength of our study was the identification of the association between ADT in
18
19 PC patients and subsequent IDA risk using a large population-based database. The
20
21 Taiwan NHIRD provides high accessibility for medical services and an adequate
22
23 statistical power to detect IDA risk between study and comparison groups. Moreover,
24
25
26 PC patients' sociodemographic characteristics and medical comorbidities acknowledged
27
28
29 as risk factors were also included. Furthermore, we also conducted a propensity score-
30
31
32 matched strategy to eliminate what potential bias such as patients' demographic
33
34
35 characteristics and comorbidities between the study cohort and the comparison cohort.
36
37
38
39

40
41 Several limitations of our retrospective study should be addressed. The study
42
43
44 lacked information on potentially important patient characteristics, such as body mass
45
46
47 index, smoking, nutrition, or nonprescription medication use. Secondly, it is plausible
48
49
50 that the database did not include all PC patients with IDA in Taiwan, since some of these
51
52
53 patients may seek alternative medicines not recorded by the NHI program. Thirdly,
54
55
56 information on personal history, including family history of anemia was unavailable
57
58
59 when data was retrieved.
60

CONCLUSIONS

In conclusion, after adjusting for patient sociodemographic characteristics and comorbid medical disorders, the present study detected an increased IDA risk among PC patients who received ADT during the considered 3-year follow-up. Medical professionals were suggested to be aware of the adverse effect of anemia by using ADT. Clinicians and pharmacists need to consider their PC patients with ADT for the possible IDA risk and determine the efficacy of prevention and treatment modalities for anemia.

Contributors

Conceptualization: Fang-Jen Wu & Yih-Dih Cheng & Li-Ting Kao; Methodology: Yih-Dih Cheng & Li-Ting Kao & Wu-Chien Chien; Formal analysis and investigation: Fang-Jen Wu & Jui-Hu Shih & Li-Ting Kao; Writing - original draft preparation: Fang-Jen Wu & I-Hsun Li & Chin-Min Chuang & Li-Ting Kao

Ethics approval

This study was exempt from full revision by the Tri-Service General Hospital Institutional Review Board (TSGHIRB No. 2-105-05-082), since the dataset used was blinded to the public for research purposes.

Data sharing statement

No additional data available.

REFERENCES

1. Fitzmaurice C, Dicker D, Pain A, et al. The Global Burden of Cancer 2013. *JAMA oncology* 2015;1(4):505-27. doi: 10.1001/jamaoncol.2015.0735
2. Attard G, Parker C, Eeles RA, et al. Prostate cancer. *Lancet* 2016;387(10013):70-82. doi: 10.1016/s0140-6736(14)61947-4
3. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2018. *CA Cancer J Clin* 2018;68(1):7-30. doi: 10.3322/caac.21442
4. Shahinian VB, Kuo YF, Gilbert SM. Reimbursement policy and androgen-deprivation therapy for prostate cancer. *N Engl J Med* 2010;363(19):1822-32. doi: 10.1056/NEJMsa0910784
5. Bryant AK, D'Amico AV, Nguyen PL, et al. Three-month posttreatment prostate-specific antigen level as a biomarker of treatment response in patients with intermediate-risk or high-risk prostate cancer treated with androgen deprivation therapy and radiotherapy. *Cancer* 2018;124(14):2939-47. doi: 10.1002/cncr.31400
6. Harris WP, Mostaghel EA, Nelson PS, et al. Androgen deprivation therapy: progress in understanding mechanisms of resistance and optimizing androgen depletion. *Nat Clin Pract Urol* 2009;6(2):76-85. doi: 10.1038/ncpuro1296
7. Basaria S, Muller DC, Carducci MA, et al. Hyperglycemia and insulin resistance in

- 1
2
3
4
5 men with prostate carcinoma who receive androgen-deprivation therapy. *Cancer*
6
7
8 2006;106(3):581-8. doi: 10.1002/cncr.21642
9
- 10
11 8. Grossmann M, Hamilton EJ, Gilfillan C, et al. Bone and metabolic health in patients
12
13 with non-metastatic prostate cancer who are receiving androgen deprivation
14
15 therapy. *Med J Aust* 2011;194(6):301-6.
16
17
- 18
19 9. Sharifi N, Gulley JL, Dahut WL. Androgen deprivation therapy for prostate cancer.
20
21
22
23 *Jama* 2005;294(2):238-44. doi: 10.1001/jama.294.2.238
24
25
- 26 10. Wu FJ, Sheu SY, Lin HC, et al. Increased Fall Risk in Patients Receiving Androgen
27
28 Deprivation Therapy for Prostate Cancer. *Urology* 2016;95:145-50. doi:
29
30 10.1016/j.urology.2016.05.058
31
32
33
- 34 11. Johnson-Wimbley TD, Graham DY. Diagnosis and management of iron deficiency
35
36 anemia in the 21st century. *Therap Adv Gastroenterol* 2011;4(3):177-84. doi:
37
38 10.1177/1756283X11398736
39
40
41
42
- 43 12. Caro JJ, Salas M, Ward A, et al. Anemia as an independent prognostic factor for
44
45 survival in patients with cancer: a systemic, quantitative review. *Cancer*
46
47 2001;91(12):2214-21.
48
49
50
51
- 52 13. Dicato M, Plawny L, Diederich M. Anemia in cancer. *Ann Oncol* 2010;21 Suppl
53
54 7:vii167-72. doi: 10.1093/annonc/mdq284
55
56
57
- 58 14. Dueregger A, Heidegger I, Ofer P, et al. The use of dietary supplements to alleviate
59
60

- 1
2
3
4 androgen deprivation therapy side effects during prostate cancer treatment.
5
6
7
8 *Nutrients* 2014;6(10):4491-519. doi: 10.3390/nu6104491
9
10
11 15. Lopez A, Cacoub P, Macdougall IC, et al. Iron deficiency anaemia. *Lancet*
12
13 2016;387(10021):907-16. doi: 10.1016/s0140-6736(15)60865-0
14
15
16 16. Ferrucci L, Maggio M, Bandinelli S, et al. Low testosterone levels and the risk of
17
18 anemia in older men and women. *Arch Intern Med* 2006;166(13):1380-8. doi:
19
20 10.1001/archinte.166.13.1380
21
22
23 17. Bachman E, Feng R, Travison T, et al. Testosterone suppresses hepcidin in men: a
24
25 potential mechanism for testosterone-induced erythrocytosis. *J Clin Endocrinol*
26
27 *Metab* 2010;95(10):4743-7. doi: 10.1210/jc.2010-0864
28
29
30
31
32
33 18. Latour C, Kautz L, Besson-Fournier C, et al. Testosterone perturbs systemic iron
34
35 balance through activation of epidermal growth factor receptor signaling in the
36
37 liver and repression of hepcidin. *Hepatology* 2014;59(2):683-94. doi:
38
39 10.1002/hep.26648
40
41
42
43
44
45
46 19. Moriyama Y, Fisher JW. Effects of testosterone and erythropoietin on erythroid
47
48 colony formation in human bone marrow cultures. *Blood* 1975;45(5):665-70.
49
50
51
52 20. D'Amico AV, Saegaert T, Chen MH, et al. Initial decline in hemoglobin during
53
54 neoadjuvant hormonal therapy predicts for early prostate specific antigen failure
55
56 following radiation and hormonal therapy for patients with intermediate and
57
58
59
60

- 1
2
3
4
5 high-risk prostate cancer. *Cancer* 2002;95(2):275-80. doi: 10.1002/cncr.10673
6
7
8 21. Timilshina N, Hussain S, Breunis H, et al. Predictors of hemoglobin decline in non-
9
10 metastatic prostate cancer patients on androgen deprivation therapy: a matched
11
12 cohort study. *Support Care Cancer* 2011;19(11):1815-21. doi: 10.1007/s00520-
13
14 010-1023-6
15
16
17
18
19 22. Hicks BM, Klil-Drori AJ, Yin H, et al. Androgen Deprivation Therapy and the Risk
20
21 of Anemia in Men with Prostate Cancer. *Epidemiology* 2017;28(5):712-18. doi:
22
23 10.1097/ede.0000000000000678
24
25
26
27
28 23. Rebbeck TR. Prostate Cancer Genetics: Variation by Race, Ethnicity, and
29
30 Geography. *Semin Radiat Oncol* 2017;27(1):3-10. doi:
31
32 10.1016/j.semradonc.2016.08.002
33
34
35
36
37 24. Liu CY, Hung YT, Chuang YL, et al. Incorporating development stratification of
38
39 Taiwan townships into sampling design of large scale health interview survey. *J*
40
41 *Health Manag* 2006;4(1):1-22.
42
43
44
45
46 25. Strum SB, McDermed JE, Scholz MC, et al. Anaemia associated with androgen
47
48 deprivation in patients with prostate cancer receiving combined hormone
49
50 blockade. *Br J Urol* 1997;79(6):933-41. doi: 10.1046/j.1464-410x.1997.00234.x
51
52
53
54
55 26. Choo R, Chander S, Danjoux C, et al. How are hemoglobin levels affected by
56
57 androgen deprivation in non-metastatic prostate cancer patients? *The Canadian*
58
59
60

- 1
2
3
4
5 *journal of urology* 2005;12(1):2547-52.
6
7
8 27. Shahani S, Braga-Basaria M, Maggio M, et al. Androgens and erythropoiesis: past
9
10 and present. *J Endocrinol Invest* 2009;32(8):704-16. doi: 10.1007/bf03345745
11
12
13 28. Dai D, Han S, Li L, et al. Anemia is associated with poor outcomes of metastatic
14
15 castration-resistant prostate cancer, a systematic review and meta-analysis.
16
17 *American journal of translational research* 2018;10(12):3877-86.
18
19
20 29. Pan X, Suzuki N, Hirano I, et al. Isolation and characterization of renal
21
22 erythropoietin-producing cells from genetically produced anemia mice. *PLoS*
23
24 *One* 2011;6(10):e25839. doi: 10.1371/journal.pone.0025839
25
26
27
28 30. Mirand EA, Gordon AS, Wenig J. Mechanism of testosterone action in
29
30 erythropoiesis. *Nature* 1965;206(981):270-2. doi: 10.1038/206270a0
31
32
33
34 31. Guo W, Li M, Bhasin S. Testosterone supplementation improves anemia in aging
35
36 male mice. *J Gerontol A Biol Sci Med Sci* 2014;69(5):505-13. doi:
37
38 10.1093/gerona/glt127
39
40
41
42
43
44
45
46 32. Guo W, Bachman E, Li M, et al. Testosterone administration inhibits hepcidin
47
48 transcription and is associated with increased iron incorporation into red blood
49
50 cells. *Aging cell* 2013;12(2):280-91. doi: 10.1111/accel.12052
51
52
53
54
55
56
57
58
59
60

1
2
3
4 **Figure Legends**
5
6

7 **Figure 1** Kaplan–Meier curve for iron-deficiency anemia (IDA) among prostate cancer
8
9
10 patients during a 3-year follow-up period, stratified by previous administration or non-
11
12
13 administration of androgen deprivation therapy (ADT)
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

Table 1 Demographic characteristics of prostate cancer patients, stratified by previous administration or non-administration of androgen deprivation therapy (ADT) ($n = 12,711$)

Characteristic	Patients not receiving ADT ($n = 4,237$)		Patients receiving ADT ($n = 8,474$)		<i>p</i> value
	Total no.	Column %	Total no.	Column %	
	Age (years), mean (SD)	70.99 (8.65)		71.03 (8.59)	
Urbanization level					0.9921
1 (most urbanized)	1,168	27.57	2,326	27.45	
2	990	23.37	1,985	23.42	
3	584	13.78	1,163	13.72	
4	537	12.67	1,056	12.46	
5 (least urbanized)	958	22.61	1,944	22.94	
Geographic region					0.9757
Northern	2,255	53.22	4,500	53.10	
Central	994	23.46	1,969	23.24	
Southern	948	22.37	1,923	22.69	
Eastern	40	0.94	82	0.97	
Monthly income					0.5705
NT\$0–15,840	2,840	67.03	5,741	67.75	
NT\$15,841–25,000	1,036	24.45	2,053	24.23	
≥NT\$25,001	361	8.52	680	8.02	
Hypertension	2,338	55.18	4,820	56.88	0.0686
Hyperlipidemia	894	21.10	1,705	20.12	0.1968
Diabetes	911	21.50	1,790	21.1	0.6237
Coronary heart disease	952	22.47	1,802	21.27	0.1205
Inflammatory bowel disease	37	0.87	70	0.83	0.7836

Notes: The average exchange rate in 2015 was US\$1.00≈New Taiwan (NT)\$30.

Table 2 Crude and adjusted hazard ratios (HRs) for anemia among prostate cancer patients during a 3-year follow-up period, stratified by previous administration or non-administration of androgen deprivation therapy (ADT)

Presence of iron-deficiency anemia	Total sample (n = 12,711)	Patients receiving ADT (n = 8,474)	Patients not receiving ADT (n = 4,237)
Three-year follow-up period			
Incidence rate per 100,000 person-years (95% CI)	3.20 (2.89–3.53)	3.68 (3.29–4.13)	2.26 (1.85–2.77)
Crude HR (95% CI)	--	1.62*** (1.29–2.05)	1.00
Adjusted ^a HR (95% CI)	--	1.61*** (1.28–2.03)	1.00

Notes: ^a Using a Cox proportional regression with cases censored if individuals died during the 3-year follow-up period; ^a Adjustments were made for patients’ geographic location, monthly income, urbanization level, age, hyperlipidemia, diabetes, hypertension, coronary heart disease, and inflammatory bowel disease.

* $p < 0.05$. ** $p < 0.01$. *** $p \leq 0.001$

CI, confidence interval




Figure 1 Kaplan–Meier curve for iron-deficiency anemia (IDA) among prostate cancer patients during a 3-year follow-up period, stratified by previous administration or non-administration of androgen deprivation therapy (ADT)

199x149mm (300 x 300 DPI)

STROBE 2007 (v4) Statement—Checklist of items that should be included in reports of *cohort studies*

Section/Topic	Item #	Recommendation	Reported on page #
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	2
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	2
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	5
Objectives	3	State specific objectives, including any prespecified hypotheses	6
Methods			
Study design	4	Present key elements of study design early in the paper	7
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	7-8
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up	7-8
		(b) For matched studies, give matching criteria and number of exposed and unexposed	7-8
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	7-8
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	7-8
Bias	9	Describe any efforts to address potential sources of bias	7-8
Study size	10	Explain how the study size was arrived at	7-8
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	7-8
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	9
		(b) Describe any methods used to examine subgroups and interactions	9
		(c) Explain how missing data were addressed	9
		(d) If applicable, explain how loss to follow-up was addressed	9
		(e) Describe any sensitivity analyses	9
Results			

Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	10-11
		(b) Give reasons for non-participation at each stage	10-11
		(c) Consider use of a flow diagram	10-11
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	10-11
		(b) Indicate number of participants with missing data for each variable of interest	10-11
		(c) Summarise follow-up time (eg, average and total amount)	10-11
Outcome data	15*	Report numbers of outcome events or summary measures over time	10-11
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	10-11
		(b) Report category boundaries when continuous variables were categorized	10-11
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	10-11
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	10-11
Discussion			
Key results	18	Summarise key results with reference to study objectives	13
Limitations			
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	15
Generalisability	21	Discuss the generalisability (external validity) of the study results	15
Other information			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	4

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.

BMJ Open

Androgen Deprivation Therapy and the Risk of Iron-Deficiency Anaemia among Patients with Prostate Cancer: A Population-based Cohort Study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2019-034202.R1
Article Type:	Original research
Date Submitted by the Author:	24-Dec-2019
Complete List of Authors:	Wu, Fang-Jen ; West Garden Hospital Li, I-Hsun; Tri-Service General Hospital Chien, Wu-Chien ; National Defense Medical Center Shih, Jui-Hu; Tri-Service General Hospital, Department of Pharmacy Practice Lin, Yi-Chun ; Research Center of Biostatistics, Taipei Medical University Chuang, Chin-Min ; China Medical University Hospital Cheng, Yih-Dih ; China Medical University Hospital, Department of Pharmacy Kao, Li-Ting; Tri-Service General Hospital, Department of Pharmacy Practice
Primary Subject Heading:	Urology
Secondary Subject Heading:	Epidemiology, Pharmacology and therapeutics
Keywords:	androgen deprivation therapy, iron-deficiency anemia, prostate cancer, anemia, drug safety

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

1
2
3
4 **Androgen Deprivation Therapy and the Risk of Iron-Deficiency Anaemia**
5
6 **among Patients with Prostate Cancer: A Population-based Cohort Study**
7

8 Fang-Jen Wu¹ I-Hsun Li^{2,3} Wu-Chien Chien^{4,5,6} Jui-Hu Shih^{2,3}
9 Yi-Chun Lin⁷ Chin-Min Chuang⁸ Yih-Dih Cheng^{9*} Li-Ting Kao^{2,3,4*}
10
11
12
13

14 ¹ Department of Pharmacy, West Garden Hospital, Taipei, Taiwan

15
16 ² Department of Pharmacy Practice, Tri-Service General Hospital, Taipei, Taiwan

17
18 ³ School of Pharmacy, National Defense Medical Center, Taipei, Taiwan

19
20 ⁴ Graduate Institute of Life Sciences, National Defense Medical Center, Taipei, Taiwan

21
22
23 ⁵ Department of Medical Research, Tri-Service General Hospital, National Defense
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Medical Center, Taipei, Taiwan.

⁶ School of Public Health, National Defense Medical Center, Taipei, Taiwan.

⁷ Research Center of Biostatistics, Taipei Medical University

⁸ Emergency Department, China Medical University Hospital

⁹ Department of Pharmacy, China Medical University Hospital, Taichung City, Taiwan

Corresponding author: Li-Ting Kao, Department of Pharmacy Practice, Tri-Service
General Hospital, No.325, Sec.2, Chenggong Rd., Neihu District, Taipei City 11490,
Taiwan. Tel: 886-2-8792-3311 ext. 17629, Fax: 886-2-8792-3171, E-mail:
kaoliting@gmail.com

* Li-Ting Kao and Yih-Dih Cheng have equal contributions to this study

Running title: Androgen Deprivation Therapy and Iron-Deficiency Anaemia

Disclosure: No conflict of financial interest to declare.

No funding source.

Word count for text: 2195, **number of tables/figures:** 3/1, **number of references:**

37

Abstract

Objectives: The administration of androgen deprivation therapy (ADT) to patients with metastatic prostate cancer might be associated with some adverse effects such as anaemia; however, few studies have been performed in East Asian populations. This study aimed to investigate the association between ADT and iron-deficiency anaemia (IDA) among patients with prostate cancer in a population-based nationwide cohort.

Design: Cohort study.

Setting: Taiwan.

Participants: Data for the cohort study were retrieved from the Taiwan National Health Insurance Research Database. Propensity score matching was used to select 8474 patients with prostate cancer who received ADT as the study group and 4237 patients with 4237 who did not receive ADT as the control group.

Primary and secondary outcome measures: This study individually tracked patients over a 3-year study period and identified those who were subsequently diagnosed with IDA following the index date.

Results: The incidence rates of IDA in the study and control groups were 1.35 (95% confidence interval [CI] = 1.20–1.51) and 0.83 per 100 person-years (95% CI = 0.67–1.01), respectively. Furthermore, proportional Cox regression revealed a hazard ratio

1
2
3
4
5 (HR) of 1.61 (95% CI = 1.28–2.03) for IDA in the study group after adjusting for
6
7
8 patients' age, monthly income, geographic location, residential urbanisation level and
9
10
11 incidence of hyperlipidemia, diabetes, hypertension, coronary heart disease,
12
13
14 inflammatory bowel disease, other cancers and gastrointestinal bleeding.
15
16

17
18 **Conclusion:** Compared with its non-use among patients with prostate cancer, ADT use
19
20 was associated with a higher risk of IDA.
21
22
23
24
25
26
27
28

29 **Key words:** androgen deprivation therapy, iron-deficiency anaemia, prostate cancer,
30
31 anaemia, drug safety
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Strengths and limitations of this study

1. Data for this cohort study were retrieved from a large Asian population-based dataset in Taiwan.
2. This study used a propensity score matching strategy to eliminate potential biases such as patients' demographics and comorbidities between the study and control cohorts.
3. The sample size and statistical power were sufficient in this study.
4. Selection bias may have been eliminated by the use of a population-based dataset.
5. Information on personal history and lifestyle was unavailable when data were retrieved.

Funding source: This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors

Competing interests: None declared.

INTRODUCTION

Prostate cancer (PCa) is the leading cause of cancer in males, and it accounted for 293,000 deaths globally in 2013.^{1 2} In 2018, an estimated 164,690 new diagnoses in males and 29,430 deaths from the disease were reported in the United States.³ Androgen deprivation therapy (ADT), also known as hormone therapy, has been a fundamental component of metastatic PCa management for more than half a century.⁴ This therapy can improve long-term survival for many patients.^{5 6} However, the decreased serum levels of endogenous androgen associated with ADT might result in some adverse effects, including decreased muscle mass, increased insulin resistance and fall risk, among patients with PCa.⁷⁻¹²

Iron-deficiency anaemia (IDA) is a common disease that affects approximately 5% of females and 2% of males in the United States.¹³ Increasing numbers of studies have recognised that anaemia may have a potentially detrimental impact on the quality of life and survival of patients with PCa.¹⁴⁻¹⁷ A previous study (InCHIANTI study) further suggested that androgen deficiency contributes to the development of anaemia in men with PCa.¹⁸ Some studies proposed that androgens may stimulate erythropoiesis through direct effects on erythroid progenitor cells and indirect inhibitory effects on hepcidin.¹⁹⁻²¹ Therefore, it is plausible that ADT may be associated with the incidence of IDA.

1
2
3
4
5 Accordingly, the association between ADT and anaemia has been reported in
6
7 several Western studies using Caucasian populations.²²⁻²⁴ However, to date, few studies
8
9 have been conducted in East Asian populations even though the hereditary risk of PCa
10
11 substantially varies by ethnicity and geography.^{25 26} In addition, the haemoglobin
12
13 concentration, which is a critical indicator of anaemia, is recognised to vary by
14
15 race/ethnicity, lifestyle, demographics and other variables.²⁷ Furthermore, a recent study
16
17 of 32 patients receiving ADT demonstrated that haematologic toxicities such as anaemia
18
19 were more frequent in Chinese patients with PCa than in their Western counterparts.²⁸
20
21 Consequently, this study examined whether ADT is associated with a subsequent risk
22
23 of anaemia in patients with PCa by employing a propensity score matching strategy
24
25 using an Asian population-based dataset in Taiwan.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

METHODS

Database

Data for the retrospective cohort study were retrieved from the Taiwan National Health Insurance Research Database (NHIRD). The NHIRD, which derived from the Taiwan Bureau of National Health Insurance, consists of inpatient and ambulatory medical benefit claims from approximately 22 million enrollees, representing more than 99% of Taiwan's population. Therefore, this database provides scientists in Taiwan with an exclusive opportunity to analyse and track medical service use by enrollees since the beginning of the Taiwan National Health Insurance programme in 1995. This study was exempt from full revision by the Tri-Service General Hospital Institutional Review Board (TSGHIRB No. 2-105-05-082) because the dataset used was blinded to the public for research purposes.

Study Sample

This cohort study included study and control groups. For the study group, 28,367 patients who received a first-time diagnosis of PCa (ICD-9-CM code 185) between January 1, 2001 and December 31, 2010 were selected from the Registry of Catastrophic Illness Patient Database in NHIRD. In total, 454 patients younger than 40 years were excluded because the prevalence of PCa is extremely low in this age strata. The date of ADT administration (including gonadotropin-releasing hormone agonists, anti-

1
2
3
4
5 androgens, ketoconazole and estrogens; ATC codes L02AE03, L02AE02, L02AE04,
6
7 L02BB03, G03HA01, L02BB01, J02AB02 and L02AA) was assigned as the index date
8
9
10 for patients with PCa who subsequently received ADT treatment. Furthermore, the date
11
12
13 of a randomly selected outpatient visit was assigned as the index date for the diagnosis
14
15
16 of PCa for patients who did not receive ADT. Meanwhile, 681 patients who had received
17
18
19 a diagnosis of IDA (ICD-9-CM codes 280, 280.0, 280.1, 280.8 and 280.9) and 2730
20
21
22 patients who had undergone orchiectomy prior to their index date were excluded. As a
23
24
25 result, 24,502 patients with PCa remained in the study, including 20,265 and 4237
26
27
28 patients with ADT use and non-use, respectively, during the 3-year study period.
29
30
31 Specifically, 49.50, 17.98, 15.74, and 16.78% of the ADT prescriptions were for
32
33
34 cyproterone, bicalutamide, flutamide and other types of medications, respectively. The
35
36
37 average time from PCa diagnosis was approximately 111.17 days in our study.

40
41 Propensity score matching was used to identify 8474 patients with PCa who
42
43
44 received ADT. This methodology could eliminate the differences attributable to
45
46
47 differences in patient demographics and medical history between the groups. We used
48
49
50 propensity score matching using the Mahalanobis metric (caliper of 0.25 standard
51
52
53 deviations of the propensity score). The matching variables included age, monthly
54
55
56 income, geographic location, residential urbanisation level (divided into five levels, with
57
58
59 1 being the most urbanised and 5 being the least urbanised),²⁹ hyperlipidaemia, diabetes,
60

1
2
3
4 hypertension, coronary heart disease and inflammatory bowel disease. We performed a
5
6
7
8 2:1 matching study, because increasing sample size of cases or controls could increase
9
10
11 the statistic power of the findings. Ultimately, 12,711 patients were enrolled in this
12
13
14 study, including 8474 who received ADT and 4237 who did not receive ADT. We
15
16
17 categorised the patients receiving ADT into two levels according to the median duration
18
19
20 of ADT use (median = 353 days). The power for this study was adequate (power =
21
22
23 0.987). Each patient was subsequently followed individually for a 3-year period to
24
25
26 identify those who received an IDA diagnosis following the index date.
27
28
29

30 **Statistical Analysis**

31
32 All statistical analyses were performed using the SAS System for Windows (ver.
33
34 9.4, SAS Institute). Chi-squared tests were used to investigate differences in
35
36 sociodemographic characteristics and medical comorbidities between the study and
37
38 control groups. Additionally, Cox proportional hazard regression analysis was
39
40
41 conducted to examine the relationship between ADT use and IDA risk during the 3-year
42
43
44 study period. Data for patients who died or who were lost to follow-up during the study
45
46
47 period were censored in the Cox regression. In total, 2716 patients died during the 3-
48
49
50 year study period (2020 ADT users and 796 non-users). A two-sided p value < 0.05
51
52
53 denoted statistical significance.
54
55
56
57
58
59
60

Patient and Public Involvement

Patients and the public were not involved in the development of the research question, outcome measures, study design and recruitment/conduct of the present study.

For peer review only

RESULTS

Data for patients' sociodemographic characteristics and medical comorbidities are shown in Table 1. No statistically significant differences were found regarding age, monthly income, geographic location, residential urbanisation level and the incidence of hyperlipidaemia, diabetes, hypertension, coronary heart disease and inflammatory bowel disease between the study and control groups.

The incidence of IDA during the 3-year study period is presented in Table 2. The IDA incidence rate per 100 person-years for the entire 12,711-patient cohort was 1.17 (95% confidence interval [CI] = 1.05–1.29). The incidence rates per 100 person-years for the study and control groups were 1.35 (95% CI = 1.20–1.51) and 0.83 (95% CI = 0.67–1.01), respectively. The log-rank test revealed that the study group had a significantly lower 3-year IDA-free survival rate than the control group ($p < 0.001$, Figure 1).

Table 2 further illustrates that the hazard ratio (HR) for IDA in the study group relative to the findings in the control group was 1.62 (95% CI = 1.29–2.05; $p \leq 0.001$). After adjusting for patients' age, monthly income, geographic location, residential urbanisation level and the incidence of hyperlipidemia, diabetes, hypertension, coronary heart disease, inflammatory bowel disease, other cancers and gastrointestinal

1
2
3
4
5 bleeding, the HR for the study group compared with the control group was 1.61 (95%
6
7
8 CI = 1.28–2.03; $p \leq 0.001$). This study further classified patients in the study group as
9
10 short-term and long-term ADT users based on the median duration of use. Data
11
12 presented in Table 3 reveal that both short-term and long-term and ADT use can
13
14 increase the risk of anaemia. Compared with the findings in the control group, the
15
16 adjusted HRs for long-term and short-term ADT use in the study group were 1.65 (95%
17
18
19 CI = 1.28–2.13) and 1.56 (95% CI = 1.19–2.04), respectively.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

DISCUSSION

To the best of our knowledge, this is the first study to investigate the relationship between ADT use in men with PCa and the 3-year IDA risk in an East Asian population. This study only included patients with PCa in the relevant analyses to avoid the potential effects attributable to cancer. To mitigate selection bias, a propensity score matching strategy (by age, monthly income, geographical location, residential urbanisation level and medical comorbidities) was used to facilitate comparisons with patients with PCa who did not receive ADT. In this population-based retrospective cohort study of patients with PCa, a significant increase in IDA risk was observed with men treated with ADT.

The findings were consistent with those of prior Western studies. For instance, in the United States, Strum et al. demonstrated that the haemoglobin levels of patients who received ADT for PCa declined from a mean of 149 g/L at baseline to 139, 132 and 131 g/L after 1, 2 and 3 months, respectively.³⁰ In Canada, Timilshina et al. performed an observational study of 250 patients with non-metastatic PCa and found that ADT was independently associated with a reduction of haemoglobin levels over 12 months.²³ Among 110 patients with PCa who received ADT, one cohort study reported a decline in haemoglobin levels from 14.8 g/dL at baseline to 12.9 g/dL at evaluation.²² Another study of 72 patients with non-metastatic PCa who received adjuvant radiotherapy plus ADT reported that the haemoglobin level had significant declined after 2 years of

1
2
3
4
5 androgen suppression.³¹ All of these studies investigated the association of ADT with
6
7
8 haemoglobin levels, which is vital given that haemoglobin levels are important for
9
10
11 identifying anaemia. However, these clinical studies included small sample sizes and
12
13
14 featured a short duration.^{22 23 30 31} Separately, Hicks et al. performed a cohort study using
15
16
17 the United Kingdom Clinical Practice Research Database linked to the Hospital Episode
18
19
20 Statistics repository. Their findings revealed that patients with non-metastatic PCa who
21
22
23 received ADT had a nearly 3-fold greater risk of anaemia than non-users (HR = 2.90,
24
25
26 95% CI = 2.67–3.16).²⁴ However, this study was performed in a Western population,
27
28
29 and generalisation of its findings to other ethnic groups is not possible.

30
31
32 Our study examined the association between ADT and the 3-year IDA risk among
33
34
35 patients with PCa; however, the risk of anaemia caused by ADT use has remained
36
37
38 unclear. To date, increasing biological evidence has suggested a potential connection
39
40
41 between the receipt of ADT and subsequent IDA risk.³² A possible explanation for the
42
43
44 underlying mechanism may be that androgen significantly increases the levels of bone
45
46
47 marrow erythroid precursors and enhances the differentiation of bone marrow erythroid
48
49
50 stem cells to erythrocytes.^{21 33} Furthermore, testosterone might influence erythropoietin
51
52
53 secretion via peritubular fibroblast-like cells in the kidneys.³⁴ In animal experiments,
54
55
56 testosterone induced erythropoiesis-stimulating factor production in a bioassay using a
57
58
59 mouse model of polycythaemia vera.³⁵ Guo et al. found that sensitivity to erythropoietin
60

1
2
3
4
5 was induced and stress erythropoiesis was increased by testosterone, which upregulated
6
7
8 the expression of GATA binding protein 1 and GATA-dependent genes, in a study of
9
10
11 ageing male mice.³⁶ In addition, Bachman et al. observed that treatment with
12
13
14 testosterone in men was related to serum hepcidin suppression in a randomised double-
15
16
17 blind clinical study.¹⁹ Testosterone directly interacts with the bone morphogenetic
18
19
20 protein-Smad signalling pathway in hepatocytes and induces the downregulation of
21
22
23 hepcidin transcription.³⁷ Furthermore, these effects of androgen on hepatic hepcidin and
24
25
26 renal erythropoietin gene expression could increase systemic iron transport and
27
28
29 erythropoietin levels.³⁷

30
31
32 The strengths of our study included the identification of the association between
33
34
35 ADT use in patients with PCa and subsequent IDA risk using a large population-based
36
37
38 database. The Taiwan NHIRD provides high accessibility to medical service data and
39
40
41 adequate to detect IDA risk between the study and control groups. Moreover, the
42
43
44 sociodemographic characteristics and medical comorbidities of patients with PCa were
45
46
47 acknowledged as risk factors. Furthermore, we also used a propensity score matching
48
49
50 strategy to eliminate potential biases such as patients' demographics and comorbidities
51
52
53 between the study and control cohorts.

54
55
56 Several limitations of our retrospective study should be addressed. The study
57
58
59 lacked information on potentially important patient characteristics, such as body mass
60

1
2
3
4
5 index, smoking, nutrition and non-prescription medication use. Second, it is plausible
6
7
8 that the database did not include all patients with PCa and IDA in Taiwan because some
9
10
11 patients might have sought alternative medicines not recorded by the NHI programme
12
13
14 and some patients with mild or normocytic anaemia may not have immediately sought
15
16
17 medical treatment. Therefore, these patients might not have been captured via diagnosis
18
19
20 codes. Third, information on the family history of anaemia, cancer stage and grade for
21
22
23 the presence of metastases, chemotherapy, radiotherapy and surgery was unavailable
24
25
26 this study. Finally, this study lacked information regarding blood variables.
27
28
29 Nevertheless, this study identified patients with PCa who used ADT as cases and
30
31
32 patients with PCa patients who did not use ADT as controls. In general, physicians in
33
34
35 Taiwan perform complete blood counts for patients with PCa to identify suitable
36
37
38 treatments. Consequently, we considered that these factors may not have affected the
39
40
41 findings in this study.

42 43 44 45 **CONCLUSIONS**

46
47
48 In conclusion, the present study detected an increased IDA risk during a 3-year
49
50
51 follow-up period among patients with PCa who received ADT. Medical professionals
52
53
54 are recommended to be aware of the risk of anaemia following ADT. Clinicians and
55
56
57 pharmacists need to consider the possible risk of IDA among patients with PCa who
58
59
60 received ADT and assess the efficacy of preventative and treatment modalities for

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

anaemia.

For peer review only

Contributors

Conceptualisation: Fang-Jen Wu, Yih-Dih Cheng and Li-Ting Kao; Methodology: Yih-Dih Cheng, Yi-Chun Lin, Li-Ting Kao and Wu-Chien Chien; Formal analysis and investigation: Fang-Jen Wu, Jui-Hu Shih and Li-Ting Kao; Writing - original draft preparation: Fang-Jen Wu, I-Hsun Li, Chin-Min Chuang and Li-Ting Kao

Ethics approval

This study was exempt from full revision by the Tri-Service General Hospital Institutional Review Board (TSGHIRB No. 2-105-05-082) because the dataset used was blinded to the public for research purposes.

Data sharing statement

No additional data available.

REFERENCES

1. Fitzmaurice C, Dicker D, Pain A, et al. The Global Burden of Cancer 2013. *JAMA oncology* 2015;1(4):505-27. doi: 10.1001/jamaoncol.2015.0735
2. Attard G, Parker C, Eeles RA, et al. Prostate cancer. *Lancet* 2016;387(10013):70-82. doi: 10.1016/S0140-6736(14)61947-4
3. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2018. *CA Cancer J Clin* 2018;68(1):7-30. doi: 10.3322/caac.21442
4. Shahinian VB, Kuo YF, Gilbert SM. Reimbursement policy and androgen-deprivation therapy for prostate cancer. *N Engl J Med* 2010;363(19):1822-32. doi: 10.1056/NEJMsa0910784
5. Bryant AK, D'Amico AV, Nguyen PL, et al. Three-month posttreatment prostate-specific antigen level as a biomarker of treatment response in patients with intermediate-risk or high-risk prostate cancer treated with androgen deprivation therapy and radiotherapy. *Cancer* 2018;124(14):2939-47. doi: 10.1002/cncr.31400
6. Harris WP, Mostaghel EA, Nelson PS, et al. Androgen deprivation therapy: progress in understanding mechanisms of resistance and optimizing androgen depletion. *Nat Clin Pract Urol* 2009;6(2):76-85. doi: 10.1038/ncpuro1296
7. Owen PJ, Daly RM, Livingston PM, et al. Lifestyle guidelines for managing adverse

- 1
2
3
4 effects on bone health and body composition in men treated with androgen
5
6
7 deprivation therapy for prostate cancer: an update. *Prostate Cancer Prostatic*
8
9
10 *Dis* 2017;20(2):137-45. doi: 10.1038/pcan.2016.69
11
12
13
14 8. Nguyen PL, Alibhai SM, Basaria S, et al. Adverse effects of androgen deprivation
15
16 therapy and strategies to mitigate them. *Eur Urol* 2015;67(5):825-36. doi:
17
18 10.1016/j.eururo.2014.07.010
19
20
21
22
23 9. Basaria S, Muller DC, Carducci MA, et al. Hyperglycemia and insulin resistance in
24
25 men with prostate carcinoma who receive androgen-deprivation therapy. *Cancer*
26
27 2006;106(3):581-8. doi: 10.1002/cncr.21642
28
29
30
31
32 10. Grossmann M, Hamilton EJ, Gilfillan C, et al. Bone and metabolic health in patients
33
34 with non-metastatic prostate cancer who are receiving androgen deprivation
35
36 therapy. *Med J Aust* 2011;194(6):301-6.
37
38
39
40
41 11. Sharifi N, Gulley JL, Dahut WL. Androgen deprivation therapy for prostate cancer.
42
43 *JAMA* 2005;294(2):238-44. doi: 10.1001/jama.294.2.238
44
45
46
47 12. Wu FJ, Sheu SY, Lin HC, et al. Increased Fall Risk in Patients Receiving Androgen
48
49 Deprivation Therapy for Prostate Cancer. *Urology* 2016;95:145-50. doi:
50
51 10.1016/j.urology.2016.05.058
52
53
54
55
56 13. Johnson-Wimbley TD, Graham DY. Diagnosis and management of iron deficiency
57
58 anemia in the 21st century. *Therap Adv Gastroenterol* 2011;4(3):177-84. doi:
59
60

1
2
3
4
5 10.1177/1756283X11398736
6
7

- 8 14. Caro JJ, Salas M, Ward A, et al. Anemia as an independent prognostic factor for
9
10 survival in patients with cancer: a systemic, quantitative review. *Cancer*
11
12 2001;91(12):2214-21.
13
14
15
16 15. Dicato M, Plawny L, Diederich M. Anemia in cancer. *Ann Oncol* 2010;21 Suppl
17
18 7:vii167-72. doi: 10.1093/annonc/mdq284
19
20
21
22 16. Dueregger A, Heidegger I, Ofer P, et al. The use of dietary supplements to alleviate
23
24 androgen deprivation therapy side effects during prostate cancer treatment.
25
26 *Nutrients* 2014;6(10):4491-519. doi: 10.3390/nu6104491
27
28
29
30
31 17. Lopez A, Cacoub P, Macdougall IC, et al. Iron deficiency anaemia. *Lancet*
32
33 2016;387(10021):907-16. doi: 10.1016/s0140-6736(15)60865-0
34
35
36
37 18. Ferrucci L, Maggio M, Bandinelli S, et al. Low testosterone levels and the risk of
38
39 anemia in older men and women. *Arch Intern Med* 2006;166(13):1380-8. doi:
40
41 10.1001/archinte.166.13.1380
42
43
44
45
46 19. Bachman E, Feng R, Travison T, et al. Testosterone suppresses hepcidin in men: a
47
48 potential mechanism for testosterone-induced erythrocytosis. *J Clin Endocrinol*
49
50 *Metab* 2010;95(10):4743-7. doi: 10.1210/jc.2010-0864
51
52
53
54
55 20. Latour C, Kautz L, Besson-Fournier C, et al. Testosterone perturbs systemic iron
56
57 balance through activation of epidermal growth factor receptor signaling in the
58
59
60

- 1
2
3
4
5 liver and repression of hepcidin. *Hepatology* 2014;59(2):683-94. doi:
6
7
8 10.1002/hep.26648
9
- 10
11 21. Moriyama Y, Fisher JW. Effects of testosterone and erythropoietin on erythroid
12
13 colony formation in human bone marrow cultures. *Blood* 1975;45(5):665-70.
14
15
- 16
17 22. D'Amico AV, Saegaert T, Chen MH, et al. Initial decline in hemoglobin during
18
19 neoadjuvant hormonal therapy predicts for early prostate specific antigen failure
20
21 following radiation and hormonal therapy for patients with intermediate and
22
23 high-risk prostate cancer. *Cancer* 2002;95(2):275-80. doi: 10.1002/cncr.10673
24
25
26
27
- 28
29 23. Timilshina N, Hussain S, Breunis H, et al. Predictors of hemoglobin decline in non-
30
31 metastatic prostate cancer patients on androgen deprivation therapy: a matched
32
33 cohort study. *Support Care Cancer* 2011;19(11):1815-21. doi: 10.1007/s00520-
34
35
36
37 010-1023-6
38
39
- 40
41 24. Hicks BM, Klil-Drori AJ, Yin H, et al. Androgen Deprivation Therapy and the Risk
42
43 of Anemia in Men with Prostate Cancer. *Epidemiology* 2017;28(5):712-18. doi:
44
45
46
47 10.1097/ede.0000000000000678
48
49
- 50
51 25. Rebbeck TR. Prostate Cancer Genetics: Variation by Race, Ethnicity, and
52
53 Geography. *Semin Radiat Oncol* 2017;27(1):3-10. doi:
54
55
56 10.1016/j.semradonc.2016.08.002
57
58
- 59
60 26. Taitt HE. Global Trends and Prostate Cancer: A Review of Incidence, Detection,

- 1
2
3
4
5 and Mortality as Influenced by Race, Ethnicity, and Geographic Location.
6
7
8 *American journal of men's health* 2018;12(6):1807-23. doi:
9
10 10.1177/1557988318798279
11
12
13
14 27. McLean E, Cogswell M, Egli I, et al. Worldwide prevalence of anaemia, WHO
15
16 Vitamin and Mineral Nutrition Information System, 1993-2005. *Public Health*
17
18 *Nutr* 2009;12(4):444-54. doi: 10.1017/s1368980008002401
19
20
21
22
23 28. Poon DMC, Chan T, Chan K, et al. Preliminary efficacy and tolerability of
24
25 chemohormonal therapy in metastatic hormone-naive prostate cancer: The first
26
27 real-life experience in Asia. *Asia Pac J Clin Oncol* 2018;14(4):347-52. doi:
28
29 10.1111/ajco.12874
30
31
32
33
34
35 29. Liu CY, Hung YT, Chuang YL, et al. Incorporating development stratification of
36
37 Taiwan townships into sampling design of large scale health interview survey. *J*
38
39 *Health Manag* 2006;4(1):1-22.
40
41
42
43
44 30. Strum SB, McDermed JE, Scholz MC, et al. Anaemia associated with androgen
45
46 deprivation in patients with prostate cancer receiving combined hormone
47
48 blockade. *Br J Urol* 1997;79(6):933-41. doi: 10.1046/j.1464-410x.1997.00234.x
49
50
51
52
53 31. Choo R, Chander S, Danjoux C, et al. How are hemoglobin levels affected by
54
55 androgen deprivation in non-metastatic prostate cancer patients? *The Canadian*
56
57 *journal of urology* 2005;12(1):2547-52.
58
59
60

- 1
2
3
4
5 32. Shahani S, Braga-Basaria M, Maggio M, et al. Androgens and erythropoiesis: past
6
7 and present. *J Endocrinol Invest* 2009;32(8):704-16. doi: 10.1007/bf03345745
8
9
- 10 33. Dai D, Han S, Li L, et al. Anemia is associated with poor outcomes of metastatic
11
12 castration-resistant prostate cancer, a systematic review and meta-analysis.
13
14 *American journal of translational research* 2018;10(12):3877-86.
15
16
17
18
19
- 20 34. Pan X, Suzuki N, Hirano I, et al. Isolation and characterization of renal
21
22 erythropoietin-producing cells from genetically produced anemia mice. *PLoS*
23
24 *One* 2011;6(10):e25839. doi: 10.1371/journal.pone.0025839
25
26
27
28
- 29 35. Mirand EA, Gordon AS, Wenig J. Mechanism of testosterone action in
30
31 erythropoiesis. *Nature* 1965;206(981):270-2. doi: 10.1038/206270a0
32
33
34
- 35 36. Guo W, Li M, Bhasin S. Testosterone supplementation improves anemia in aging
36
37 male mice. *J Gerontol A Biol Sci Med Sci* 2014;69(5):505-13. doi:
38
39 10.1093/gerona/glt127
40
41
42
43
- 44 37. Guo W, Bachman E, Li M, et al. Testosterone administration inhibits hepcidin
45
46 transcription and is associated with increased iron incorporation into red blood
47
48 cells. *Aging cell* 2013;12(2):280-91. doi: 10.1111/accel.12052
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 **Figure Legend**
5

6
7 **Figure 1** Kaplan–Meier curve of the risk of iron-deficiency anaemia (IDA) among
8
9
10 patients with prostate cancer during a 3-year follow-up period as stratified by the
11
12
13 previous use or non-use of androgen deprivation therapy (ADT).
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

Table 1 Demographic characteristics of patients with prostate cancer stratified by the previous use or non-use of androgen deprivation therapy (ADT) ($n = 12,711$)

Characteristic	Patients not receiving ADT ($n = 4237$)		Patients receiving ADT ($n = 8474$)		<i>p</i> value
	Total no.	Column %	Total no.	Column %	
Age (years), mean (SD)	70.99 (8.65)		71.03 (8.59)		0.7971
Urbanisation level					0.9921
1 (most urbanised)	1168	27.57	2326	27.45	
2	990	23.37	1985	23.42	
3	584	13.78	1163	13.72	
4	537	12.67	1056	12.46	
5 (least urbanised)	958	22.61	1944	22.94	
Geographic region					0.9757
Northern	2255	53.22	4500	53.10	
Central	994	23.46	1969	23.24	
Southern	948	22.37	1923	22.69	
Eastern	40	0.94	82	0.97	
Monthly income					0.5705
NT\$0–15,840	2840	67.03	5741	67.75	
NT\$15,841–25,000	1036	24.45	2053	24.23	
\geq NT\$25,001	361	8.52	680	8.02	
Hypertension	2338	55.18	4820	56.88	0.0686
Hyperlipidemia	894	21.10	1705	20.12	0.1968
Diabetes	911	21.50	1790	21.1	0.6237
Coronary heart disease	952	22.47	1802	21.27	0.1205
Inflammatory bowel disease	37	0.87	70	0.83	0.7836
Gastrointestinal bleeding	116	2.74	216	2.55	0.5292
Previous cancers	258	6.09	490	5.78	0.4884

Notes: The average exchange rate in 2015 was US\$1.00 \approx New Taiwan (NT)\$30.

Table 2 Crude and adjusted hazard ratios (HRs) for anaemia among patients with prostate cancer during a 3-year follow-up period stratified by the previous use or non-use of androgen deprivation therapy (ADT)

Presence of iron-deficiency anaemia	Total sample (<i>n</i> = 12,711)	Patients receiving ADT (<i>n</i> = 8474)	Patients not receiving ADT (<i>n</i> = 4237)
Three-year follow-up period			
Incidence rate per 100 person-years (95% CI)	1.17 (1.05–1.29)	1.35 (1.20–1.51)	0.83 (0.67–1.01)
Crude HR (95% CI)	--	1.62 (1.29–2.05)	1.00
Adjusted HR (95% CI) ^{a,b}	--	1.61 (1.28–2.03)	1.00
Adjusted HR (95% CI) ^{a,c}	--	1.61 (1.28–2.03)	1.00

Notes: ^a Using a Cox proportional regression with data censored if individuals died during the 3-year follow-up period; ^b Adjustments were made for patients' geographic location, monthly income, urbanisation level, age and incidence of hyperlipidaemia, diabetes, hypertension, coronary heart disease and inflammatory bowel disease. ^c Adjustments were made for patients' geographic location, monthly income, urbanisation level, age and incidence of hyperlipidaemia, diabetes, hypertension, coronary heart disease, inflammatory bowel disease, other cancers and gastrointestinal bleeding.

CI, confidence interval

Table 3 Crude and adjusted hazard ratios (HRs) for anaemia among patients with prostate cancer during a 3-year follow-up period stratified by the duration of androgen deprivation therapy (ADT)

Presence of iron-deficiency anaemia	Patients receiving long-term ADT (n =4239) ^d	Patients receiving short-term ADT (n =4235) ^d	Patients not receiving ADT (n = 4237)
Three-year follow-up period			
Incidence rate per 100 person-years (95% CI)	1.41 (1.20–1.64)	1.27 (1.06–1.51)	0.83 (0.67–1.01)
Crude HR (95% CI)	1.70 (1.32–2.19)	1.53 (1.17–2.00)	1.00
Adjusted HR (95% CI) ^{a,b}	1.65 (1.28–2.13)	1.56 (1.20–2.04)	1.00
Adjusted HR (95% CI) ^{a,c}	1.65 (1.28–2.13)	1.56 (1.19–2.04)	1.00

Notes: ^a Using a Cox proportional regression with data censored if individuals died during the 3-year follow-up period; ^b Adjustments were made for patients’ geographic location, monthly income, urbanisation level, age and incidence of hyperlipidemia, diabetes, hypertension, coronary heart disease and inflammatory bowel disease. ^c Adjustments were made for patients’ geographic location, monthly income, urbanisation level, age and incidence of hyperlipidaemia, diabetes, hypertension, coronary heart disease, inflammatory bowel disease, other cancers and gastrointestinal bleeding. ^d We categorised the patients receiving ADT into two levels according to the median duration of ADT use (353 days). CI, confidence interval




Figure 1 Kaplan–Meier curve of the risk of iron-deficiency anaemia (IDA) among patients with prostate cancer during a 3-year follow-up period as stratified by the previous use or non-use of androgen deprivation therapy (ADT).

199x150mm (500 x 500 DPI)

BMJ Open

Androgen Deprivation Therapy and the Risk of Iron-Deficiency Anaemia among Patients with Prostate Cancer: A Population-based Cohort Study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2019-034202.R2
Article Type:	Original research
Date Submitted by the Author:	18-Feb-2020
Complete List of Authors:	Wu, Fang-Jen ; West Garden Hospital Li, I-Hsun; Tri-Service General Hospital Chien, Wu-Chien ; National Defense Medical Center Shih, Jui-Hu; Tri-Service General Hospital, Department of Pharmacy Practice Lin, Yi-Chun ; Research Center of Biostatistics, Taipei Medical University Chuang, Chin-Min ; China Medical University Hospital Cheng, Yih-Dih ; China Medical University Hospital, Department of Pharmacy Kao, Li-Ting; Tri-Service General Hospital, Department of Pharmacy Practice
Primary Subject Heading:	Urology
Secondary Subject Heading:	Epidemiology, Pharmacology and therapeutics
Keywords:	androgen deprivation therapy, iron-deficiency anemia, prostate cancer, anemia, drug safety

SCHOLARONE™
Manuscripts



I, the Submitting Author has the right to grant and does grant on behalf of all authors of the Work (as defined in the below author licence), an exclusive licence and/or a non-exclusive licence for contributions from authors who are: i) UK Crown employees; ii) where BMJ has agreed a CC-BY licence shall apply, and/or iii) in accordance with the terms applicable for US Federal Government officers or employees acting as part of their official duties; on a worldwide, perpetual, irrevocable, royalty-free basis to BMJ Publishing Group Ltd ("BMJ") its licensees and where the relevant Journal is co-owned by BMJ to the co-owners of the Journal, to publish the Work in this journal and any other BMJ products and to exploit all rights, as set out in our [licence](#).

The Submitting Author accepts and understands that any supply made under these terms is made by BMJ to the Submitting Author unless you are acting as an employee on behalf of your employer or a postgraduate student of an affiliated institution which is paying any applicable article publishing charge ("APC") for Open Access articles. Where the Submitting Author wishes to make the Work available on an Open Access basis (and intends to pay the relevant APC), the terms of reuse of such Open Access shall be governed by a Creative Commons licence – details of these licences and which [Creative Commons](#) licence will apply to this Work are set out in our licence referred to above.

Other than as permitted in any relevant BMJ Author's Self Archiving Policies, I confirm this Work has not been accepted for publication elsewhere, is not being considered for publication elsewhere and does not duplicate material already published. I confirm all authors consent to publication of this Work and authorise the granting of this licence.

1
2
3
4 **Androgen Deprivation Therapy and the Risk of Iron-Deficiency Anaemia**
5
6 **among Patients with Prostate Cancer: A Population-based Cohort Study**
7

8 Fang-Jen Wu¹ I-Hsun Li^{2,3,4} Wu-Chien Chien^{5,6,7} Jui-Hu Shih^{2,3}

9 Yi-Chun Lin⁸ Chin-Min Chuang⁹ Yih-Dih Cheng^{10*} Li-Ting Kao^{2,3,5,7*}
10
11
12

13 ¹ Department of Pharmacy, West Garden Hospital, Taipei, Taiwan

14 ² Department of Pharmacy Practice, Tri-Service General Hospital, Taipei, Taiwan

15 ³ School of Pharmacy, National Defense Medical Center, Taipei, Taiwan

16 ⁴ Department of Pharmacology, National Defense Medical Center, Taipei, Taiwan

17 ⁵ Graduate Institute of Life Sciences, National Defense Medical Center, Taipei, Taiwan

18 ⁶ Department of Medical Research, Tri-Service General Hospital, National Defense
19 Medical Center, Taipei, Taiwan.

20 ⁷ School of Public Health, National Defense Medical Center, Taipei, Taiwan.

21 ⁸ Research Center of Biostatistics, Taipei Medical University, Taipei, Taiwan.

22 ⁹ Emergency Department, China Medical University Hospital, Taichung City, Taiwan

23 ¹⁰ Department of Pharmacy, China Medical University Hospital, Taichung City, Taiwan

24
25
26
27
28 **Corresponding author:** Li-Ting Kao, Department of Pharmacy Practice, Tri-Service
29 General Hospital, No.325, Sec.2, Chenggong Rd., Neihu District, Taipei City 11490,
30 Taiwan. Tel: 886-2-8792-3311 ext. 17629, Fax: 886-2-8792-3171, E-mail:
31 kaoliting@gmail.com
32
33
34

35
36
37 * Li-Ting Kao and Yih-Dih Cheng have equal contributions to this study
38
39

40 **Running title:** Androgen Deprivation Therapy and Iron-Deficiency Anaemia

41
42 **Disclosure:** No conflict of financial interest to declare.

43
44 **No funding source.**

45 **Word count for text:** 2213, **number of tables/figures:** 3/1, **number of references:** 37
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Objectives: The administration of androgen deprivation therapy (ADT) to patients with metastatic prostate cancer might be associated with some adverse effects such as anaemia; however, few studies have been performed in East Asian populations. This study aimed to investigate the association between ADT and iron-deficiency anaemia (IDA) among patients with prostate cancer in a population-based nationwide cohort.

Design: Cohort study.

Setting: Taiwan.

Participants: Data for the cohort study were retrieved from the Taiwan National Health Insurance Research Database. Propensity score matching was used to select 7,262 patients with prostate cancer who received ADT as the study group and 3,631 patients who did not receive ADT as the control group.

Primary and secondary outcome measures: This study individually tracked patients over a 3-year study period and identified those who were subsequently diagnosed with IDA following the index date.

Results: The incidence rates of IDA in the study and control groups were 1.66 (95% confidence interval [CI] = 1.45–1.86) and 1.01 per 100 person-years (95% CI = 0.78–1.25), respectively. Furthermore, proportional Cox regression revealed a hazard ratio

1
2
3
4
5 (HR) of 1.62 (95% CI = 1.24–2.12) for IDA in the study group after adjusting for
6
7
8 patients' age, monthly income, geographic location, residential urbanisation level and
9
10
11 incidence of hyperlipidemia, diabetes, hypertension, coronary heart disease,
12
13
14 inflammatory bowel disease, other cancers and gastrointestinal bleeding.
15
16

17
18 **Conclusion:** Compared with its non-use among patients with prostate cancer, ADT use
19
20 was associated with a higher risk of IDA.
21
22
23
24
25
26
27
28

29 **Key words:** androgen deprivation therapy, iron-deficiency anaemia, prostate cancer,
30
31 anaemia, drug safety
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Strengths and limitations of this study

1. Data for this cohort study were retrieved from a large Asian population-based dataset in Taiwan.
2. This study used a propensity score matching strategy to eliminate potential biases such as patients' demographics and comorbidities between the study and control cohorts.
3. The sample size and statistical power were sufficient in this study.
4. Selection bias may have been eliminated by the use of a population-based dataset.
5. Information on personal history and lifestyle was unavailable when data were retrieved.

Funding source: This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors

Competing interests: None declared.

INTRODUCTION

Prostate cancer (PCa) is the leading cause of cancer in males, and it accounted for 293,000 deaths globally in 2013.^{1 2} In 2018, an estimated 164,690 new diagnoses in males and 29,430 deaths from the disease were reported in the United States.³ Androgen deprivation therapy (ADT), also known as hormone therapy, has been a fundamental component of metastatic PCa management for more than half a century.⁴ This therapy can improve long-term survival for many patients.^{5 6} However, the decreased serum levels of endogenous androgen associated with ADT might result in some adverse effects, including decreased muscle mass, increased insulin resistance and fall risk, among patients with PCa.⁷⁻¹²

Iron-deficiency anaemia (IDA) is a common disease that affects approximately 5% of females and 2% of males in the United States.¹³ Increasing numbers of studies have recognised that anaemia may have a potentially detrimental impact on the quality of life and survival of patients with PCa.¹⁴⁻¹⁷ A previous study (InCHIANTI study) further suggested that androgen deficiency contributes to the development of anaemia in men with PCa.¹⁸ Some studies proposed that androgens may stimulate erythropoiesis through direct effects on erythroid progenitor cells and indirect inhibitory effects on hepcidin.¹⁹⁻
²¹ Therefore, it is plausible that ADT may be associated with the incidence of IDA.

1
2
3
4
5 Accordingly, the association between ADT and anaemia has been reported in
6
7 several Western studies using Caucasian populations.²²⁻²⁴ However, to date, few studies
8
9 have been conducted in East Asian populations even though the hereditary risk of PCa
10
11 substantially varies by ethnicity and geography.^{25 26} In addition, the haemoglobin
12
13 concentration, which is a critical indicator of anaemia, is recognised to vary by
14
15 race/ethnicity, lifestyle, demographics and other variables.²⁷ Furthermore, a recent study
16
17 of 32 patients receiving ADT demonstrated that haematologic toxicities such as anaemia
18
19 were more frequent in Chinese patients with PCa than in their Western counterparts.²⁸
20
21 Consequently, this study examined whether ADT is associated with a subsequent risk
22
23 of anaemia in patients with PCa by employing a propensity score matching strategy
24
25 using an Asian population-based dataset in Taiwan.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

METHODS

Database

Data for the retrospective cohort study were retrieved from the Taiwan National Health Insurance Research Database (NHIRD). The NHIRD, which derived from the Taiwan Bureau of National Health Insurance, consists of inpatient and ambulatory medical benefit claims from approximately 22 million enrollees, representing more than 99% of Taiwan's population. Therefore, this database provides scientists in Taiwan with an exclusive opportunity to analyse and track medical service use by enrollees since the beginning of the Taiwan National Health Insurance programme in 1995. This study was exempt from full revision by the Tri-Service General Hospital Institutional Review Board (TSGHIRB No. 2-105-05-082) because the dataset used was blinded to the public for research purposes.

Study Sample

This cohort study included study and control groups. For the study group, 28,367 patients who received a first-time diagnosis of PCa (ICD-9-CM code 185) between January 1, 2001 and December 31, 2010 were selected from the Registry of Catastrophic Illness Patient Database in NHIRD. In total, 454 patients younger than 40 years were excluded because the prevalence of PCa is extremely low in this age strata. The date of ADT administration (including gonadotropin-releasing hormone agonists, anti-

1
2
3
4
5 androgens, ketoconazole and estrogens; ATC codes L02AE03, L02AE02, L02AE04,
6
7 L02BB03, G03HA01, L02BB01, J02AB02 and L02AA) was assigned as the index date
8
9
10 for patients with PCa who subsequently received ADT treatment. Furthermore, the date
11
12
13 of a randomly selected outpatient visit was assigned as the index date for the diagnosis
14
15
16 of PCa for patients who did not receive ADT. Meanwhile, 1,605 patients who had
17
18
19 received a diagnosis of IDA (ICD-9-CM codes 280, 280.0, 280.1, 280.8 and 280.9) and
20
21
22 1,533 patients who had undergone orchiectomy prior to their index date were excluded.
23
24
25 As a result, 24,775 patients with PCa remained in the study, including 20,272 and 4,503
26
27
28 patients with ADT use and non-use, respectively, during the 3-year study period.
29
30
31 Specifically, 44.06, 26.06, 14.65, and 15.23% of the ADT prescriptions were for
32
33
34 cyproterone, bicalutamide, flutamide and other types of medications, respectively. The
35
36
37 average time from PCa diagnosis date to patient entry date for both non-users and users
38
39
40 were 758.4 days and 796.9 days, respectively.
41
42
43

44 Propensity score matching was used to identify 7,262 patients with PCa who
45
46
47 received ADT. This methodology could eliminate the differences attributable to
48
49
50 differences in patient demographics and medical history between the groups. We used
51
52
53 propensity score matching using the Mahalanobis metric (caliper of 0.25 standard
54
55
56 deviations of the propensity score). The matching variables included year of the entry
57
58
59 date, age, monthly income, geographic location, residential urbanisation level (divided
60

1
2
3
4 into five levels, with 1 being the most urbanised and 5 being the least urbanised),²⁹
5
6
7
8 hyperlipidaemia, diabetes, hypertension, coronary heart disease and inflammatory
9
10
11 bowel disease. We performed a 2:1 matching study, because increasing sample size of
12
13
14 cases or controls could increase the statistic power of the findings. Ultimately, 10,893
15
16
17 patients were enrolled in this study, including 7,262 who received ADT and 3,631 who
18
19
20 did not receive ADT. We categorised the patients receiving ADT into two levels
21
22
23 according to the median duration of ADT use (median = 144 days). Those patients who
24
25
26 received ADT <144 days were identified as short-term ADT users. Moreover, patients
27
28
29 receiving ADT \geq 144 days were defined as long-term ADT users. The power for this
30
31
32 study was adequate (power > 0.9). Each patient was subsequently followed individually
33
34
35 for a 3-year period to identify those who received an IDA diagnosis following the index
36
37
38 date.

41 **Statistical Analysis**

42
43
44 All statistical analyses were performed using the SAS System for Windows (ver.
45
46
47 9.4, SAS Institute). Chi-squared tests were used to investigate differences in
48
49
50 sociodemographic characteristics and medical comorbidities between the study and
51
52
53 control groups. Additionally, Cox proportional hazard regression analysis was
54
55
56 conducted to examine the relationship between ADT use and IDA risk during the 3-year
57
58
59 study period. Data for patients who died or who were lost to follow-up during the study
60

1
2
3
4
5 period were censored in the Cox regression. In total, 3,152 patients died during the 3-
6
7
8 year study period (2,153 ADT users and 999 non-users). A two-sided p value < 0.05
9
10 denoted statistical significance.
11
12
13

14 **Patient and Public Involvement**

15
16
17 Patients and the public were not involved in the development of the research
18
19 question, outcome measures, study design and recruitment/conduct of the present
20
21 study.
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

RESULTS

Data for patients' sociodemographic characteristics and medical comorbidities are shown in Table 1. No statistically significant differences were found regarding age, monthly income, geographic location, residential urbanisation level and the incidence of hyperlipidaemia, diabetes, hypertension, coronary heart disease and inflammatory bowel disease between the study and control groups.

The incidence of IDA during the 3-year study period is presented in Table 2. The IDA incidence rate per 100 person-years for the entire 10,893-patient cohort was 1.45 (95% confidence interval [CI] = 1.29–1.61). The incidence rates per 100 person-years for the study and control groups were 1.66 (95% CI = 1.45–1.86) and 1.01 (95% CI = 0.78–1.25), respectively. The log-rank test revealed that the study group had a significantly lower 3-year IDA-free survival rate than the control group ($p < 0.001$, Figure 1).

Table 2 further illustrates that the hazard ratio (HR) for IDA in the study group relative to the findings in the control group was 1.64 (95% CI = 1.25–2.14; $p \leq 0.001$). After adjusting for patients' age, monthly income, geographic location, residential urbanisation level and the incidence of hyperlipidemia, diabetes, hypertension, coronary heart disease, inflammatory bowel disease, other cancers and gastrointestinal

1
2
3
4
5 bleeding, the HR for the study group compared with the control group was 1.62 (95%
6
7
8 CI = 1.24–2.12; $p \leq 0.001$). This study further classified patients in the study group as
9
10 short-term and long-term ADT users based on the median duration of use. Compared
11
12
13 with the findings in the control group, the adjusted HRs for short-term ADT use in the
14
15
16 study group were 2.06 (95% CI = 1.53–2.76) (Table 3). Furthermore, sensitivity
17
18
19 analyses were displayed in Supplementary Table 1 and Supplementary Table 2.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

DISCUSSION

To the best of our knowledge, this is the first study to investigate the relationship between ADT use in men with PCa and the 3-year IDA risk in an East Asian population. This study only included patients with PCa in the relevant analyses to avoid the potential effects attributable to cancer. To mitigate selection bias, a propensity score matching strategy (by year of the entry date, age, monthly income, geographical location, residential urbanisation level and medical comorbidities) was used to facilitate comparisons with patients with PCa who did not receive ADT. In this population-based retrospective cohort study of patients with PCa, a significant increase in IDA risk was observed with men treated with ADT.

The findings were consistent with those of prior Western studies. For instance, in the United States, Strum et al. demonstrated that the haemoglobin levels of patients who received ADT for PCa declined from a mean of 149 g/L at baseline to 139, 132 and 131 g/L after 1, 2 and 3 months, respectively.³⁰ In Canada, Timilshina et al. performed an observational study of 250 patients with non-metastatic PCa and found that ADT was independently associated with a reduction of haemoglobin levels over 12 months.²³ Among 110 patients with PCa who received ADT, one cohort study reported a decline in haemoglobin levels from 14.8 g/dL at baseline to 12.9 g/dL at evaluation.²² Another study of 72 patients with non-metastatic PCa who received adjuvant radiotherapy plus

1
2
3
4
5 ADT reported that the haemoglobin level had significant declined after 2 years of
6
7
8 androgen suppression.³¹ All of these studies investigated the association of ADT with
9
10
11 haemoglobin levels, which is vital given that haemoglobin levels are important for
12
13
14 identifying anaemia. However, these clinical studies included small sample sizes and
15
16
17 featured a short duration.^{22 23 30 31} Separately, Hicks et al. performed a cohort study using
18
19
20 the United Kingdom Clinical Practice Research Database linked to the Hospital Episode
21
22
23 Statistics repository. Their findings revealed that patients with non-metastatic PCa who
24
25
26 received ADT had a nearly 3-fold greater risk of anaemia than non-users (HR = 2.90,
27
28
29 95% CI = 2.67–3.16).²⁴ However, this study was performed in a Western population,
30
31
32 and generalisation of its findings to other ethnic groups is not possible.

33
34
35 Our study examined the association between ADT and the 3-year IDA risk among
36
37
38 patients with PCa; however, the risk of anaemia caused by ADT use has remained
39
40
41 unclear. To date, increasing biological evidence has suggested a potential connection
42
43
44 between the receipt of ADT and subsequent IDA risk.³² A possible explanation for the
45
46
47 underlying mechanism may be that androgen significantly increases the levels of bone
48
49
50 marrow erythroid precursors and enhances the differentiation of bone marrow erythroid
51
52
53 stem cells to erythrocytes.^{21 33} Furthermore, testosterone might influence erythropoietin
54
55
56 secretion via peritubular fibroblast-like cells in the kidneys.³⁴ In animal experiments,
57
58
59 testosterone induced erythropoiesis-stimulating factor production in a bioassay using a
60

1
2
3
4
5 mouse model of polycythaemia vera.³⁵ Guo et al. found that sensitivity to erythropoietin
6
7
8 was induced and stress erythropoiesis was increased by testosterone, which upregulated
9
10
11 the expression of GATA binding protein 1 and GATA-dependent genes, in a study of
12
13
14 ageing male mice.³⁶ In addition, Bachman et al. observed that treatment with
15
16
17 testosterone in men was related to serum hepcidin suppression in a randomised double-
18
19
20 blind clinical study.¹⁹ Testosterone directly interacts with the bone morphogenetic
21
22
23 protein-Smad signalling pathway in hepatocytes and induces the downregulation of
24
25
26 hepcidin transcription.³⁷ Furthermore, these effects of androgen on hepatic hepcidin and
27
28
29 renal erythropoietin gene expression could increase systemic iron transport and
30
31
32 erythropoietin levels.³⁷

33
34
35 The strengths of our study included the identification of the association between
36
37
38 ADT use in patients with PCa and subsequent IDA risk using a large population-based
39
40
41 database. The Taiwan NHIRD provides high accessibility to medical service data and
42
43
44 adequate to detect IDA risk between the study and control groups. Moreover, the
45
46
47 sociodemographic characteristics and medical comorbidities of patients with PCa were
48
49
50 acknowledged as risk factors. Furthermore, we also used a propensity score matching
51
52
53 strategy to eliminate potential biases such as patients' demographics and comorbidities
54
55
56 between the study and control cohorts.

57
58
59 Several limitations of our retrospective study should be addressed. The study
60

1
2
3
4
5 lacked information on potentially important patient characteristics, such as body mass
6
7
8 index, smoking, nutrition and non-prescription medication use. Second, it is plausible
9
10
11 that the database did not include all patients with PCa and IDA in Taiwan because some
12
13
14 patients might have sought alternative medicines not recorded by the NHI programme
15
16
17 and some patients with mild or normocytic anaemia may not have immediately sought
18
19
20 medical treatment. Therefore, these patients might not have been captured via diagnosis
21
22
23 codes. Third, information on the family history of anaemia, cancer stage and grade for
24
25
26 the presence of metastases, chemotherapy, radiotherapy and surgery was unavailable
27
28
29 this study. Finally, this study lacked information regarding blood variables.
30
31
32 Nevertheless, this study identified patients with PCa who used ADT as cases and
33
34
35 patients with PCa patients who did not use ADT as controls. In general, physicians in
36
37
38 Taiwan perform complete blood counts for patients with PCa to identify suitable
39
40
41 treatments. Consequently, we considered that these factors may not have affected the
42
43
44 findings in this study.

45 46 47 **CONCLUSIONS**

48
49
50
51 In conclusion, the present study detected an increased IDA risk during a 3-year
52
53
54 follow-up period among patients with PCa who received ADT. Medical professionals
55
56
57 are recommended to be aware of the risk of anaemia following ADT. Clinicians and
58
59
60 pharmacists need to consider the possible risk of IDA among patients with PCa who

1
2
3
4
5 received ADT and assess the efficacy of preventative and treatment modalities for
6
7
8 anaemia.
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

Contributors

Conceptualisation: Fang-Jen Wu, Yih-Dih Cheng, I-Hsun Li and Li-Ting Kao;

Methodology: Fang-Jen Wu, Yih-Dih Cheng, Yi-Chun Lin, and Li-Ting Kao; Formal

analysis and investigation: Fang-Jen Wu and Wu-Chien Chien; Writing - original draft

preparation: Fang-Jen Wu, Jui-Hu Shih, Chin-Min Chuang and Yih-Dih Cheng

Ethics approval

This study was exempt from full revision by the Tri-Service General Hospital

Institutional Review Board (TSGHIRB No. 2-105-05-082) because the dataset used was

blinded to the public for research purposes.

Data sharing statement

No additional data available.

REFERENCES

1. Fitzmaurice C, Dicker D, Pain A, et al. The Global Burden of Cancer 2013. *JAMA oncology* 2015;1(4):505-27. doi: 10.1001/jamaoncol.2015.0735
2. Attard G, Parker C, Eeles RA, et al. Prostate cancer. *Lancet* 2016;387(10013):70-82. doi: 10.1016/S0140-6736(14)61947-4
3. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2018. *CA Cancer J Clin* 2018;68(1):7-30. doi: 10.3322/caac.21442
4. Shahinian VB, Kuo YF, Gilbert SM. Reimbursement policy and androgen-deprivation therapy for prostate cancer. *N Engl J Med* 2010;363(19):1822-32. doi: 10.1056/NEJMsa0910784
5. Bryant AK, D'Amico AV, Nguyen PL, et al. Three-month posttreatment prostate-specific antigen level as a biomarker of treatment response in patients with intermediate-risk or high-risk prostate cancer treated with androgen deprivation therapy and radiotherapy. *Cancer* 2018;124(14):2939-47. doi: 10.1002/cncr.31400
6. Harris WP, Mostaghel EA, Nelson PS, et al. Androgen deprivation therapy: progress in understanding mechanisms of resistance and optimizing androgen depletion. *Nat Clin Pract Urol* 2009;6(2):76-85. doi: 10.1038/ncpuro1296
7. Owen PJ, Daly RM, Livingston PM, et al. Lifestyle guidelines for managing adverse

- 1
2
3
4 effects on bone health and body composition in men treated with androgen
5
6
7 deprivation therapy for prostate cancer: an update. *Prostate Cancer Prostatic*
8
9
10 *Dis* 2017;20(2):137-45. doi: 10.1038/pcan.2016.69
11
12
13
14 8. Nguyen PL, Alibhai SM, Basaria S, et al. Adverse effects of androgen deprivation
15
16 therapy and strategies to mitigate them. *Eur Urol* 2015;67(5):825-36. doi:
17
18 10.1016/j.eururo.2014.07.010
19
20
21
22
23 9. Basaria S, Muller DC, Carducci MA, et al. Hyperglycemia and insulin resistance in
24
25 men with prostate carcinoma who receive androgen-deprivation therapy. *Cancer*
26
27 2006;106(3):581-8. doi: 10.1002/cncr.21642
28
29
30
31
32 10. Grossmann M, Hamilton EJ, Gilfillan C, et al. Bone and metabolic health in patients
33
34 with non-metastatic prostate cancer who are receiving androgen deprivation
35
36 therapy. *Med J Aust* 2011;194(6):301-6.
37
38
39
40
41 11. Sharifi N, Gulley JL, Dahut WL. Androgen deprivation therapy for prostate cancer.
42
43 *JAMA* 2005;294(2):238-44. doi: 10.1001/jama.294.2.238
44
45
46
47 12. Wu FJ, Sheu SY, Lin HC, et al. Increased Fall Risk in Patients Receiving Androgen
48
49 Deprivation Therapy for Prostate Cancer. *Urology* 2016;95:145-50. doi:
50
51 10.1016/j.urology.2016.05.058
52
53
54
55
56 13. Johnson-Wimbley TD, Graham DY. Diagnosis and management of iron deficiency
57
58 anemia in the 21st century. *Therap Adv Gastroenterol* 2011;4(3):177-84. doi:
59
60

1
2
3
4
5 10.1177/1756283X11398736
6
7

8 14. Caro JJ, Salas M, Ward A, et al. Anemia as an independent prognostic factor for
9
10 survival in patients with cancer: a systemic, quantitative review. *Cancer*
11
12 2001;91(12):2214-21.
13
14

15
16 15. Dicato M, Plawny L, Diederich M. Anemia in cancer. *Ann Oncol* 2010;21 Suppl
17
18 7:vii167-72. doi: 10.1093/annonc/mdq284
19
20

21
22 16. Dueregger A, Heidegger I, Ofer P, et al. The use of dietary supplements to alleviate
23
24 androgen deprivation therapy side effects during prostate cancer treatment.
25
26 *Nutrients* 2014;6(10):4491-519. doi: 10.3390/nu6104491
27
28

29
30 17. Lopez A, Cacoub P, Macdougall IC, et al. Iron deficiency anaemia. *Lancet*
31
32 2016;387(10021):907-16. doi: 10.1016/s0140-6736(15)60865-0
33
34
35

36
37 18. Ferrucci L, Maggio M, Bandinelli S, et al. Low testosterone levels and the risk of
38
39 anemia in older men and women. *Arch Intern Med* 2006;166(13):1380-8. doi:
40
41 10.1001/archinte.166.13.1380
42
43
44

45
46 19. Bachman E, Feng R, Travison T, et al. Testosterone suppresses hepcidin in men: a
47
48 potential mechanism for testosterone-induced erythrocytosis. *J Clin Endocrinol*
49
50 *Metab* 2010;95(10):4743-7. doi: 10.1210/jc.2010-0864
51
52
53

54
55 20. Latour C, Kautz L, Besson-Fournier C, et al. Testosterone perturbs systemic iron
56
57 balance through activation of epidermal growth factor receptor signaling in the
58
59
60

- 1
2
3
4
5 liver and repression of hepcidin. *Hepatology* 2014;59(2):683-94. doi:
6
7
8 10.1002/hep.26648
9
- 10
11 21. Moriyama Y, Fisher JW. Effects of testosterone and erythropoietin on erythroid
12
13 colony formation in human bone marrow cultures. *Blood* 1975;45(5):665-70.
14
15
- 16
17 22. D'Amico AV, Saegaert T, Chen MH, et al. Initial decline in hemoglobin during
18
19 neoadjuvant hormonal therapy predicts for early prostate specific antigen failure
20
21 following radiation and hormonal therapy for patients with intermediate and
22
23 high-risk prostate cancer. *Cancer* 2002;95(2):275-80. doi: 10.1002/cncr.10673
24
25
26
27
- 28
29 23. Timilshina N, Hussain S, Breunis H, et al. Predictors of hemoglobin decline in non-
30
31 metastatic prostate cancer patients on androgen deprivation therapy: a matched
32
33 cohort study. *Support Care Cancer* 2011;19(11):1815-21. doi: 10.1007/s00520-
34
35 010-1023-6
36
37
38
- 39
40 24. Hicks BM, Klil-Drori AJ, Yin H, et al. Androgen Deprivation Therapy and the Risk
41
42 of Anemia in Men with Prostate Cancer. *Epidemiology* 2017;28(5):712-18. doi:
43
44 10.1097/ede.0000000000000678
45
46
47
48
- 49
50 25. Rebbeck TR. Prostate Cancer Genetics: Variation by Race, Ethnicity, and
51
52 Geography. *Semin Radiat Oncol* 2017;27(1):3-10. doi:
53
54 10.1016/j.semradonc.2016.08.002
55
56
57
- 58
59 26. Taitt HE. Global Trends and Prostate Cancer: A Review of Incidence, Detection,
60

- 1
2
3
4
5 and Mortality as Influenced by Race, Ethnicity, and Geographic Location.
6
7
8 *American journal of men's health* 2018;12(6):1807-23. doi:
9
10 10.1177/1557988318798279
11
12
13
14 27. McLean E, Cogswell M, Egli I, et al. Worldwide prevalence of anaemia, WHO
15
16 Vitamin and Mineral Nutrition Information System, 1993-2005. *Public Health*
17
18 *Nutr* 2009;12(4):444-54. doi: 10.1017/s1368980008002401
19
20
21
22
23 28. Poon DMC, Chan T, Chan K, et al. Preliminary efficacy and tolerability of
24
25 chemohormonal therapy in metastatic hormone-naive prostate cancer: The first
26
27 real-life experience in Asia. *Asia Pac J Clin Oncol* 2018;14(4):347-52. doi:
28
29 10.1111/ajco.12874
30
31
32
33
34
35 29. Liu CY, Hung YT, Chuang YL, et al. Incorporating development stratification of
36
37 Taiwan townships into sampling design of large scale health interview survey. *J*
38
39 *Health Manag* 2006;4(1):1-22.
40
41
42
43
44 30. Strum SB, McDermed JE, Scholz MC, et al. Anaemia associated with androgen
45
46 deprivation in patients with prostate cancer receiving combined hormone
47
48 blockade. *Br J Urol* 1997;79(6):933-41. doi: 10.1046/j.1464-410x.1997.00234.x
49
50
51
52
53 31. Choo R, Chander S, Danjoux C, et al. How are hemoglobin levels affected by
54
55 androgen deprivation in non-metastatic prostate cancer patients? *The Canadian*
56
57 *journal of urology* 2005;12(1):2547-52.
58
59
60

- 1
2
3
4
5 32. Shahani S, Braga-Basaria M, Maggio M, et al. Androgens and erythropoiesis: past
6
7 and present. *J Endocrinol Invest* 2009;32(8):704-16. doi: 10.1007/bf03345745
8
9
- 10 33. Dai D, Han S, Li L, et al. Anemia is associated with poor outcomes of metastatic
11
12 castration-resistant prostate cancer, a systematic review and meta-analysis.
13
14 *American journal of translational research* 2018;10(12):3877-86.
15
16
17
18
19
- 20 34. Pan X, Suzuki N, Hirano I, et al. Isolation and characterization of renal
21
22 erythropoietin-producing cells from genetically produced anemia mice. *PLoS*
23
24 *One* 2011;6(10):e25839. doi: 10.1371/journal.pone.0025839
25
26
27
- 28 35. Mirand EA, Gordon AS, Wenig J. Mechanism of testosterone action in
29
30 erythropoiesis. *Nature* 1965;206(981):270-2. doi: 10.1038/206270a0
31
32
33
- 34 36. Guo W, Li M, Bhasin S. Testosterone supplementation improves anemia in aging
35
36 male mice. *J Gerontol A Biol Sci Med Sci* 2014;69(5):505-13. doi:
37
38 10.1093/gerona/glt127
39
40
41
42
- 43 37. Guo W, Bachman E, Li M, et al. Testosterone administration inhibits hepcidin
44
45 transcription and is associated with increased iron incorporation into red blood
46
47 cells. *Aging cell* 2013;12(2):280-91. doi: 10.1111/accel.12052
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 **Figure Legend**
5
6

7 **Figure 1** Kaplan–Meier curve of the risk of iron-deficiency anaemia (IDA) among
8
9
10 patients with prostate cancer during a 3-year follow-up period as stratified by the
11
12
13 previous use or non-use of androgen deprivation therapy (ADT).
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

Table 1 Demographic characteristics of patients with prostate cancer stratified by the previous use or non-use of androgen deprivation therapy (ADT) ($n=10,893$)

Characteristic	Patients not receiving ADT ($n=3,631$)		Patients receiving ADT ($n=7,262$)		<i>p</i> value
	Total no.	Column %	Total no.	Column %	
Age (years), mean (SD)	74.26 (8.81)		74.46 (8.81)		0.284
Urbanisation level					0.792
1 (most urbanised)	943	25.97	1,816	25.01	
2	871	23.99	1,734	23.88	
3	492	13.55	1,024	14.10	
4	492	13.55	1,011	13.92	
5 (least urbanised)	833	22.94	1,677	23.09	
Geographic region					0.454
Northern	1,876	51.67	3,676	50.62	
Central	850	23.41	1,688	23.24	
Southern	860	23.68	1,789	24.64	
Eastern	45	1.24	109	1.50	
Monthly income					0.385
NT\$0–15,840	2,426	66.81	4,826	66.46	
NT\$15,841–25,000	932	25.67	1,931	26.59	
\geq NT\$25,001	273	7.52	505	6.95	
Hypertension	2,047	56.38	4,099	56.44	0.946
Hyperlipidemia	718	19.77	1,308	18.01	0.026
Diabetes	792	21.81	1,511	20.81	0.226
Coronary heart disease	838	23.08	1,679	23.12	0.962
Inflammatory bowel disease	29	0.80	65	0.90	0.608
Gastrointestinal bleeding	122	3.36	265	3.65	0.442
Previous cancers	348	9.58	533	7.34	<0.001

Notes: The average exchange rate in 2015 was US\$1.00 \approx New Taiwan (NT)\$30.

Table 2 Crude and adjusted hazard ratios (HRs) for anaemia among patients with prostate cancer during a 3-year follow-up period stratified by the previous use or non-use of androgen deprivation therapy (ADT)

Presence of iron-deficiency anaemia	Total sample (<i>n</i> = 10,893)	Patients receiving ADT (<i>n</i> = 7,262)	Patients not receiving ADT (<i>n</i> = 3,631)
Three-year follow-up period			
Incidence rate per 100 person-years (95% CI)	1.45 (1.29-1.61)	1.66 (1.45-1.86)	1.01 (0.78-1.25)
Crude HR (95% CI)	--	1.64 (1.25-2.14)	1.00
Adjusted HR (95% CI) ^{a,b}	--	1.58 (1.21-2.07)	1.00
Adjusted HR (95% CI) ^{a,c}	--	1.62 (1.24-2.12)	1.00


Notes: ^a Using a Cox proportional regression with data censored if individuals died during the 3-year follow-up period; ^b Adjustments were made for patients' geographic location, monthly income, urbanisation level, age and incidence of hyperlipidaemia, diabetes, hypertension, coronary heart disease and inflammatory bowel disease. ^c Adjustments were made for patients' geographic location, monthly income, urbanisation level, age and incidence of hyperlipidaemia, diabetes, hypertension, coronary heart disease, inflammatory bowel disease, other cancers and gastrointestinal bleeding.

CI, confidence interval

Table 3 Crude and adjusted hazard ratios (HRs) for anaemia among patients with prostate cancer during a 3-year follow-up period stratified by the duration of androgen deprivation therapy (ADT)

Presence of iron-deficiency anaemia	Patients receiving short-term ADT (n =3,631) ^d	Patients receiving long-term ADT (n =3,634) ^d	Patients not receiving ADT (n = 3,631)
Three-year follow-up period			
Incidence rate per 100 person-years (95% CI)	2.24 (1.87-2.62)	1.25 (1.02-1.48)	1.01 (0.78-1.25)
Crude HR (95% CI)	2.19 (1.64-2.92)	1.25 (0.93-1.69)	1.00
Adjusted HR (95% CI) ^{a,b}	2.20 (1.64-2.94)	1.22 (0.90-1.65)	1.00
Adjusted HR (95% CI) ^{a,c}	2.06 (1.53-2.76)	1.23 (0.91-1.66)	1.00

Notes: ^a Using a Cox proportional regression with data censored if individuals died during the 3-year follow-up period; ^b Adjustments were made for patients’ geographic location, monthly income, urbanisation level, age and incidence of hyperlipidemia, diabetes, hypertension, coronary heart disease and inflammatory bowel disease. ^c Adjustments were made for patients’ geographic location, monthly income, urbanisation level, age and incidence of hyperlipidaemia, diabetes, hypertension, coronary heart disease, inflammatory bowel disease, other cancers and gastrointestinal bleeding. ^d We categorised the patients receiving ADT into two levels according to the median duration of ADT use (144 days).
 CI, confidence interval



99x72mm (300 x 300 DPI)

Supplementary Material

Supplementary Table 1 Crude and adjusted hazard ratios (HRs) for anaemia among patients with prostate cancer during a 3-year follow-up period stratified by the different androgen deprivation therapy (ADT) exposure definition

Presence of iron-deficiency anaemia	Current ADT users ($n = 109$) ^d	Past ADT users ($n = 7,153$) ^d	Patients not receiving ADT ($n = 3,631$)
Three-year follow-up period			
Incidence rate per 100 person-years (95% CI)	93.65 (76.07-111.24)	0.94 (0.78-1.09)	1.01 (0.78-1.25)
Crude HR (95% CI)	90.74 (68.88-119.55)	0.93 (0.69-1.24)	1.00
Adjusted HR (95% CI) ^{a,b}	89.64 (67.19-119.58)	0.91 (0.68-1.22)	1.00
Adjusted HR (95% CI) ^{a,c}	81.52 (60.58-109.69)	0.90 (0.67-1.20)	1.00

Notes: ^a Using a Cox proportional regression with data censored if individuals died during the 3-year follow-up period; ^b Adjustments were made for patients' geographic location, monthly income, urbanisation level, age and incidence of hyperlipidemia, diabetes, hypertension, coronary heart disease and inflammatory bowel disease. ^c Adjustments were made for patients' geographic location, monthly income, urbanisation level, age and incidence of hyperlipidaemia, diabetes, hypertension, coronary heart disease, inflammatory bowel disease, other cancers and gastrointestinal bleeding. ^d We categorised the patients receiving ADT into current users (patients who had received ADT prior to the outcome date within 1 month) and past users (other remaining users).

CI, confidence interval

Supplementary Table 2 Sensitivity analysis

Outcome variable	Outcome definition: cases with anaemia diagnosis		Outcome definition: cases with anaemia diagnosis and receiving iron supplementation	
	Patients receiving ADT	Patients not receiving ADT	Patients receiving ADT	Patients not receiving ADT
	(n = 7,262)	(n = 3,631)	(n = 7,262)	(n = 3,631)
Presence of iron-deficiency anaemia				
Three-year follow-up period				
Incidence rate per 100 person-years (95% CI)	1.66 (1.45-1.86)	1.01 (0.78-1.25)	1.66 (1.45-1.86)	0.08 (0.02-0.15)
Crude HR (95% CI)	1.64 (1.25-2.14)	1.00	19.56 (8.69-44.02)	1.00
Adjusted HR (95% CI) ^{a,b}	1.58 (1.21-2.07)	1.00	19.43 (8.63-43.76)	1.00
Adjusted HR (95% CI) ^{a,c}	1.62 (1.24-2.12)	1.00	19.14 (8.50-43.10)	1.00

Notes: ^a Using a Cox proportional regression with data censored if individuals died during the 3-year follow-up period; ^b Adjustments were made for patients' geographic location, monthly income, urbanisation level, age and incidence of hyperlipidaemia, diabetes, hypertension, coronary heart disease and inflammatory bowel disease. ^c Adjustments were made for patients' geographic location, monthly income, urbanisation level, age and incidence of hyperlipidaemia, diabetes, hypertension, coronary heart disease, inflammatory bowel disease, other cancers and gastrointestinal bleeding.

CI, confidence interval